

Correlates of HIV Infection Among Sexually Active Adults in Kenya: A National Population-Based Survey

Tom Oluoch^{*1}, Ibrahim Mohammed², Rebecca Bunnell¹, Reinhard Kaiser¹, Andrea A. Kim³, Anthony Gichangi¹, Mary Mwangi¹, Sufia Dadabhai⁴, Lawrence Marum³, Alloys Orago⁵ and Jonathan Mermin⁶ for the 2007 KAIS Study Group

¹US Centers for Disease Control and Prevention – Division of Global HIV/AIDS, Nairobi, Kenya

²National AIDS and STI Control Program, Ministry of Medical Services, Nairobi, Kenya

³US Centers for Disease Control and Prevention – Division of Global HIV/AIDS, Atlanta, Georgia, USA

⁴Johns Hopkins Bloomberg School of Public Health, Nairobi, Kenya

⁵National AIDS Control Council, Nairobi, Kenya

⁶US Centers for Disease Control and Prevention--Division of HIV/AIDS Prevention, National Center for HIV, Viral Hepatitis, STD, and TB Prevention, Atlanta, USA

Abstract: *Objective:* To identify factors associated with prevalent HIV in a national HIV survey in Kenya.

Methods: The Kenya AIDS Indicator Survey was a nationally representative population-based sero-survey that examined demographic and behavioral factors and serologic testing for HIV, HSV-2 and syphilis in adults aged 15-64 years. We analyzed questionnaire and blood testing data to identify significant correlates of HIV infection among sexually active adults.

Results: Of 10,957 eligible women and 8,883 men, we interviewed 10,239 (93%) women and 7,731 (87%) men. We collected blood specimens from 9,049 women and 6,804 men of which 6,447 women and 5,112 men were sexually active during the 12 months prior to the survey. HIV prevalence among sexually active adults was 7.4%. Factors independently associated with HIV among women were region (Nyanza vs Nairobi: adjusted OR [AOR] 1.6, 95%CI 1.1-2.3), number of lifetime sex partners (6-9 vs 0-1 partners: AOR 3.0, 95%CI 1.6-5.9), HSV-2 (AOR 6.5, 95%CI 4.9-8.8), marital status (widowed vs never married: AOR 2.7, 95%CI 1.5-4.8) and consistent condom use with last sex partner (AOR 2.3, 95%CI 1.6-3.4). Among men, correlates of HIV infection were 30-to-39-year-old age group (AOR 5.2, 95%CI 2.6-10.5), number of lifetime sex partners (10+ vs 0-1 partners, AOR 3.5, 95%CI 1.4-9.0), HSV-2 (AOR 4.7, 95%CI 3.2-6.8), syphilis (AOR 2.4, 95%CI 1.4-4.0), consistent condom use with last sex partner (AOR 2.1, 95% CI 1.5-3.1) and lack of circumcision (AOR 4.0, 95%CI 2.8 - 5.5).

Conclusion: Kenya's heterogeneous epidemic will require regional and gender-specific prevention approaches.

Keywords: AIDS Indicator Survey, HIV, HIV prevalence, Kenya, Nyanza, Population-based, Sexually transmitted infection.

INTRODUCTION

Lack of a comprehensive understanding of national HIV epidemics remains a major challenge for targeting effective HIV programs. Antenatal clinic (ANC) sentinel surveillance has been useful to document trends of generalized epidemics but lacks adequate sexual-behavioral and other associated risk factors for HIV infection, and is not generalizable to men and non-pregnant women [1]. Studies conducted in a number of sub-Saharan African countries showed that ANC data over-estimated prevalence in 15-19 year olds and under-estimated prevalence in older ANC clinic attendees [2, 3].

Hospital-based studies have also been conducted in low prevalence areas to understand risk factors [4], but the findings are not easily generalizable to populations living in countries with high HIV prevalence.

Concerns about the representativeness and accuracy of national HIV estimates derived from ANC surveillance have led to an increased demand for surveys and more data on the prevalence and distribution of HIV in the whole population [5]. In sub-Saharan Africa, more than 38 countries have conducted national population-based HIV surveys [6]. These surveys most often were restricted to reproductive ages, collected dried blood spots for biological tests [7], and avoided questions about sexual behavior and HIV status which are considered sensitive. Thus, laboratory and risk factor analyses have been limited. Programmatic data from voluntary counseling and testing (VCT) centers as well as HIV test results from prevention of mother-to-child

*Address correspondence to this author at the Division of Global HIV/AIDS, US Centers for Disease Control and Prevention, P.O. Box 606, 00621, Village Market, Nairobi, Kenya; Tel: +254 20 286 7000; Fax: +254 20 271 4745; E-mail: toluoch@ke.cdc.gov

transmission (PMTCT) programs have been used in some cases to estimate HIV prevalence and investigate associations with demographic, behavioral, and biological factors [8-10]. However, these are limited due to their lack of representativeness of the populations that seek VCT and PMTCT services [11]. For example, PMTCT services are only offered to pregnant women. Additionally, risk behavior, perception of risk behavior, or attitudes towards HIV/AIDS have been shown to influence uptake of VCT services [12].

Sexual transmission is the major route for HIV infection in sub-Saharan Africa [13,14]. In Kenya, sexual transmission contributes over 90% of all HIV infections [13]. The World Health Organization (WHO) categorizes Kenya as having a generalized HIV epidemic, defined as having an HIV prevalence of more than 1% in the adult general population and more than 5% in vulnerable adults such as sexually transmitted infection (STI) clinic patients. Kenya has wide variability in the distribution of HIV infection by geographic, demographic, sexual-behavioral, and biological factors. The Kenya Demographic and Health Survey (KDHS) conducted in 2003 included HIV serology but was limited in the scope and depth of sexual-behavioral and biological indicators collected, and included consenting adults aged 15-49 years [15]. In this era of expansion of prevention, care and treatment programs for HIV, additional variables of interest are needed to facilitate accurate interpretation of HIV prevalence data and associated risk factors [16].

We conducted the Kenya AIDS Indicator Survey (KAIS) in 2007 to provide nationally-representative and comprehensive data on demographic, behavioral, and biologic indicators of HIV/AIDS, beyond that of previous national HIV surveys in Kenya. In addition to providing national prevalence estimates for HIV and sexually transmitted infections, these data provided the opportunity to link HIV status with key demographic, behavioral, and biologic information to identify significant correlates associated with HIV infection in Kenya. We examined factors independently associated with the risk of HIV infection among individuals who were sexually active during the 12 months prior to the survey.

METHODS

Ethics Statement

Ethical approval was obtained from the Ethical Review Committee at the Kenya Medical Research Institute (KEMRI) and the Institutional Review Board at the Centers for Disease Control and Prevention (CDC). Respondents provided consent separately for the interviews and blood draws.

Study Setting

KAIS was conducted among a nationally representative sample of households selected from all the eight provinces in Kenya, covering both rural and urban areas. Various studies show that Nyanza province continues to have the highest HIV prevalence in Kenya [15,17,18]. Nairobi province is the capital city and has a cosmopolitan population with diverse ethnic groups and cultures. Nyanza on the other hand is predominantly inhabited by the *Luo* ethnic group.

Survey Design, Sampling and Participation

Briefly, we conducted a cross-sectional, stratified two-stage cluster survey designed to achieve a nationally representative sample and sufficient statistical power to provide prevalence estimates for each of the 8 provinces (Fig. 2). The first stage involved selecting 415 clusters out of 1,800 in the national sampling frame and the second stage involved the selection of households per cluster with equal probability of selection in the rural-urban strata. In order to reflect the population distribution, we sampled 294 and 121 clusters in rural and urban areas respectively. We collected data between August and December, 2007. Eligible participants were individuals aged 15-64 who were residents or household visitors the night prior to the survey team's visit. We administered household and individual questionnaires and collected venous blood from respondents. Key questions included respondent's HIV status, testing history, age at sexual debut, condom use, number of lifetime sex partners, alcohol use during last sexual intercourse, male circumcision status and mobility (which was defined as the number of separate occasions when the respondent traveled away from home/community and slept away in the previous 12 months) and access to cotrimoxazole and antiretroviral therapy (ART) for those with HIV infection. Partner-specific information on HIV status, testing history, disclosure, condom use and family planning preference were collected for up to four sex partners in the last year. With exception of condom use and family planning preferences, many of the indicators were unique to KAIS and were not asked in the KDHS. For population size estimates, we used census data from the Kenya National Bureau of Statistics, using the 2006 sampling frame based on the 1999 census. We calculated the corresponding 95% confidence intervals taking into account the sampling weights and study design. Full details of methods used in KAIS have been published elsewhere [18].

Laboratory Testing

Blood specimens were transported to Kenya's National Public Health Laboratory in Nairobi and tested for HIV, syphilis and HSV-2. CD4 cell count enumeration was conducted for HIV-infected participants. HIV testing was performed according to the national guidelines for HIV testing using a validated HIV testing algorithm for the country: the Vironostika HIV Uni-Form II antigen/antibody (BioMérieux Bv, Beseind, Netherlands) for screening and the Murex HIV antigen/antibody combination (Abbott/Murex-Biotech Ltd, Kent, UK) for confirmation in a serial testing algorithm. Specimens with discordant results were re-tested with the two assays. Polymerase Chain Reaction (PCR) testing was conducted on specimens that still had two discordant results after re-testing. For HSV-2 testing, the Kalon HSV Type 2- specific IgG EIA (Kalon Biologicals, Guildford, UK) was used; this was a recombinant type 2 antigen (gG2) modified to eliminate reactivity arising from HSV type 1 infection, at the same time retaining the natural antigenic characteristics of HSV-2. For syphilis infection, serum specimens were first screened using a *Treponema pallidum* particle agglutination assay (TPPA) (Serodia-TPPA, Fujirebio Diagnostics Inc, Tokyo, Japan) to detect previous exposure to syphilis antigens. All TPPA-positive specimens were then tested using a rapid plasma reagin (RPR) (Becton, Dickinson and Company, Franklin Lakes,

NJ, USA) test on undiluted (i.e. neat) serum specimen to test for reagenic antigens. Specimens with positive TPPA and RPR serologies were defined an active syphilis infection. For quality control of each test, all positive specimens and 5% of negative specimens were retested in a different laboratory using the same testing algorithms.

Data Analysis

Data from questionnaires were entered into a CSPro database (version 3.3, US Census Bureau, Washington DC, USA) by two different data clerks. All discrepancies between the two entries were resolved by a data manager during the data cleaning stage. To obtain nationally representative estimates, we calculated sampling weights for each individual and household based on selection probability and taking into account cluster-level non-participation. In addition, special weights were calculated for those who participated in the blood draw to take into account cluster-level non-participation.

For the purposes of this paper, we restricted analysis to a sub-set of questions from the individual questionnaire among participants who were sexually active in the last year before the survey. Sexually active individuals were those who self-reported having had sexual intercourse. Additionally, we conducted a sub-analysis to further investigate correlates of HIV infection that were specific to Nyanza province, the province with the highest HIV prevalence rate for men and women compared to all other provinces. We used the Rao-Scott chi-square test which allowed adjusting for the cluster survey design when testing for associations between categorical variables and HIV infection. Bivariate analysis was used to quantify the association between the demographic, behavioral and biological variables and HIV infection. We conducted multivariate logistic regression by constructing separate models for males and females to assess factors independently and significantly associated with HIV infection among sexually active persons. All variables were first included in the models and model selection was carried out using a backward elimination procedure. All variables that had a p-value of greater than 0.5 were removed from the models in the first step unless they were suspected to be confounders. Variables were then removed sequentially from the models starting with the one with the highest p-value until all variables had a p-value of less than 0.05. All confounders were retained in the model irrespective of the p-value. Odds ratios, adjusted odds ratios (AOR), and associated 95% confidence intervals were calculated based on pre-specified reference groups.

RESULTS

We collected information from 9,691 households. A total of 19,840 individuals were eligible, of which 17,970 (90%) consented to be interviewed, including 10,239 women (93%) and 7,731 men (87%) (Fig. 1). Among the eligible respondents, 9,049 (83%) women and 6,804 (77%) men consented for and provided a blood sample for HIV, HSV-2, and syphilis testing, a response rate of 88% of those interviewed, with similar uptakes rates among men and women. Among those consenting to blood draw 6,447 women (71%) and 5,112 men (75%) were sexually active in the last year. Overall, 57% of KAIS participants were

female, the median age was 30 years, 74% resided in rural areas, 70% were currently employed, 61% were married, and <1% of women and 6.6% of men drank alcohol before their last sexual encounter (Table 1). Among the laboratory-confirmed HIV positive respondents, 56% had never tested for HIV infection, 28% self-reported HIV negative and 16% self-reported HIV positive status. We found ethnicity to be co-linear with province and circumcision and choose to present differential risks with a focus on circumcision status and province rather than with a focus on ethnicity.

HIV Prevalence

Overall, HIV prevalence among all adults aged 15-64 years was 7.1% (95% CI, 6.5-7.7), representing an estimated 1.4 million people nationwide. Among sexually active adults aged 15-64 years old, 7.4% (95% CI, 6.7-8.2) were infected with HIV. Women were more likely to be infected (8.2%) than men (6.4%). Young women aged 15-24 years were nearly 5 times more likely to be infected (7.8%) than young men of the same age group (1.7%). HIV prevalence increased with age, with the highest prevalence (10.1%) between 30 and 39 years. Among 30-39 year olds, HIV prevalence was 11.0% among women compared to 9.0% among men. Prevalence among individuals aged 60-64 was 2.9%.

Although there was no difference in HIV prevalence between rural and urban areas, prevalence varied greatly across provinces (Table 1). Women had higher HIV prevalence than men in all provinces. In Nyanza Province 17.7% (95% CI, 14.9-20.5) of women and 14.1% (95% CI, 11.4-16.9) of men were infected. HIV prevalence was higher among married or cohabitating (7.4%, 95% CI 6.6-8.2), separated/ divorced (12.7%, 95% CI 9.1-16.2), and widowed individuals (21.3%, 95% CI 15.0-27.6) compared with never married individuals (4.3%, 95% CI 3.2-5.4). We observed no difference in HIV prevalence by wealth status for women and men; however, men who were currently employed had higher prevalence (6.8%, 95% CI 5.9-7.6) than those that were not employed (2.9%, 95% CI 1.0-4.8) (Table 1).

Behavioral Factors and HIV Infection

Among sexually active men, HIV prevalence was 18.1% (95% CI 14.8-21.4) among uncircumcised men and 4.5% (95% CI 3.8-5.2) among the circumcised men. HIV prevalence in uncircumcised men rose sharply between the 15-24 year age group (3.2%) and the 30-39 year age group (30.4%) and remained high in uncircumcised men through age 59.

Among participants who reported being sexually active in the last 12 months, consistent condom use with the last sex partner was associated with higher HIV prevalence compared to no condom use with the last sex partner (Table 1). However, a sub-analysis showed that HIV-infected persons who knew their sero-positive status were 4 times more likely to use a condom compared to those who self-reported an HIV negative status based on the result of their last test (53.1% vs 13.9%) or those who had never had an HIV test (11.2%).

HIV prevalence also varied with STI infection. Respondents who reported any STI in the last 12 months had

Table 1. HIV Prevalence by Demographic, Behavioral and Biologic Factors Among Sexually Active Adults Aged 15-64 Years in Kenya in 2007

Variable	Female			Male			Total		
	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)
Total	6447	536	8.2 (7.3 - 9.2)	5112	323	6.4 (5.6 - 7.2)	11559	859	7.4 (6.7 - 8.2)
Age									
15-24	1669	147	7.8 (6.2 - 9.4)	977	20	1.7 (0.8 - 2.5)	2646	167	5.6 (4.5 - 6.6)
25-29	1181	108	10.0 (7.9 - 12.1)	720	56	8.3 (6.0 - 10.7)	1901	164	9.4 (7.8 - 11.0)
30-39	1825	198	11.0 (9.1 - 12.9)	1354	115	9.0 (7.1 - 10.9)	3179	313	10.1 (8.7 - 11.6)
40-49	1156	63	5.2 (3.9 - 6.5)	1040	88	8.1 (6.3 - 10.0)	2196	151	6.6 (5.4 - 7.7)
50-59	520	18	3.5 (1.7 - 5.2)	740	37	5.2 (3.4 - 7.0)	1260	55	4.5 (3.1 - 5.9)
60-64	96	2	3.4 (0.0 - 8.1)	281	7	2.7 (0.6 - 4.9)	377	9	2.9 (0.5 - 5.3)
Gender									
Male	0	0	. (. -)	5112	323	6.4 (5.6 - 7.2)	5112	323	6.4 (5.6 - 7.2)
Female	6447	536	8.2 (7.3 - 9.2)	0	0	. (. -)	6447	536	8.2 (7.3 - 9.2)
Marital status									
never married/cohabitating	631	49	7.0 (4.6 - 9.4)	1017	30	2.6 (1.5 - 3.6)	1648	79	4.3 (3.2 - 5.4)
separated/divorced	299	54	17.1 (11.6 - 22.5)	198	14	5.8 (2.3 - 9.3)	497	68	12.7 (9.1 - 16.2)
widowed	145	37	21.0 (14.3 - 27.8)	31	8	22.4 (7.0 - 37.8)	176	45	21.3 (15.0 - 27.6)
currently married/cohabitating	5372	396	7.5 (6.6 - 8.4)	3866	271	7.3 (6.3 - 8.3)	9238	667	7.4 (6.6 - 8.2)
Widowhood*									
Yes	213	56	23.4 (17.3 - 29.6)	-	-	-	213	56	23.4 (17.3 - 29.6)
No	6234	480	7.7 (6.8 - 8.6)	-	-	-	6234	480	7.7 (6.8 - 8.6)
Level of education									
No primary	1076	55	6.2 (4.3 - 8.2)	478	16	4.4 (2.0 - 6.8)	1554	71	5.7 (4.3 - 7.1)
Incomplete primary	1846	202	10.6 (8.8 - 12.3)	1270	89	6.8 (5.1 - 8.6)	3116	291	9.0 (7.7 - 10.4)
Complete primary	1701	155	8.6 (7.2 - 10.1)	1347	94	6.9 (5.3 - 8.4)	3048	249	7.9 (6.7 - 9.0)
Secondary or higher	1824	124	6.4 (4.7 - 8.0)	2017	124	6.2 (5.0 - 7.4)	3841	248	6.3 (5.2 - 7.3)
Wealth Quintiles									
Lowest	1196	91	8.7 (6.7 - 10.7)	883	56	6.6 (4.5 - 8.7)	2079	147	7.8 (6.0 - 9.5)
Second	1148	100	9.0 (7.0 - 11.0)	898	58	7.1 (5.3 - 8.9)	2046	158	8.2 (6.7 - 9.7)
Middle	1225	94	7.6 (5.9 - 9.2)	919	56	6.1 (4.3 - 7.8)	2144	150	6.9 (5.6 - 8.3)
Fourth	1270	120	8.8 (6.8 - 10.8)	1023	59	5.4 (3.8 - 6.9)	2293	179	7.3 (5.9 - 8.7)
Highest	1608	131	7.4 (5.4 - 9.5)	1389	94	7.0 (5.1 - 8.8)	2997	225	7.2 (5.6 - 8.8)
Current employment									
Currently employed	4405	396	8.7 (7.6 - 9.8)	4620	308	6.8 (5.9 - 7.6)	9025	704	7.7 (6.9 - 8.5)
Not currently employed	2042	140	7.2 (5.7 - 8.6)	492	15	2.9 (1.0 - 4.8)	2534	155	6.4 (5.1 - 7.7)
Province									
Nairobi	696	68	10.6 (7.9 - 13.4)	623	44	8.4 (4.5 - 12.2)	1319	112	9.7 (6.8 - 12.5)
Central	887	35	4.0 (2.3 - 5.7)	746	24	3.9 (1.7 - 6.2)	1633	59	3.9 (2.2 - 5.7)
Coast	759	57	8.6 (6.4 - 10.7)	610	40	6.5 (4.2 - 8.8)	1369	97	7.6 (6.1 - 9.2)
Eastern	1048	62	6.1 (3.6 - 8.5)	751	20	2.7 (1.5 - 4.0)	1799	82	4.7 (3.0 - 6.4)
North-Eastern	305	3	1.2 (0.1 - 2.3)	225	4	1.1 (0.0 - 2.9)	530	7	1.2 (0.1 - 2.2)

(Table 1) contd.....

Variable	Female			Male			Total		
	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)
Nyanza	980	180	17.7 (14.9 - 20.5)	770	114	14.1 (11.4 - 16.9)	1750	294	16.1 (13.8 - 18.5)
Rift-Valley	933	74	6.5 (4.2 - 8.9)	734	49	5.9 (4.3 - 7.5)	1667	123	6.3 (4.6 - 8.0)
Western	839	57	6.2 (4.0 - 8.4)	653	28	4.4 (2.6 - 6.3)	1492	85	5.4 (3.7 - 7.1)
Urban/Rural									
Rural	4825	370	7.7 (6.8 - 8.7)	3763	220	6.1 (5.2 - 6.9)	8588	590	7.0 (6.3 - 7.8)
Urban	1622	166	9.7 (7.1 - 12.3)	1349	103	7.5 (5.4 - 9.6)	2971	269	8.8 (6.8 - 10.8)
Circumcision									
Yes				4418	189	4.5 (3.8 - 5.2)	4418	189	4.5 (3.8 - 5.2)
No				680	132	18.1 (14.8 - 21.4)	680	132	18.1 (14.8 - 21.4)
Mobility**									
0	3348	251	7.5 (6.3 - 8.7)	2735	158	5.8 (4.8 - 6.8)	6083	409	6.7 (5.8 - 7.7)
1	1911	168	8.4 (6.9 - 9.8)	1005	74	7.3 (5.5 - 9.1)	2916	242	8.0 (6.9 - 9.2)
2	1188	117	10.0 (8.1 - 12.0)	1372	91	7.0 (5.5 - 8.6)	2560	208	8.4 (7.1 - 9.8)
Age at first sex									
<10	10	1	8.6 (0.0 - 25.1)	44	0	. (-)	54	1	1.9 (0.0 - 5.8)
10-14	712	94	12.6 (9.6 - 15.7)	853	60	7.0 (5.1 - 8.8)	1565	154	9.6 (7.8 - 11.4)
15-19	3723	334	8.6 (7.4 - 9.9)	2886	196	6.8 (5.8 - 7.8)	6609	530	7.9 (7.0 - 8.7)
20-24	979	58	5.6 (3.9 - 7.3)	816	52	6.2 (4.4 - 8.0)	1795	110	5.9 (4.6 - 7.1)
25+	119	1	0.4 (0.0 - 1.3)	239	11	5.0 (2.1 - 7.9)	358	12	3.4 (1.5 - 5.4)
Condom use first sex									
Yes	427	35	6.8 (4.3 - 9.4)	280	6	1.7 (0.3 - 3.1)	707	41	4.9 (3.2 - 6.6)
No	1202	105	7.9 (6.1 - 9.6)	665	11	1.4 (0.5 - 2.3)	1867	116	5.5 (4.4 - 6.7)
Sex partners in the last 12 months									
0	0	0	. (-)	0	0	. (-)	0	0	. (-)
1	6268	505	8.0 (7.1 - 9.0)	4241	246	6.0 (5.2 - 6.8)	10509	751	7.2 (6.5 - 7.9)
2	136	24	15.5 (9.1 - 21.8)	695	55	7.5 (5.2 - 9.9)	831	79	8.9 (6.7 - 11.1)
3+	14	3	19.5 (0.0 - 42.2)	145	20	13.7 (7.3 - 20.1)	159	23	14.2 (8.0 - 20.4)
Lifetime sex partners									
0	0	0	. (-)	0	0	. (-)	0	0	. (-)
1	2579	81	3.7 (2.7 - 4.6)	554	7	1.5 (0.3 - 2.6)	3133	88	3.3 (2.5 - 4.1)
2-3	2905	291	8.9 (7.6 - 10.2)	1479	62	4.1 (2.8 - 5.3)	4384	353	7.3 (6.3 - 8.3)
4-5	654	105	16.2 (12.6 - 19.7)	1040	71	5.9 (4.2 - 7.6)	1694	176	10.0 (8.2 - 11.7)
6-9	124	26	20.1 (12.2 - 28.0)	719	46	7.7 (5.4 - 10.1)	843	72	9.7 (7.1 - 12.3)
10+	88	12	16.4 (5.1 - 27.6)	981	91	9.3 (7.1 - 11.4)	1069	103	9.8 (7.5 - 12.2)
Condom use with last sexual partner									
Yes	568	95	17.8 (14.1 - 21.6)	873	68	8.2 (6.1 - 10.2)	1441	163	12.0 (9.7 - 14.3)
No	5879	441	7.3 (6.4 - 8.2)	4239	255	6.0 (5.2 - 6.9)	10118	696	6.8 (6.1 - 7.5)

(Table 1) contd.....

Variable	Female			Male			Total		
	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)	Total (N)	Total - Positive (n)	Total - Positive % (95% CI)
Alcohol use at last sex									
Yes	934	89	8.9 (6.6 - 11.1)	560	42	7.5 (4.8 - 10.2)	1494	131	8.3 (6.6 - 10.1)
No	5513	447	8.1 (7.1 - 9.1)	4552	281	6.3 (5.4 - 7.1)	10065	728	7.3 (6.5 - 8.1)
HSV-2 infection									
HSV-2 negative	3592	79	2.2 (1.6 - 2.7)	3530	83	2.4 (1.7 - 3.0)	7122	162	2.3 (1.8 - 2.7)
HSV-2 positive	2793	455	15.8 (14.0 - 17.5)	1546	240	15.3 (13.3 - 17.3)	4339	695	15.6 (14.1 - 17.1)
Active Syphilis infection***									
Syphilis negative	6272	521	8.2 (7.3 - 9.2)	4962	301	6.2 (5.4 - 6.9)	11234	822	7.3 (6.6 - 8.0)
Syphilis positive	106	13	11.8 (5.1 - 18.5)	107	20	18.6 (9.0 - 28.2)	213	33	15.3 (9.3 - 21.4)
STI									
Yes	154	28	17.0 (9.3 - 24.6)	135	17	15.9 (7.2 - 24.7)	289	45	16.5 (10.5 - 22.6)
No	5708	479	8.1 (7.2 - 9.1)	4670	297	6.4 (5.6 - 7.2)	10378	776	7.4 (6.6 - 8.1)

Widowhood*: Widowhood variable was defined as previous marriage that ended due to death of a spouse. Some previously widowed women had remarried when the survey was conducted.

Mobility**: Mobility was defined as the number of separate occasions the respondent traveled away from home or community and slept away in the last 12 months.

Active Syphilis infection***: Active syphilis infection was defined as positive RPR and TPPA serologies.

higher prevalence of HIV (16.5%) compared to those not reporting an STI (7.4%), and 44.7% of women and 32.0% of men were infected with HSV-2. HIV prevalence among those co-infected with HSV-2 was 15.6% compared to 2.3% among those not infected with HSV-2. Syphilis prevalence was 1.9% (1.6% and 2.3% among women and men, respectively). HIV prevalence was significantly higher among those infected with syphilis (16.5%) compared to those without syphilis infection (7.4%).

Factors Independently Associated with HIV

Women

In multivariate analyses, independent correlates of HIV infection among sexually active women were geographical area of residence (Nyanza Province vs Nairobi: adjusted OR [AOR] 1.6, 95% CI 1.1-2.3), higher number of lifetime sex partners (6-9 vs 0-1 partners: AOR 3.0, 95% CI 1.6-5.9), HSV-2 infection (AOR 6.5, 95% CI 4.9-8.8), marital status (separated/divorced vs never married: AOR 2.2, 95% CI 1.2-3.9; widowed vs never married: AOR 2.7, 95% CI 1.5-4.8) and consistent condom use with the last sex partner (AOR 2.3, 95% CI 1.6-3.4) (Table 2).

Men

Among men, correlates of HIV infection were age group (30-39 years vs 15-24 years: AOR 5.2, 95% CI 2.6-10.5), number of lifetime sex partners (≥ 10 partners vs 0-1, AOR 3.5, 95% CI 1.4-9.0), HSV-2 infection (AOR 4.7 95% CI 3.2-6.8), syphilis infection (AOR 2.4, 95% CI 1.4-4.0), consistent condom use with the last sex partner (AOR 2.1, 95% CI 1.5-3.1) and lack of male circumcision (AOR 4.0, 95% CI 2.8 - 5.5) (Table 2).

Factors Associated with HIV in Nyanza Province

Prevalence among sexually active adults in Nyanza province was highest in the 25-to-29-year-old age group for women (23.4%) and in the 30-to-39-year old age group for men (25.9%). Of separated/divorced individuals, 42.2% were HIV-infected; uncircumcised men had a prevalence of 20.8% compared to the 6.8% among those circumcised. In multivariate analysis, factors independently associated with HIV prevalence among recently sexually active women were age group 25-29 years vs 15-24 years (AOR 1.3, 95% CI 0.8-2.3); current marital status (separated/divorced vs never married: AOR 5.7, 95% CI 1.5-20.9; widowed vs never married: AOR 4.4, 95% CI 1.0-19.5), HSV-2 infection (AOR 6.4, 95% CI 3.6-11.4), higher number of lifetime sex partners (6-9 vs 0-1: AOR 3.2, 95% CI 1.2-9.0), consistent condom use with the last sex partner (AOR 3.2, 95% CI 1.9-5.5), mobility (2 days vs 0 days: AOR 1.4, 95% CI 0.9-2.3). Factors associated with HIV infection among recently sexually active men were age group 30-39 years vs 15-24 years (AOR 19.1, 95% CI 5.9-61.6); education (incomplete primary vs no primary: AOR 1.4, 95% CI 0.5-4.0), HSV-2 infection (AOR 4.6, 95% CI 2.2-9.4), consistent condom use with the last sex partner (AOR 3.7, 95% CI 2.0-7.1), and lack of male circumcision among men (AOR 2.8, 95% CI 1.7-4.8).

DISCUSSION

In 2007, an estimated 7.4% of sexually active Kenyan adults aged 15-64 years were infected with HIV. Correlates of HIV infection among women and men were age, number of lifetime sex partners, residence in Nyanza province, HSV-2




Fig. (1). Study Profile.

infection, consistent condom use with the last sex partner and lack of circumcision among men. The strongest independent predictors for HIV infection for both women and men were HSV-2 co-infection and higher number of lifetime sex partners.

HIV prevalence was highest in Nyanza Province, where 16.9% of the sexually active adults were HIV-infected. In a sub-analysis for Nyanza Province, we found that age, HSV-2 infection, multiple lifetime sex partners, consistent condom use with the last sex partner and lack of male circumcision were independently associated with HIV. Many of these factors are similar to predictors of HIV infection found at the national level and are consistent with findings from other studies [1,8,16,19].

In Kenya, the adjusted odds of having HIV among sexually active persons with HSV-2 infection were 5-6-fold higher than those uninfected with HSV-2. However, awareness of HSV-2 is very low, even among health care providers, despite the high prevalence of HSV-2 and the

potential role of HSV-2 in driving the HIV epidemic [20,21]. HIV-induced immune compromise can result in frequent and persistent HSV disease, while poorly managed HSV replication may influence HIV transmission [22]. Researchers estimate that in settings with high HSV-2 prevalence, such as Nyanza province, HSV-2 infection could contribute to the risk of HIV-infection in more than one in four new cases of HIV [19,21]. Unfortunately, randomized controlled trials that have examined daily acyclovir treatment of HSV-2 among persons with HIV co-infection, or acyclovir prophylaxis among persons without HIV have not demonstrated a protective effect [23].




Fig. (2). Map of Kenya, with provincial HIV prevalence among all adults aged 15-64 years.

The results from KAIS show a non-statistically significant increase in overall HIV prevalence from KDHS 2003 (7.4% vs 6.7%) among those aged 15-49. The potential increase in HIV prevalence may be in part due to the survival effects of antiretroviral therapy. It may also indicate increasing incidence and a need to improve and expand HIV prevention programs throughout Kenya, and particularly in Nyanza Province. Appropriate messages on delaying sexual debut, knowledge of HIV status, male circumcision, consistent and correct use of condom with partner of unknown HIV status or known discordant HIV status, and reduction of number of sex partners should be reinforced [8, 24, 25]. Our study showed that widowhood and divorce were significantly associated with higher HIV prevalence. This corroborates findings from other sub-Saharan African countries which show that women who encounter marital disruption through divorce or widowhood were more likely to be HIV infected [26]. On the other hand, divorce is more common among HIV-infected women, particularly those in HIV discordant unions [27].

Ethnicity and province influence both the distribution of circumcision practice and HIV prevalence. Several studies have showed that male circumcision reduces the risk of HIV acquisition among men [28]. High prevalence in Nyanza

Table 2. Factors Independently Associated with HIV Infection Among Sexually Active Adults in Kenya in 2007

Variable	Females	Males
	Adjusted Odds Ratio (95% Confidence Interval)	Adjusted Odds Ratio (95% Confidence Interval)
Province		
Nairobi	ref	-
Central	0.4(0.3 - 0.8)	-
Coast	0.8(0.5 - 1.3)	-
Eastern	0.7(0.4 - 1.1)	-
North-Eastern	0.4(0.1 - 1.2)	-
Nyanza	1.6(1.1 - 2.3)	-
Rift-Valley	0.7(0.4 - 1.1)	-
Western	0.6(0.3 - 0.9)	-
Age		
15-24	ref	ref
25-29	1.2(0.9 - 1.7)	4.3(2.1 - 9.1)
30-39	1.1(0.8 - 1.5)	5.2(2.6 - 10.5)
40-49	0.4(0.3 - 0.6)	3.1(1.4 - 6.6)
50-59	0.3(0.1 - 0.5)	1.9(0.8 - 4.4)
60-64	0.6(0.1 - 2.3)	1.2(0.4 - 3.6)
Marital status		
Never married/cohabitating	ref	-
Currently married/cohabitating	1.5(0.9 - 2.4)	-
Separated/divorced	2.2(1.2 - 3.9)	-
Widowed	2.7(1.5 - 4.8)	-
HSV-2 infection		
HSV-2 negative	ref	ref
HSV-2 positive	6.5(4.9 - 8.8)	4.7(3.2 - 6.8)
Circumcision		
Yes	-	Ref
No	-	4.0(2.8 - 5.5)
Lifetime sexual partners		
0	ref	ref
10+	2.4(1.0 - 5.8)	3.5(1.4 - 9.0)
2-3	1.5(1.1 - 2.1)	2.0(0.8 - 4.8)
4-5	2.3(1.6 - 3.5)	2.2(0.8 - 5.8)
6-9	3.0(1.6 - 5.9)	2.6(1.0 - 6.9)
Consistent condom use with last sex partner		
No	ref	ref
Yes	2.3(1.6 - 3.4)	2.1(1.5 - 3.1)
Age at first sex		
<10	-	ref
10-14	-	1.3(0.6 - 2.9)
15-19	-	1.1(0.5 - 2.4)
20-24	-	1.2(0.6 - 2.6)
<10	-	0.0(0.0 - 0.0)
Active Syphilis infection*		
Syphilis negative	-	ref
Syphilis positive	-	2.4(1.4 - 4.0)

*Active Syphilis infection: Active syphilis infection was defined as positive RPR and TPPA serologies.

province could be attributed to low male circumcision rates. Among the general population, 85% of men were circumcised nationally while 48.2% were circumcised in Nyanza Province. In addition, cultural practices such as widow inheritance practiced among the Luo community (the predominant ethnic group in Nyanza) may be a factor [28, 29]. Widow inheritance is a traditional practice in which a designated man takes social and economic responsibility over a woman following the death of her husband [30]. A study by Agot *et al.* [30] in Nyanza province showed that inherited widows were more likely to have HIV infection compared to those not inherited. Luo women are believed to acquire contagious cultural impurity following the death of a husband. Often, a professional “cleanser” is hired who performs sexual rituals to cleanse the widow. If the spouse of the deceased is HIV-infected, the cleanser acts as a bridge for HIV transmission to other widows hence putting widowed women at a high risk [31]. The Kenya Ministries of Health published The National Guidance for Voluntary Medical Male Circumcision (VMMC) in January 2008 and in November 2008 launched the VMMC program focusing on Nyanza province and other traditionally non-circumcising communities with the aim of reducing new HIV infections and other STIs. The guidance provides a broad policy framework for the integration of VMMC into existing HIV prevention programs.

Respondents reporting ever having used condoms were more likely to have HIV infection. Condom use reduces risk of HIV acquisition and transmission [32] and HIV-infected persons who are aware of their HIV infection are more likely to use condoms [33, 34]. KAIS showed a 4-fold increase in condom use among those who knew their HIV-positive status. Though the association between condom use and HIV infection was not expected, the finding may reflect the success of positive prevention interventions and condom promotion efforts to increase condom use by people living with HIV. These data highlight the need for future HIV surveys to collect more detailed partner-specific information on condom use and knowledge of HIV status of participants and their partners which could help interpret the complex associations between condom use, sexual behavior, and HIV. The application of a laboratory assay that can accurately distinguish recent from established infection could also suggest the temporality of any associations between HIV infection and condom use, and more accurately highlight areas for programmatic focus.

Our study was limited by several factors. About 20% of eligible residents were either not present or declined to participate in the interview and blood draw. Although we do not expect that there is significant participation bias, we were not able to conduct these analyses; however, appropriate weighting was applied to adjust for non-response. These results cannot be generalized to all Kenyans, but only to those that reported recent sexual activity. Additionally, key sexual behavior indicators were based on self reported data. Though KAIS interviewers were trained on asking sensitive questions around sexual behavior and ensuring respondent confidentiality, there is a possibility that these questions were not accurately answered. We did not ask how long after circumcision the men engaged in sexual intercourse. The cross-sectional design of the study limited our interpretation of the temporality of association between the factors

examined and HIV infection. The survey also did not ask questions on men having sex with men or injecting drug use activities that are practiced in Kenya and may contribute to new HIV infections [13]. We did not include children due to the relatively low HIV prevalence among this group.

HIV remains a major public health challenge in Kenya. Although various prevention, care and treatment programs have been initiated and expanded in Kenya, evidence based prevention efforts that target known behavioral and biologic factors such as reduction of sex partners, condom use, delayed sexual debut and male circumcision should be enhanced. The wide regional variation in HIV prevalence reinforces the need for targeted prevention interventions focusing on provinces with high infection rates, while at the same time addressing the key behavioral factors that are associated with the risk of HIV infection nationally.

ACKNOWLEDGEMENTS

Special thanks to Professor George Rutherford of the University of California, San Francisco, for his useful comments and assistance in editing this paper

Supported by the U.S. President’s Emergency Plan for AIDS Relief, through CDC/HHS and USAID.

CONFLICTS OF INTEREST

None declared.

DISCLAIMER

The findings and conclusions in this paper are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

REFERENCES

- [1] Kiptoo M, Mpoke S, Nganga Z, Mueke J, Okoth F, Songok E. Survey on prevalence and risk factors on HIV-1 among pregnant women in North-Rift, Kenya: a hospital based cross-sectional study conducted between 2005 and 2006. *BMC Int Health Hum Rights* 2009; 9: 10.
- [2] Musinguzi J, Kirungi W, Opio A, *et al.* Comparison of HIV Prevalence Estimates From Sentinel Surveillance and a National Population-Based Survey in Uganda, 2004–2005. *J Acquir Immune Defic Syndr* 2009; 51(1): 78–84.
- [3] Montana L, Mishra V, Hong R. Comparison of HIV prevalence estimates from antenatal care surveillance and population-based surveys in sub-Saharan Africa. *Sex Transm Infect* 2008; 84(Suppl 1): i78–84.
- [4] Croce F, Fedeli P, Dahoma M, *et al.* Risk factors for HIV/AIDS in a low HIV prevalence site of sub-Saharan Africa. *Trop Med Int Health* 2007; 12(9): 1011–7.
- [5] UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Guidelines for measuring national HIV prevalence in population-based surveys. 2005.
- [6] MEASURE DHS. HIV/AIDS Survey Indicators Database. MEASURE DHS Tools 2009 [cited: 2009 Oct 15]; Available from: URL: <http://www.measuredhs.com/hivdata/surveys/start.cfm>
- [7] Garcia Calleja JM, Gouws E, Ghys PD. National population based HIV prevalence surveys in sub-Saharan Africa: results and implications for HIV and AIDS estimates. *Sex Transm Infect* 2006; 82 (Suppl III): iii64–70.
- [8] Landman KZ, Ostermann J, Crump JA, *et al.* Gender differences in the risk of HIV infection among persons reporting abstinence, monogamy, and multiple sexual partners in northern Tanzania. *PLoS One* 2008; 3(8): e3075.
- [9] Fabiani M, Cawthorne A, Nattabi B, Ayella EO, Ogwang M, Declich S. Investigating factors associated with uptake of HIV voluntary counselling and testing among pregnant women living in North Uganda. *AIDS Care* 2007; 19(6): 733–9.

- [10] Baryarama F, Bunnell RE, Montana L, *et al.* HIV prevalence in voluntary counseling and testing centers compared with national HIV serosurvey data in Uganda. *J Acquir Immune Defic Syndr* 2008; 49(2): 183-9.
- [11] Baryarama F, Bunnell RE, Ransom RL, *et al.* Using HIV voluntary counseling and testing data for monitoring the Uganda HIV epidemic, 1992-2000. *J Acquir Immune Defic Syndr* 2004; 37(1): 1180-6.
- [12] Bradley H, Tsui A, Kidanu A, Gillespie D. Client characteristics and HIV risk associated with repeat HIV testing among women in Ethiopia. *AIDS Behav* 2011; 15(4): 725-33.
- [13] Kenya National AIDS Control Council, UNAIDS, World Bank, Global AIDS M&E Team. Kenya - HIV Response and Modes of Transmission Analysis - Final Report. 2009.
- [14] Potterat JJ. AIDS epidemiology in Africa: a changing of the guard. *Int J STD AIDS* 2009; 20(12): 812-5.
- [15] Central Bureau of Statistics (CBS) [Kenya], Ministry of Health (MOH) [Kenya], ORC Macro. Kenya Demographic and Health Survey 2003-2004.
- [16] Chen L, Jha P, Stirling B, *et al.* Sexual risk factors for HIV infection in early and advanced HIV epidemics in sub-Saharan Africa: systematic overview of 68 epidemiological studies. *PLoS One* 2007; 2(10): e1001.
- [17] Central Bureau of Statistics (CBS) [Kenya], Ministry of Health (MOH) [Kenya], ORC Macro. Kenya Demographic and Health Survey 2008-09. 2010.
- [18] National AIDS and STI Control Program (NASCOP) K. 2007 Kenya AIDS Indicator Survey: Final Report. 2009.
- [19] Amornkul PN, Vandenhoudt H, Nasokho P, *et al.* HIV prevalence and associated risk factors among individuals aged 13-34 years in Rural Western Kenya. *PLoS One* 2009; 4(7): e6470.
- [20] Abu-Raddad LJ, Magaret AS, Celum C, *et al.* Genital herpes has played a more important role than any other sexually transmitted infection in driving HIV prevalence in Africa. *PLoS One* 2008; 3(5): e2230.
- [21] Freeman EE, Weiss HA, Glynn JR, Cross PL, Whitworth JA, Hayes RJ. Herpes simplex virus 2 infection increases HIV acquisition in men and women: systematic review and meta-analysis of longitudinal studies. *AIDS* 2006; 20(1): 73-83.
- [22] Ramaswamy M, Geretti AM. Interactions and management issues in HSV and HIV coinfection. *Expert Rev Anti Infect Ther* 2007; 5(2): 231-43.
- [23] Celum C, Wald A, Lingappa JR, *et al.* Acyclovir and transmission of HIV-1 from persons infected with HIV-1 and HSV-2. *N Engl J Med* 2010; 362(5): 427-39.
- [24] Stoneburner RL, Low-Beer D. Sexual partner reductions explain human immunodeficiency virus declines in Uganda: comparative analyses of HIV and behavioural data in Uganda, Kenya, Malawi, and Zambia. *Int J Epidemiol* 2004; 33(3): 624.
- [25] Williams BG, Lloyd-Smith JO, Gouws E, *et al.* The potential impact of male circumcision on HIV in Sub-Saharan Africa. *PLoS Med* 2006; 3(7): e262.
- [26] Boileau C, Clark S, Bignami-Van AS, *et al.* Sexual and marital trajectories and HIV infection among ever-married women in rural Malawi. *Sex Transm Infect* 2009; 85 (Suppl 1): i27-33.
- [27] Porter L, Hao L, Bishai D, *et al.* HIV status and union dissolution in sub-Saharan Africa: the case of Rakai, Uganda. *Demography* 2004; 41(3): 465-82.
- [28] Luginaah I, Elkins D, Maticka-Tyndale E, Landry T, Mathui M. Challenges of a pandemic: HIV/AIDS-related problems affecting Kenyan widows. *Soc Sci Med* 2005; 60(6): 1219-28.
- [29] Shiino W. Death and Rituals Among The Luo in South Nyanza. *Afr Study Monogr* 1997; 18: 213-28.
- [30] Agot KE, Vander SA, Tracy M, *et al.* Widow inheritance and HIV prevalence in Bondo District, Kenya: baseline results from a prospective cohort study. *PLoS One* 2010; 5(11): e14028.
- [31] Ambasa-Shisanya CR. Widowhood in the era of HIV/AIDS: a case study of Slaya District, Kenya. *SAHARA J* 2007; 4(2): 606-15.
- [32] Pinkerton SD, Galletly CL. Reducing HIV transmission risk by increasing serostatus disclosure: a mathematical modeling analysis. *AIDS Behav* 2007; 11(5): 698-705.
- [33] Crepaz N, Lyles CM, Wolitski RJ, *et al.* Do prevention interventions reduce HIV risk behaviours among people living with HIV? A meta-analytic review of controlled trials. *AIDS* 2006; 20(2): 143-57.
- [34] Bunnell R, Opio A, Musinguzi J, *et al.* HIV transmission risk behavior among HIV-infected adults in Uganda: results of a nationally representative survey. *AIDS* 2008; 22(5): 617-24.

Received: May 23, 2011

Revised: August 25, 2011

Accepted: September 26, 2011

© Oluoch *et al.*; Licensee Bentham Open.This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.