

On the Approximate Solution of Non-Linear Stochastic Diffusion Equation Using Symbolic WHEP

Noha A. Al-Mulla*

The University Dammam, Faculty of Science, Mathematics Department, Dammam, Saudi Arabia

Abstract: In this paper, a stochastic perturbed nonlinear diffusion equation is studied under a stochastic nonlinear non-homogeneity. The Pickard approximation method is used to introduce a reference first order approximate solution. Under different correction levels, the WHEP technique is used to obtain approximate solutions. Using Mathematica-5, the solution algorithm is operated and several comparisons among correction levels together with error curves have been demonstrated. The method of solution is illustrated through case studies and figures.

Keywords: Stochastic nonlinear diffusion equation, Eigenfunction expansion, Pickard approximation, WHEP technique.

1. INTRODUCTION

The study of random solutions of partial differential equations was initiated by Kampe de Fariet in 1955 [1]. In his valuable survey on the theory of random equations, Bharucha-Reid showed how a stochastic heat equation of Cauchy type can be solved using the stochastic integrals theory [2]. In 1973, Lo Dato V. [3] considered the stochastic velocity field and the Navier-Stokes equation and discussed the mathematical problems associated with it. Becus A. Georges [4] introduced a general solution for the heat conduction problem with a random source term and random initial and boundary conditions. Many authors investigated the stochastic diffusion equation under different views, see [5-11].

El-Tawil M. used the Wiener-Hermite expansion together with perturbation theory (WHEP technique) to solve a perturbed nonlinear stochastic diffusion equation [12]. The technique has been then developed to be applied on non-perturbed differential equations using the homotopy perturbation method and is called Homotopy WHEP [13].

In this paper, the diffusion equations with linear and nonlinear losses and stochastic nonhomogeneity are solved using WHEP technique under different correction levels. The main goal of the paper is to show that the correction levels are true corrections for the solution where the errors are decreased whenever the corrections are updated.

Two techniques are used to obtain the ensemble average, covariance and variance of the solution process, mainly the WHEP technique and the Pickard approximation which is used to get a reference solution. In section 2, the linear case is solved. The nonlinear case is analysed in section 3 using Pickard approximation to get a reference solution. The WHEP technique is processed in section 4 using different correction levels where the results of the execution of the

algorithms and comments on figures are included. Comparisons are concluded in section 5.

2. THE DIFFUSION EQUATION UNDER LINEAR LOSSES

Considering the following stochastic diffusion equation with $f(t, x; \omega)$ as random nonhomogeneity

$$\frac{\partial u(t, x; \omega)}{\partial t} = \frac{\partial^2 u}{\partial x^2} - \lambda u + f(t, x; \omega); (t, x) \in (0, \infty) \times (0, L), \quad (1)$$

$$u(t, 0) = 0, u(t, L) = 0 \text{ and } u(0, x) = g(x),$$

where ω is a random outcome of a triple probability space (Ω, χ, P) in which Ω is a sample space, χ is σ -field associated with Ω and P is a probability measure and λu is a linear losses term.

Applying the eigenfunction expansion technique [14], the following general expressions for the ensemble average, covariance and variance of the solution process are obtained as follows:

$$\mu_u(t, x) = e^{-\lambda t} \left(\sum_{n=0}^{\infty} T_n e^{-\left(\frac{n\pi\alpha}{L}\right)^2 t} \sin \frac{n\pi}{L} x + \sum_{n=0}^{\infty} E I_n(t) \sin \frac{n\pi}{L} x \right),$$

$$T_n = \frac{2}{L} \int_0^L g(x) \sin \frac{n\pi}{L} x dx,$$

$$E I_n = e^{-\left(\frac{n\pi\alpha}{L}\right)^2 t} \int_0^t e^{\left(\frac{n\pi\alpha}{L}\right)^2 \tau} E F_n(\tau) d\tau, \quad (2)$$

$$E F_n(\tau) = \frac{2e^{\lambda \tau}}{L} \int_0^L E f(t, x; \omega) \sin \frac{n\pi}{L} x dx.$$

where E denotes the average operator,

*Address correspondence to this author at the University Dammam, Faculty of science, Mathematics Department, Dammam, Saudi Arabia; Tel: 00966505846330; Fax: 0096638151026; E-mail: n_almulla@yahoo.com

$$Cov(u(t_1, x), u(t_2, x)) = e^{-\lambda(t_1+t_2)} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} V_{nm} \sin \frac{n\pi}{L} x \sin \frac{m\pi}{L} x$$

$$V_{nm} = e^{-\left(\frac{n\pi\alpha}{L}\right)^2 t_1} e^{-\left(\frac{m\pi\alpha}{L}\right)^2 t_2} \int_0^{t_1} \int_0^{t_2} e^{\left(\frac{n\pi\alpha}{L}\right)^2 \tau_1} e^{\left(\frac{m\pi\alpha}{L}\right)^2 \tau_2} MFnFm d\tau_2 d\tau_1$$

$$MFnFm = \frac{4}{L^2} e^{\lambda(\tau_1+\tau_2)} \int_0^L \int_0^L E f(\tau_1, x_1; \omega) f(\tau_2, x_2; \omega) \sin \frac{n\pi}{L} x_1 \sin \frac{m\pi}{L} x_2 dx_2 dx_1.$$

$$Varu(t, x) = Cov(u(t_1, x), u(t_2, x)) \Big|_{t_1=t_2=t} \cdot (4)$$

Considering white noise, $\sigma n(x; \omega)$, or simply $\sigma n(x)$ as a random nonhomogeneity with a deterministic scale σ , the following results are obtained using the statistical properties of white noise

$$\mu_u(t, x) = e^{-\lambda t} \left(\sum_{n=0}^{\infty} T_n e^{-\left(\frac{n\pi\alpha}{L}\right)^2 t} \sin \frac{n\pi}{L} x \right). \tag{5}$$

$$Cov(u(t_1, x), u(t_2, x)) = e^{-\lambda(t_1+t_2)} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} V_{nm} \sin \frac{n\pi}{L} x \sin \frac{m\pi}{L} x$$

$$V_{nm} = e^{-\left(\frac{n\pi\alpha}{L}\right)^2 t_1} e^{-\left(\frac{m\pi\alpha}{L}\right)^2 t_2} \int_0^{t_1} \int_0^{t_2} e^{\left(\frac{n\pi\alpha}{L}\right)^2 \tau_1} e^{\left(\frac{m\pi\alpha}{L}\right)^2 \tau_2} MFnFm d\tau_2 d\tau_1 \tag{6}$$

$$MFnFm = \frac{4\sigma^2}{L^2} e^{\lambda(\tau_1+\tau_2)} \int_0^L \sin \frac{n\pi}{L} x \sin \frac{m\pi}{L} x dx$$

Expressions (5) and (6) are the exact solutions of the linear problem. For the sake of validating the WHEP technique [12], the technique was applied on the linear problem where the Gaussian solution can be expanded, for a first order approximation, as follows:

$$u(t, x; \omega) = u^{(0)}(t, x) + \int_0^L u^{(1)}(t, x; x_1) H^{(1)}(x_1; \omega) dx_1. \tag{7}$$

where $u^{(0)}(t, x)$ and $u^{(1)}(t, x; x_1)$ are deterministic kernels to be evaluated and $H^{(1)}(x_1; \omega)$ is the Wiener-Hermite polynomial of order one; mainly is the same as white noise $n(x_1)$. In this case, the average takes the form

$$\mu_u = u^{(0)}(t, x), \tag{8}$$

while the covariance takes the form

$$Cov(u(t_1, x), u(t_2, x)) = \int_0^L u^{(1)}(t_1, x; x_1) u^{(1)}(t_2, x; x_1) dx_1. \tag{9}$$

In the WHEP technique, the deterministic kernels are expanded as a power series in the parameter λ as follows:

$$u^{(j)}(t, x) = u_0^{(j)}(t, x) + \lambda u_1^{(j)}(t, x) + \lambda^2 u_2^{(j)}(t, x) + \dots, j=0,1. \tag{10}$$

Following the technique algorithm [12], the following results are obtained for the first correction (up to λ):

$$u_0^{(0)}(t, x) = \sum_{n=0}^{\infty} T_n e^{-\left(\frac{n\pi\alpha}{L}\right)^2 t} \sin \frac{n\pi}{L} x, \tag{11}$$

$$u_1^{(0)}(t, x) = \sum_{n=0}^{\infty} I_0 \sin \frac{n\pi}{L} x,$$

$$I_0 = \int_0^t e^{-\left(\frac{n\pi\alpha}{L}\right)^2 (t-\tau)} F_n(\tau) d\tau, \tag{12}$$

$$F_n(t) = \frac{-2}{L} \int_0^L u_0^{(0)}(t, x) \sin \frac{n\pi}{L} x dx,$$

$$u_0^{(1)}(t, x; x_1) = \sum_{n=0}^{\infty} I_1 \sin \frac{n\pi}{L} x,$$

$$I_1 = \int_0^t e^{-\left(\frac{n\pi\alpha}{L}\right)^2 (t-\tau)} F_{n1}(\tau) d\tau, \tag{13}$$

$$F_{n1}(t) = \frac{2}{L} \int_0^L \delta(x - x_1) \sin \frac{n\pi}{L} x dx$$

$$= \frac{2}{L} \sin \frac{n\pi}{L} x_1,$$

$$u_1^{(1)}(t, x; x_1) = \sum_{n=0}^{\infty} I_{11} \sin \frac{n\pi}{L} x,$$

$$I_{11} = \int_0^t e^{-\left(\frac{n\pi\alpha}{L}\right)^2 (t-\tau)} F_{n11}(\tau) d\tau, \tag{14}$$

$$F_{n11}(t) = \frac{-2}{L} \int_0^L u_0^{(1)}(t, x; x_1) \sin \frac{n\pi}{L} x dx.$$

The first order, first correction solution is

$$\mu_{u1} = u_0^{(0)}(t, x) + \lambda u_1^{(0)}(t, x) \tag{15}$$

$$Cov(u(t_1, x), u(t_2, x)) = \int_0^L u_0^{(1)}(t_1, x; y) u_0^{(1)}(t_2, x; y) dy$$

$$+ \lambda \left[\int_0^L u_1^{(1)}(t_1, x; y) u_0^{(1)}(t_2, x; y) dy + \int_0^L u_1^{(1)}(t_2, x; y) u_0^{(1)}(t_1, x; y) dy \right]$$

$$+ \lambda^2 \left[\int_0^L u_1^{(1)}(t_1, x; y) u_1^{(1)}(t_2, x; y) dy \right] \tag{16}$$

The results for the second correction (up to λ^2) are

$$u_2^{(0)}(t, x) = \sum_{n=0}^{\infty} I_{02} \sin \frac{n\pi}{L} x,$$

$$I_{02} = \int_0^t e^{-\left(\frac{n\pi\alpha}{L}\right)^2 (t-\tau)} F_{n02}(\tau) d\tau, \tag{17}$$

$$F_{n02}(t) = \frac{-2}{L} \int_0^L u_1^{(0)}(t, x) \sin \frac{n\pi}{L} x dx,$$

$$F_{n_{n+1}}(t) = \frac{2\sigma}{L} \int_0^L n(x) \sin \frac{m\pi}{L} x dx - \frac{2\varepsilon}{L} \int_0^L u_n^2(t,x) \sin \frac{m\pi}{L} x dx \quad (30)$$

If the convergence of the process is insured, one can obtain the solution as

$$u(t,x) = \lim_{n \rightarrow \infty} u_n(t,x). \quad (31)$$

One can notice that all order of approximations are stochastic processes. The ensemble average of the zero order approximation is obtained as

$$\mu_{u_0}(t,x) = \sum_{n=0}^{\infty} T_n e^{-\left(\frac{n\pi}{L}\right)^2 t} \sin \frac{n\pi}{L} x. \quad (32)$$

The covariance of u_0 is given by

$$\begin{aligned} Cov(u_0(t,x_1), u_0(t,x_2)) &= \frac{4\sigma^2}{L^2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} I_{n,m} \left(\int_0^t e^{-\left(\frac{n\pi}{L}\right)^2 (t-\tau)} d\tau \right) \\ &\quad \left(\int_0^t e^{-\left(\frac{m\pi}{L}\right)^2 (t-\tau)} d\tau \right) \sin \frac{n\pi}{L} x_1 \sin \frac{m\pi}{L} x_2 \end{aligned} \quad (33)$$

where

$$I_{n,m} = \int_0^L \sin \frac{n\pi}{L} x \sin \frac{m\pi}{L} x dx. \quad (34)$$

The variance is

$$\begin{aligned} \sigma_{u_0}^2(t,x) &= \frac{4\sigma^2}{L^2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} I_{n,m} \left(\int_0^t e^{-\left(\frac{n\pi}{L}\right)^2 (t-\tau)} d\tau \right) \left(\int_0^t e^{-\left(\frac{m\pi}{L}\right)^2 (t-\tau)} d\tau \right) \\ &\quad \sin \frac{n\pi}{L} x \sin \frac{m\pi}{L} x. \end{aligned} \quad (35)$$

The following results for the first order approximation are obtained:

$$\mu_{u_1}(t,x) = \sum_{n=0}^{\infty} T_n e^{-\left(\frac{n\pi}{L}\right)^2 t} \sin \frac{n\pi}{L} x + \sum_{n=0}^{\infty} \left(\int_0^t e^{-\left(\frac{n\pi}{L}\right)^2 (t-\tau)} E F_{n_1}(\tau) d\tau \right) \sin \frac{n\pi}{L} x \quad (36)$$

where

$$E F_{n_1}(t) = \frac{-2\lambda}{L} \int_0^L (\sigma_{u_0}^2 + \mu_{u_0}^2) \sin \frac{n\pi}{L} x dx, \quad (37)$$

$$\begin{aligned} Cov(u_1(t,x_1), u_1(t,x_2)) &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} E I_{n_1}(t) I_{m_1}(t) \sin \frac{n\pi}{L} x_1 \sin \frac{m\pi}{L} x_2 \\ &\quad - \sum_{n=0}^{\infty} \left(\int_0^t e^{-\left(\frac{n\pi}{L}\right)^2 (t-\tau)} E F_{n_1}(\tau) d\tau \right) \sin \frac{n\pi}{L} x_2 \sum_{m=0}^{\infty} E I_{m_1}(t) \sin \frac{m\pi}{L} x_1 \\ &\quad - \sum_{m=0}^{\infty} \left(\int_0^t e^{-\left(\frac{m\pi}{L}\right)^2 (t-\tau)} E F_{m_1}(\tau) d\tau \right) \sin \frac{m\pi}{L} x_1 \sum_{n=0}^{\infty} E I_{n_1}(t) \sin \frac{n\pi}{L} x_2 \\ &\quad + \sum_{n=0}^{\infty} E I_{n_1}(t) \sin \frac{n\pi}{L} x_1 \sum_{m=0}^{\infty} E I_{m_1}(t) \sin \frac{m\pi}{L} x_2, \end{aligned} \quad (38)$$

$$E I_{n_1}(t) = \int_0^t e^{-\left(\frac{n\pi}{L}\right)^2 (t-\tau)} E F_{n_1}(\tau) d\tau, \quad (39)$$

$$E I_{n_1}(t) I_{m_1}(t) = \int_0^t \int_0^t e^{-\left(\frac{n\pi}{L}\right)^2 (t-\tau_1)} e^{-\left(\frac{m\pi}{L}\right)^2 (t-\tau_2)} E F_{n_1}(\tau_1) F_{m_1}(\tau_2) d\tau_1 d\tau_2 \quad (40)$$

in which

$$\begin{aligned} E F_{n_1}(\tau_1) F_{m_1}(\tau_1) &= \frac{4\sigma^2}{L^2} \int_0^L \sin \frac{n\pi}{L} x \sin \frac{m\pi}{L} x dx \\ &\quad - \frac{4\lambda\sigma}{L^2} \int_0^L \int_0^L \sin \frac{n\pi}{L} x_3 \sin \frac{m\pi}{L} x_4 E n(x_3) u_0^2(\tau_2, x_4) dx_3 dx_4 \\ &\quad - \frac{4\lambda\sigma}{L^2} \int_0^L \int_0^L \sin \frac{n\pi}{L} x_3 \sin \frac{m\pi}{L} x_4 E n(x_4) u_0^2(\tau_1, x_3) dx_3 dx_4 \\ &\quad + \frac{4\lambda^2}{L^2} \int_0^L \int_0^L \sin \frac{n\pi}{L} x_3 \sin \frac{m\pi}{L} x_4 E u_0^2(\tau_1, x_3) u_0^2(\tau_2, x_4) dx_3 dx_4 \end{aligned} \quad (41)$$

where

$$E n(x) u_0^2(t,y) = 2 \left(\sum_{n=0}^{\infty} T_n e^{-\left(\frac{n\pi}{L}\right)^2 t} \sin \frac{n\pi}{L} y \right) \sum_{m=0}^{\infty} \sin \frac{m\pi}{L} y E n(x) I_{m_0}(t) \quad (42)$$

in which

$$E n(x) I_{n_0}(t) = \frac{2\sigma}{L} \sin \frac{n\pi}{L} x \int_0^t e^{-\left(\frac{n\pi}{L}\right)^2 (t-\tau)} d\tau, \quad (43)$$

and

$$\begin{aligned} E u_0^2(\tau_1, x_3) u_0^2(\tau_2, x_4) &= \phi^2(\tau_1, x_3) \phi^2(\tau_2, x_4) \\ &\quad + \phi^2(\tau_1, x_3) \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} E I_{n_0}(\tau_2) I_{m_0}(\tau_2) \cdot \sin \frac{n\pi}{L} x_4 \sin \frac{m\pi}{L} x_4 \\ &\quad + 4\phi(\tau_1, x_3) \phi(\tau_2, x_4) \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} E I_{n_0}(\tau_1) I_{m_0}(\tau_2) \cdot \sin \frac{n\pi}{L} x_3 \sin \frac{m\pi}{L} x_4 \\ &\quad + \phi^2(\tau_2, x_4) \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} E I_{n_0}(\tau_1) I_{m_0}(\tau_1) \cdot \sin \frac{n\pi}{L} x_3 \sin \frac{m\pi}{L} x_3 \\ &\quad + \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} E I_{n_0}(\tau_1) I_{m_0}(\tau_1) I_{l_0}(\tau_2) I_{k_0}(\tau_2) \cdot \\ &\quad \sin \frac{n\pi}{L} x_3 \sin \frac{m\pi}{L} x_3 \sin \frac{l\pi}{L} x_4 \sin \frac{k\pi}{L} x_4 \end{aligned} \quad (44)$$

in which

$$\phi(t,x) = \sum_{n=0}^{\infty} T_n e^{-\left(\frac{n\pi}{L}\right)^2 t} \sin \frac{n\pi}{L} x, \quad (45)$$

$$E I_{n_0}(\tau_1) I_{m_0}(\tau_1) = \frac{4\sigma^2}{L^2} I_{n,m} \int_0^{\tau_1} \int_0^{\tau_1} e^{-\left(\frac{n\pi}{L}\right)^2 (\tau_1-\tau_3)} e^{-\left(\frac{m\pi}{L}\right)^2 (\tau_1-\tau_4)} d\tau_3 d\tau_4 \quad (46)$$

$$E I_{n_0}(\tau_2) I_{m_0}(\tau_2) = \frac{4\sigma^2}{L^2} I_{n,m} \int_0^{\tau_2} \int_0^{\tau_2} e^{-\left(\frac{n\pi}{L}\right)^2 (\tau_2-\tau_3)} e^{-\left(\frac{m\pi}{L}\right)^2 (\tau_2-\tau_4)} d\tau_3 d\tau_4 \quad (47)$$

$$E I_{n_0}(\tau_1) I_{m_0}(\tau_2) = \frac{4\sigma^2}{L^2} I_{n,m} \int_0^{\tau_1} \int_0^{\tau_2} e^{-\left(\frac{n\pi}{L}\right)^2 (\tau_1-\tau_3)} e^{-\left(\frac{m\pi}{L}\right)^2 (\tau_2-\tau_4)} d\tau_3 d\tau_4 \quad (48)$$

$$EI_{n_0}(\tau_1)I_{m_0}(\tau_1)I_{l_0}(\tau_2)I_{k_0}(\tau_2) = \int_0^{\tau_1} \int_0^{\tau_1} \int_0^{\tau_2} \int_0^{\tau_2} e^{-\frac{n\pi}{L}(\tau_1-\tau_3)} e^{-\frac{m\pi}{L}(\tau_1-\tau_4)} e^{-\frac{l\pi}{L}(\tau_2-\tau_5)} e^{-\frac{k\pi}{L}(\tau_2-\tau_6)} EF_{n_0}(\tau_3)F_{m_0}(\tau_4)F_{l_0}(\tau_5)F_{k_0}(\tau_6) d\tau_3 d\tau_4 d\tau_5 d\tau_6, \tag{49}$$

where

$$EF_{n_0}(\tau_3)F_{m_0}(\tau_4)F_{l_0}(\tau_5)F_{k_0}(\tau_6) = \frac{16\sigma^4}{L^4} \int_0^L \int_0^L \sin \frac{n\pi}{L} x_1 \sin \frac{m\pi}{L} x_1 \sin \frac{l\pi}{L} x_2 \sin \frac{k\pi}{L} x_2 dx_1 dx_2 + \frac{16\sigma^4}{L^4} \int_0^L \int_0^L \sin \frac{n\pi}{L} x_1 \sin \frac{l\pi}{L} x_1 \sin \frac{m\pi}{L} x_2 \sin \frac{k\pi}{L} x_2 dx_1 dx_2 + \frac{16\sigma^4}{L^4} \int_0^L \int_0^L \sin \frac{n\pi}{L} x_1 \sin \frac{k\pi}{L} x_1 \sin \frac{m\pi}{L} x_2 \sin \frac{l\pi}{L} x_2 dx_1 dx_2. \tag{50}$$

Using mathematica-5, the previous huge computations were performed and the following sample results are obtained; Figs. (5-10).




Fig. (5). The change of the mean of the zero order approximation u_0 with time t and space variable x . ($L = 1, M = 1, \sigma = 1, g(x) = x$).




Fig. (6). The change of the variance of the zero order approximation u_0 with space variable x and time t . ($L = 1, M = 1, \sigma = 1, g(x) = x$).




Fig. (7). The change of the mean of the first order approximation u_1 with space x and time t at $\lambda = 0$. ($L = 1, M = 1, \sigma = 1, g(x) = x$).




Fig. (8). The change of the mean of the first order approximation u_1 with space x and time t at $\lambda = 1000$. ($L = 1, M = 1, \sigma = 1, g(x) = x$).




Fig. (9). Comparisons between the mean of the zero and first orders approximation with time t at different λ levels, $x=0.5$. ($L = 1, M = 1, \sigma = 1, g(x) = x$).




Fig. (10). Comparisons between the mean of the zero and first orders approximation with time t at different λ levels, $x=.5$ ($L = 1, M = 1, \sigma = 1, g(x) = x$).

4. THE DIFFUSION EQUATION UNDER NONLINEAR LOSSES, THE WHEP TECHNIQUE

Since Meecham and his co-workers [15] developed a theory of turbulence involving a truncated Wiener-Hermite expansion (WHE) of the velocity field, many authors studied problems concerning turbulence [16-21] and others [22-31].

The application of the WHE aims at finding a truncated series solution to the solution process of differential equations. The truncated series composes of two major parts; the first is the Gaussian part which consists of the first two terms, while the rest of the series constitute the non-Gaussian part. In nonlinear cases, there exists always difficulties of solving the resultant set of deterministic integro-differential equations got from the applications of a set of comprehensive averages on the stochastic integro-differential equation obtained after the direct application of WHE. Many authors introduced different methods to face these obstacles. Among them, the WHEP technique was introduced in [28] using the perturbation technique to solve perturbed nonlinear problems. The WHE method utilizes the Wiener-Hermite polynomials which are the elements of a complete set of statistically orthogonal random functions [13].

Due to the completeness of the Wiener-Hermite set, any random function $G(t; \omega)$ can be expanded as

$$G(t; \omega) = G^{(0)}(t) + \int_{-\infty}^{\infty} G^{(1)}(t; t_1) H^{(1)}(t_1) dt_1 + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} G^{(2)}(t; t_1, t_2) H^{(2)}(t_1, t_2) dt_1 dt_2 + \dots \tag{51}$$

where the first two terms are the Gaussian part of $G(t; \omega)$. The rest of the terms in the expansion represent the non-Gaussian part of $G(t; \omega)$. The average of $G(t; \omega)$ is

$$\mu_G = E G(t; \omega) = G^{(0)}(t) \tag{52}$$

The covariance of $G(t; \omega)$ is

$$\begin{aligned} Cov(G(t; \omega), G(\tau; \omega)) &= E(G(t; \omega) - \mu_G(t))(G(\tau; \omega) - \mu_G(\tau)) \\ &= \int_{-\infty}^{\infty} G^{(1)}(t; t_1) G^{(1)}(\tau, t_1) dt_1 + \\ & 2 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} G^{(2)}(t; t_1, t_2) G^{(2)}(\tau, t_1, t_2) dt_1 dt_2 \end{aligned} \tag{53}$$

The variance of $G(t; \omega)$ is

$$\begin{aligned} Var G(t; \omega) &= E(G(t; \omega) - \mu_G(t))^2 \\ &= \int_{-\infty}^{\infty} [G^{(1)}(t; t_1)]^2 dt_1 + 2 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [G^{(2)}(t; t_1, t_2)]^2 dt_1 dt_2 \end{aligned} \tag{54}$$

The WHEP technique can be applied on linear or nonlinear perturbed systems described by ordinary or partial differential equations. The solution can be modified in the sense that additional parts of the Wiener-Hermite expansion can always be taken into considerations and the required order of approximations can always be made depending on the computing tool. It can be even run through a package if it is coded in some sort of symbolic languages. The technique was successfully applied to several nonlinear stochastic equations [32,33], the symbolic WHEP algorithm is charted in appendix-A.

Considering equation (21) and searching for the Gaussian part of the solution process, equation (7), with taking the necessary averages, we get the following four sets of deterministic equations:

$$\frac{\partial u_0^{(0)}(t, x)}{\partial t} = \frac{\partial^2 u_0^{(0)}}{\partial x^2} \tag{55}$$

$$u_0^{(0)}(t, 0) = 0, u_0^{(0)}(t, L) = 0 \text{ and } u_0^{(0)}(0, x) = g(x),$$

$$\frac{\partial u_1^{(0)}(t, x)}{\partial t} = \frac{\partial^2 u_1^{(0)}}{\partial x^2} - [u_0^{(0)}]^2 - \int_0^L [u_0^{(1)}]^2 dx_1, \tag{56}$$

$$u_1^{(0)}(t, 0) = 0, u_1^{(0)}(t, L) = 0 \text{ and } u_1^{(0)}(0, x) = 0,$$

$$\frac{\partial u_0^{(1)}(t, x; x_1)}{\partial t} = \frac{\partial^2 u_0^{(1)}}{\partial x^2} + \sigma \delta(x - x_1) \tag{57}$$

$$u_0^{(1)}(t, 0; x_1) = 0, u_0^{(1)}(t, L; x_1) = 0 \text{ and } u_0^{(1)}(0, x; x_1) = 0,$$

$$\frac{\partial u_1^{(1)}(t, x; x_1)}{\partial t} = \frac{\partial^2 u_1^{(1)}}{\partial x^2} - 2u_0^{(0)} u_0^{(1)} \tag{58}$$

$$u_1^{(1)}(t, 0; x_1) = 0, u_1^{(1)}(t, L; x_1) = 0 \text{ and } u_1^{(1)}(0, x; x_1) = 0.$$

The algorithm of solution is evaluating $u_0^{(0)}$ and $u_0^{(1)}$ first using the separation of variables and the eigenfunction expansion respectively and then computing the other two kernels independently using the eigenfunction expansion. The final results of the first order first correction mean and variance respectively are:

$$\mu_u(x, t) = u_0^{(0)}(x, t) + \lambda u_1^{(0)}(x, t), \tag{59}$$

$$\text{Var } u(t,x) = \int_0^L [u_0^{(1)}(t,x;x_1)]^2 dx_1 + 2\lambda \int_0^L u_0^{(1)} u_1^{(1)} dx_1. \quad (60)$$

The following are some sample results Figs. (11-18).




Fig. (11). The change of the first order approximation and fourth order correction of the mean with time t at x=0.5 and $\lambda=1$, ($L=1, M=1, \sigma=1, g(x)=x$).




Fig. (12). The change of the first order approximation and fourth order correction of the variance with time t at x=0.5 and $\lambda=1$, ($L=1, M=1, \sigma=1, g(x)=x$).




Fig. (13). The change of the first order approximation and fourth order correction of the mean with time t at x=0.5 and $\lambda=0.001$. ($L=1, M=1, \sigma=1, g(x)=x$).




Fig. (14). The change of the first order approximation and fourth order correction of the variance with time t at x=0.5 and $\lambda=0.001$. ($L=1, M=1, \sigma=1, g(x)=x$).




Fig. (15). The change of the first order approximation and fourth order correction of the mean-error with time t at x=0.5, $\lambda=1$ ($L=1, M=1, \sigma=1, g(x)=x$).




Fig. (16). The change of the first order approximation and fourth order correction of the variance-error with time t at x=0.5, $\lambda=1$ ($L=1, M=1, \sigma=1, g(x)=x$).




Fig. (17). The change of the first order approximation and fourth order correction of the mean-error with time t at x=0.5, $\lambda=0.001$ ($L=1, M=1, \sigma=1, g(x)=x$).




Fig. (18). The change of the first order approximation and fourth order correction of the variance-error with time t at x=0.5, $\lambda=0.001$ ($L=1, M=1, \sigma=1, g(x)=x$).

One can notice that the approximations are very close, especially second and higher corrections as shown in Tables 1 and 2.


Table 1. Errors of the WHEP Corrections of the First Order Mean Solution

Time \ Mean	0.0	0.1	0.2	0.3	0.4	0.5
First correction	0	0.000220419	0.000755961	0.00120395	0.00147417	0.00161483
Second correction	0	0.0000714979	0.000517808	0.00106582	0.00139979	0.00157618
Third correction	4.06506^{-20}	0.0000612266	0.000530039	0.00107424	0.00140498	0.00157945
Fourth correction	3.17478^{-20}	0.0000615896	0.000529407	0.00107373	0.00140465	0.00157926

Table 2. Errors of the WHEP Corrections of the First Order Variance Solution

Time \ Variance	0.0	0.1	0.2	0.3	0.4	0.5
First correction	0	3.73958^{-12}	8.05185^{-12}	8.58321^{-12}	8.65956^{-12}	8.87587^{-12}
Second correction	0	1.29844^{-11}	2.82209^{-11}	2.33893^{-11}	1.45704^{-11}	8.63275^{-12}
Third correction	0	1.29837^{-11}	2.82188^{-11}	2.33873^{-11}	1.45691^{-11}	8.63197^{-12}
Fourth correction	0	1.29837^{-11}	2.82188^{-11}	2.33873^{-11}	1.45691^{-11}	8.63197^{-12}

APPENDIX-A: THE SYMBOLIC WHEP ALGORITHM



CONCLUSIONS

Concerning the presented case studies in this paper, the WHEP technique offers corrections for the approximate solution which are true solution developments in the case of taking the exact solution as a reference. We Still have relative developments when taking an approximate solution as a reference.

REFERENCES

- [1] Kampe F. Random solutions of partial differential equations. Proc. 3rd Berkeley Symposium on Mathematical Statistics and Probability 1956; III: pp. 199-08.
- [2] Bhurcha R. A survey on the theory of random functions, The institute of mathematical sciences. Matscience Report India 1965; p. 31.
- [3] Lo Dato V. Stochastic processes in heat and mass transport, C Probabilistic methods in applied mathematics. Bharucha-Reid Ed. 1973; vol. 3; pp. 183-12.
- [4] Becus A. Random generalized solutions to the heat equations. J Math Anal Appl 1977; 60: 93-02.
- [5] Marcus R. Parabolic Ito equation with monotone nonlinearities. J Func Anal 1978; 29: 257-86.
- [6] Manthey R. Weak convergence of solutions of the heat equation with Gaussian noise. Math Nachr 1985; 123: 157-68.
- [7] Manthey R. Existence and uniqueness of a solution of a reaction-diffusion with polynomial nonlinearity and with noise disturbance. Math Nachr 1986; 125: 121-33.
- [8] Jetschke G. II. Most probable states of a nonlinear Brownian bridge. Forschungsergebnisse (Jena) 1986;N/86/20.
- [9] Jetschke G. III. Tunneling in a bistable infinite-dimensional potential. Forschungsergebnisse (Jena) 1986; N/86/40.
- [10] El-Tawil M. Nonhomogeneous boundary value problems. J Math Anal Appl 1996; 200: 53-65.
- [11] Uemura H. Construction of the solution of 1-dim heat equation with white noise potential and its asymptotic behaviour. Stoch Anal Appl 1996; 14(4): 487-06.
- [12] El-Tawil M. The application of WHEP technique on partial differential equations. Int J Differ Equant Appl 2003; 7(3): 325-37.
- [13] El-Tawil M. The homotopy Wiener-Hermite expansion and perturbation technique (WHEP). Trans Comput Sci (Springer) 2008; 159-80.
- [14] Farlow SJ. Partial differential equations for scientists and engineers. NY: Wiley 1982.
- [15] Crow S, Canavan G. Relationship between a Wiener-Hermite expansion and an energy cascade. J Fluid Mech 1970; 41(2): 387-03.
- [16] Saffman P. Application of Wiener-Hermite expansion to the diffusion of a passive scalar in a homogeneous turbulent flow. Phys Fluids 1969; 12(9): 1786-98.
- [17] Kahan W, Siegel A. Cameron-Martin-Wiener method in turbulence and in Burger's model : General formulae and application to late decay. J Fluid Mech 1970; 41(3) : 593-18.
- [18] Wang J, Shu S. Wiener-Hermite expansion and the inertial sub-range of a homogeneous isotropic turbulence. Phys Fluids 1974; 17(6).
- [19] Hogge H, Meecham W. Wiener-Hermite expansion applied to decaying isotropic turbulence using a renormalized time-dependent base. J Fluid Mech 1978; 85(2): 325-47.
- [20] Doi, Masaaki, Imamura , Tsutomu. Exact Gaussian solution for two-dimensional incompressible inviscid turbulent flow. J Phys Soc Jpn 1979; 46(4): 1358-59.
- [21] Kambe, Ryouichi, Doi, Masaaki, Imamura , Tsutomu. Turbulent flows near flat plates. J Phys Soc Jpn 1980; 49(2): 763-78
- [22] Joelson M, Ramamonjiarisoa A. Random fields of water surface waves using Wiener-Hermite functional series expansions. J Fluid Mech 2003; 496: 313-34.
- [23] Eftimiu C. First-order Wiener-Hermite expansion in the electromagnetic scattering by conducting rough surfaces. Radio Sci 1988; 23(5): 769-79.
- [24] Tamura Y, Nakayama J. Enhanced scattering from a thin film with one-dimensional disorder. Waves Random Complex Media 2005; 15(2): 269-95.
- [25] Tamura Y, Nakayama J. TE plane wave reflection and transmission from one-dimensional random slab. IEICE Trans Electron 2005; E88-C(4): 713-20.
- [26] Jahedi A, Ahmadi G. Application of Wiener-Hermite expansion to non-stationary random vibration of a Duffing oscillator. J Appl Mech 1983; Trans ASME 50(2): 436-42.
- [27] AbdelGawad E, El-Tawil M, Nassar M.A. Nonlinear oscillatory systems with random excitation. Modeling, Simul Control -B 1989; 23(1): 55-63.
- [28] Gawad E, El-Tawil M. General stochastic oscillatory systems. Appl Math Mod 1993; 17(6): 329-35.
- [29] El-Tawil M, Mahmoud G. The solvability of parametrically forced oscillators using WHEP technique. Mech Mech Eng 1999; 3(2): 181-88.
- [30] Tamura Y, Nakayama J. A formula on the Hermite expansion and its application to a random boundary value problem. IEICE Trans Electron 2003; E86-C(8): 1743-48.
- [31] Rubinstein R, Choudhari M. Uncertainty quantification for systems with random initial conditions using Wiener-Hermite expansions. Stud Appl Math 2005; 114(2): 167-88.
- [32] El-Tawil M, Al-Jihany A. Approximate solution of a mixed nonlinear stochastic oscillator. J. Comput Math Appl 2009; 58 : 2236-59.
- [33] Xu Y, Xu W, Mahmoud G. On a complex Duffing system with random excitation. Chaos Solit Fract 2008;35(1): 126-32 .

Received: August 04, 2009

Revised: October 27, 2009

Accepted: November 28, 2009

© Noha A. Al-Mulla; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.