



The Open Agriculture Journal

Content list available at: <https://openagriculturejournal.com>



RESEARCH ARTICLE

Growth, Nitrogen and Phosphorus Uptake of Sorghum Plants as Affected by Green Manuring with Pea or Faba Bean Shell Pod Wastes Using ^{15}N

Mohammed Al-Chammaa¹, Farid Al-Ain¹ and Fawaz Kurdali^{1,*}

¹Agriculture Department, Atomic Energy Commission of Syria, Damascus, Syria

Abstract:

Background:

During the freezing or canning preparation process of green grain leguminous, large amounts of shell pods are considered as agricultural organic wastes, which may be used as Green Manure (GM) for plant growth enhancement.

Objective:

Evaluation of the effectiveness of soil amended with shell pod wastes of pea (PGM) or faba bean (FGM) as GM on growth, nitrogen and phosphorus uptake in sorghum plants.

Methods:

Determination of the impact of adding four rates of nitrogen (0, 50, 100, and 150 kg N ha⁻¹) in the form of pea (PGM) or faba bean (FGM) shell pod wastes as GM on the performance of sorghum using the indirect ^{15}N isotopic dilution technique.

Results:

Sorghum plants responded positively and differently to the soil amendments with either GMs used, particularly, the PGM. In comparison with the control (N), soil amendment with an equivalent rate of 3.5 t ha⁻¹ of PGM (PGM₁₀₀) or with 6.5 t ha⁻¹ of FGM (FGM₁₅₀) almost doubled dry weight, N and P uptake in different plant parts of sorghum. Regardless of the GM used, estimated values of %Ndf_{gm} in sorghum plants ranged from 35% to 55% indicating that the use of pod shells as GM provided substantial portions and amounts of N requirements for sorghum. Moreover, nitrogen recoveries of added GM (%NUE_{gm}) ranged from 29 to 45% indicating that N in both of GM forms were used effectively. Accordingly, equivalent amounts to 17 - 48 kg N ha⁻¹ of inorganic fertilizer may be saved. The beneficial effect of incorporating pod shells in soil on sorghum N was mainly attributed to their N availability, besides to their effects on the improvement of soil N uptake, particularly when using PGM.

Conclusion:

The agricultural by-products of faba bean and pea pod shells could be used as GM for sorghum growth improvement by enhancing N and P uptake from soil and from the organic source.

Keywords: By-products, Green manure, Leguminous pod shells, ^{15}N , Sorghum, PGM, FGM.

Article History

Received: May 29, 2019

Revised: September 04, 2019

Accepted: September 24, 2019

1. INTRODUCTION

Green Manures (GMs) represent a promising approach to maintaining sustainable nutrients for crop growth [1]. The importance of GM is increasing in recent years because of the high cost of chemical fertilizers, increased risk of environmental pollution, and the need for sustainable cropping systems

[2]. Leguminous plants are commonly used as green manure in different cropping systems worldwide. Their residues are particularly useful as organic green manure due to their high nitrogen (N) contents, and because this N is more likely to become readily available for uptake by other plants [3, 4], providing a significant amount of N to the subsequent crop, replacing some portion of the economically and environmentally costly N fertilizers [5, 6]. Therefore, possible options to reduce the use of chemical fertilizers could be the implementation of leguminous plants as green manure in cropping

* Address correspondence to this author at the Agriculture Department, Atomic Energy Commission of Syria, Damascus, Syria; Tel: +963112132580; E-mail: fkurdali@aec.org.sy

systems and recycling of organic wastes [7]. In addition to nitrogen, the GMs also supply other nutrients that are needed for plant growth [8, 9].

To evaluate the benefits of GM for plant production, it is necessary to quantify the proportions and amounts of nitrogen derived from this organic material. The use of the ^{15}N isotope to assess the efficiency of GM fertilizers is extensively reported in the literature. The indirect ^{15}N isotopic dilution technique is widely used to determine the nitrogen fraction derived from GM. This method is based on the fact that plant received un-labeled organic residues (*e.g.* GM) with ^{15}N labeled fertilizer to the soil resulted in a lower ^{15}N in plant tissues compared to the control receiving no residues. This indicates that a dilution effect occurred and that a portion of plant N is derived from GM [3, 10 - 12].

Vicia faba L. (faba bean) and *Pisum sativum* L. (pea) are sources of high-quality protein and are often grown on a large scale worldwide [13]. These grain legumes are the most commonly used legume in the diets of people and can be used as vegetable, green, dried or canned. In Syria, as well as in other Mediterranean regions, seeds of both species are usually consumed green before maturity, after a simple process in which the pod shells are removed. They may be preserved for several months by freezing or canning for later consumption. Disposal of by-products generated by leguminous plant food processing represents a promising source of compounds that may be used for their nutritional properties [14]. During the manufacturing process of canned or frozen grain legumes, like green pea or faba beans, large amounts of shell pods are considered as agricultural organic wastes. Also, substantial amounts of these materials are usually thrown and discarded by housewives during freezing preparation of green grains at home. Effective use of such organic wastes in agriculture could be an important issue in developing countries. These plant materials can be used as unconventional feedstuffs for ruminants [15], or they are thrown away as worthless. On the other hand, Eusuf Zai *et al.* [8] reported that the indigenous agricultural wastes could be recycled and used as GM or compost to achieve the highest nutrient recovery for plant growth enhancement. The ability of grain legumes to fix atmospheric N_2 along with high N content in their tissues with a low C/N ratio would allow their residues 'like shell pods by-products' to make contributions to the N economy in farming systems, particularly in the organic agricultural system [16]. Recently, Fall *et al.* [17] reported that peanut (*Arachis hypogaea*) shells application improved soil chemical properties and tree growth under saline conditions. Thus, the use of such organic wastes, as green manure fertilizers, could be an alternative or additional way for their handling in agriculture. Therefore, the aims of this study were to evaluate the effectiveness of soil amended with shell pod wastes of pea

(PGM) or faba bean (FGM) as GM on growth, nitrogen and phosphorus uptake in sorghum plants using the indirect ^{15}N isotope dilution method.

2. MATERIALS AND METHODS

2.1. Soil Properties

The experiment was conducted in pots, each containing 10 kg of soil collected from Deir AL-Hajar research station, located southeast of Damascus, Syria, ($36^\circ 28' \text{E}$, $33^\circ 21' \text{N}$) at 617 m above sea level. The area is located within a semiarid region in which the average annual rainfall reaches 120 mm year⁻¹, and most precipitations occur between November and early April. For the last ten years, the average minimum temperature in winter was 1.3°C in January, while it increases to the average maximum temperature of 38.3°C in July. Some climatic data for the studied site collected during the growing season were close to those averaged over the last 10 years (2008 to 2017) as shown in Table 1. The soil is classified as clay loam, with an average 57.9% clay, 39.5% silt, and 2.6% sand. The main physical and chemical soil properties were pH 7.74; Electrical Conductivity (EC) 0.32 dS m⁻¹; organic matter 0.91%; total nitrogen 0.07%; available phosphorus 6.2 $\mu\text{g g}^{-1}$; ionic content (meq L⁻¹): chloride (Cl⁻) 0.74, bicarbonate (HCO_3^-) 0.97, sulfate (SO_4^{2-}) 1.27, calcium (Ca^{2+}) 1.15, potassium (K^+) 0.14, sodium (Na^+) 1.27 and magnesium (Mg^{2+}) 0.47; Cation Exchange Capacity (CEC) 29.08 meq 100 g⁻¹; nitrate (NO_3^-) 9.5 $\mu\text{g g}^{-1}$ and ammonium (NH_4^+) 6.8 $\mu\text{g g}^{-1}$.

2.2. Chemical Composition of Green Manure and Treatments

Shell pod wastes of pea (PGM) and faba bean (FGM) were used as a source of Green Manure (GM) for the growth of sorghum plants. After a young pod harvest of pea and faba bean, the green seeds were removed. Pod shells residues of both legumes' species were cut into 3 - 5 cm pieces, weighed, dried, and ground to a fine powder for total N determination by Kjeldahl method, in four replicates. The mean value of total N was 2.3% and 2.86% for faba bean and pea shell pod wastes, respectively. Moreover, total polyphenols in PGM and FGM was determined by the Folin-Ciocalteu method [18]. Lignin was determined using the Acid Detergent Fiber (ADF) method as outlined by Rowland and Roberts [19]. Total carbon in both pod shells was determined by a mass spectrometer (Integra-CN, PDZ Europa Scientific Instrument, UK). Data of polyphenols, lignin, ratio of polyphenols to total N, ratio of lignin /N, ratio of (lignin + polyphenols) to total N and C/N ratio are shown in Table 2.

Table 1. Some climatic data for the studied site during the experimental period and the last ten years average (from 2008 to 2017).

Year	Variable	July	August	September	October
2008-2017 average	T _{min} (C)	19.7	20.3	18.2	13.9
	T _{max} (C)	38.3	38.0	34.9	29.6
	RH (%)	65.6	64.0	64.2	66.1
	ET (mm day ⁻¹)	7.1	8.0	6.7	5.3

(Table 1) contd.....

2017	T _{min} (C)	21.3	20.0	18.4	13.0
	T _{max} (C)	40.6	38.5	36.4	28.8
	RH (%)	56.0	59.0	60.0	60.0
	ET (mm day ⁻¹)	6.4	7.8	5.8	4.9

T_{min} = minimum temperature, T_{max} = maximum temperature, RH= relative air humidity, ET= reference evapotranspiration.

Table 2. Data of total polyphenols (%), lignin (%), ratio of polyphenols to total N, ratio of lignin /N, ratio of (lignin + polyphenols) to total N and C/N ratio in pod shell wastes of faba bean (FGM) and pea (PGM).

Pod shells	Polyphenols (% Ph)	Lignin (L)	C/N	Ph/N	L+Ph/N	L/N
FGM	1.66±0.01 ^a	32.70±0.57 ^a	17.96±0.31 ^a	0.72±0.01 ^a	14.94±0.26 ^a	14.22±0.25 ^a
PGM	0.35±0.01 ^b	28.30±0.49 ^b	15.17±0.26 ^b	0.12±0.003 ^b	10.02±0.17 ^b	9.90±0.17 ^b
LSD _{0.05}	0.08	2.08	1.13	0.03	0.87	0.83

Note: Means within a column followed by the same letter are not significantly different ($P < 0.05$).

Table 3. Actual amounts of shell pod wastes of pea (PGP) and faba bean (FGP) added as green manure (GM) per pot, in addition to their equivalent rates per unit area.

Treatments	Amounts of Added GM (pot ⁻¹)		Equivalent Added Rates (ha ⁻¹)	
	Dry Matter (g pot ⁻¹)	Nitrogen(mg N pot ⁻¹)	Dry Matter (ton ha ⁻¹)	Nitrogen(kg N ha ⁻¹)
FGM ₅₀	9.45	217.4	2.17	50
FGM ₁₀₀	18.9	434.8	4.34	100
FGM ₁₅₀	28.36	652.2	6.51	150
PGM ₅₀	7.6	217.4	1.75	50
PGM ₁₀₀	15.2	434.8	3.5	100
PGM ₁₅₀	22.8	652.2	5.25	150

FGP or PGM₅₀₋₁₀₀₋₁₅₀: pod shells of faba bean (F) or pea (P) added as GM at rates of 50, 100, and 150 kg N ha⁻¹, respectively.

Pea and faba bean pod shells were incorporated into the soil, fifteen days before sorghum planting, at equivalent rates of 0, 50, 100, and 150 kg N ha⁻¹ (Table 3). They were mixed with the soil of each pot as GM by adopting the following treatments:

- (PGM) or (FGM) control, without green manure (N0).
- (PGM₅₀) addition of shell pod wastes of pea at a rate of 50 kg N ha⁻¹ (N50).
- (PGM₁₀₀) addition of shell pod wastes of pea at a rate of 100 kg N ha⁻¹ (N100).
- (PGM₁₅₀) addition of shell pod wastes of pea at a rate of 150 kg N ha⁻¹ (N150).
- (FGM₅₀) addition of shell pod wastes of faba bean at a rate of 50 kg N ha⁻¹ (N50).
- (FGM₁₀₀) addition of shell pod wastes of faba bean at a rate of 100 kg N ha⁻¹ (N100).
- (FGM₁₅₀) addition of shell pod wastes of faba bean at a rate of 150 kg N ha⁻¹ (N150).

2.3. Planting Procedures and ¹⁵N Application

Seeds of sorghum plants (*Sorghum bicolor* L.) were sown in pots containing soil previously amended with GMs (*i.e.* 15 days before sowing). After germination, seedlings were thinned to two plants per pot. Since soil contained a small amount of available phosphorus, an equivalent amount of 100 kg P₂O₅ ha⁻¹ in the form of triple phosphate was applied to the soil prior to

planting to all treatments. Pots were weighed every three days and water was added to maintain the soil moisture content at around 75% of field capacity throughout the experimental period. Pots were arranged in a randomized complete block design in four replicates and set outdoors under open field conditions.

To estimate nitrogen derived from green manure, using the indirect ¹⁵N isotopic dilution method, an equivalent fertilizer rate of 10 kg N ha⁻¹ (43.48 mg N pot⁻¹) in the form of urea enriched with 9.63% ¹⁵N atom excess was applied to the soils. The N fertilizer was applied to all treatments in two equally split applications (5 kg N ha⁻¹ for each application) at ten days intervals after planting. This procedure was followed to stabilize the ¹⁵N enrichment of the N pool and to minimize N immobilization.

2.4. Plant Harvest, Analysis and Calculations

Sorghum plants were harvested 90 days after planting (*i.e.* physiological maturity stage) and separated into shoots, roots, and panicles. Samples were dried to a constant weight at 70 °C and ground to a fine powder. Kjeldahl method was used to determine total nitrogen in the samples, and the ¹⁵N/¹⁴N isotopic ratio was determined using the emission spectrometry (Jasco-150, Japan). Phosphorus content was determined by dry ashing and measured calorimetrically by spectrophotometer (Termo Specronic, England).

Nitrogen fractions derived from the available sources were

calculated using the indirect ^{15}N isotopic dilution method [11, 12, 20] as follows:

The percent N derived from green manure (Ndf_{gm}) for both PGM and FGM was calculated using the following equation:

$$\% \text{Ndf}_{\text{gm}} = [1 - (\text{atom } \%^{15}\text{N excess}_{\text{treatment}} / \text{atom } \%^{15}\text{N excess}_{\text{control}})] 100.$$

Where treatment: plants amended with PGM or FGM; control: plants grown without any green manure application.

Amount of Ndf_{gm} (mg N pot^{-1}) was calculated as follows:

$$\text{Ndf}_{\text{gm}} = (\% \text{Ndf}_{\text{gm}} / 100) \text{ total N yield.}$$

Nitrogen Use Efficiency ($\% \text{NUE}_{\text{gm}}$) of added green manure was calculated by the following equation:

$$\% \text{NUE}_{\text{gm}} = (\text{Ndf}_{\text{gm}} / \text{N added as GM}) 100.$$

Percent (%) and amount (mg N pot^{-1}) of N derived from fertilizer (Ndff) were calculated using the following equations:

$$\% \text{Ndff} = (\text{atom } \%^{15}\text{N in excess}_{\text{plant}} / \text{atom } \%^{15}\text{N in excess}_{\text{fertilizer}}) 100.$$

$$\text{Ndff} = (\% \text{Ndff} / 100) \text{ total N}$$

Percent (%) and amount (mg N pot^{-1}) of N derived from soil (Ndfs) were calculated as follows:

$$\% \text{Ndfs} = 100 - (\% \text{Ndf}_{\text{gm}} + \% \text{Ndff}).$$

$$\text{Ndfs} = (\% \text{Ndfs} / 100) \text{ total N.}$$

2.5. Statistical Analysis

Data were subjected to Analysis of Variance (ANOVA) using the statistical program Statview, 4.57[®] Abacus Concepts, Berkley, Canada. Means were compared using the least significant difference (Fisher's LSD) test at a probability level of $P < 0.05$.

3. RESULTS

3.1. Dry Matter Yield, Nitrogen and Phosphorus Uptake

The data of dry matter yield (DM), nitrogen and phosphorus uptake in different plant parts of sorghum as affected by green manuring with pea pod shell wastes (PGM) are given in Table 4. Data in the whole plant of sorghum (leaves plus roots and panicles) are shown in Fig. (1). The addition of PGM at a rate of 50 kg Nha^{-1} (PGM_{50}), significantly increased total DM yield in sorghum plants as compared with the control. Raising the N rate (e.g. 100 and 150 kg Nha^{-1}) resulted in more significant increments in the DM, which did not significantly differ from each other. The percent increments

in total DM were 60, 99, and 98% in PGM_{50} , PGM_{100} , and PGM_{150} , respectively, as compared with the control PGM. It is noteworthy that DM of panicles increased by 49, 109, and 106% as compared with the control, for the same aforementioned treatments, respectively. Also, there were no significant differences in DM between PGM_{100} and PGM_{150} for different plant parts of sorghum. These results may indicate that PGM_{100} is the proper treatment for plant growth improvement. In comparison with the un-manured control treatment, PGM_{100} almost doubled DM in different plant parts of sorghum. The pattern of N yield in sorghum plants was relatively similar to that of dry matter yield. Amounts of nitrogen accumulated in the whole plant of sorghum significantly increased by 63, 123, and 132% in PGM_{50} , PGM_{100} , and PGM_{150} , respectively, as compared with the control. In panicles, the increases were 56, 124, and 126%, for the same treatments. Increasing the rate of nitrogen in the form of PGM from 100 to 150 kg N ha^{-1} did not significantly increase nitrogen yield. Also, total phosphorus content (mg P pot^{-1}) in the different plant parts of sorghum increased as a result of adding PGM. For panicles, the percent increments in total phosphorus content were 77, 185, and 200% in PGM_{50} , PGM_{100} , and PGM_{150} , respectively, as compared with the control PGM (Table 4). For the whole plant (Fig. 1), the amounts of P uptake were 86, 142, and 148% higher in PGM_{50} , PGM_{100} , and PGM_{150} than that in the control, for the abovementioned treatments, respectively. No significant difference was obtained between PGM_{100} and PGM_{150} treatments.

For Faba bean pod shells wastes (FGM), soil amended with FGM significantly increased dry matter yield of sorghum (Table 5 and Fig. 1). Soil amended with FGM at a rate of 50 kg N ha^{-1} (FGM_{50}), significantly increased DM yield in sorghum plants as compared with FGM. Raising the N rate (i.e. PGM_{100} and PGM_{150}) resulted in more significant increments in the DM. Percent increments in DM were 19, 103, and 105% for panicles, and 14, 64, and 101% for the whole plant in FGM_{50} , FGM_{100} , and FGM_{150} , respectively, as compared with the un manured control treatment (FGM). Total nitrogen yield by sorghum responded to FGM in a manner relatively similar to dry matter yield. The percent increments in N yield were 29, 111, and 109% for panicles and 21, 69, and 104% for the whole plant, in FGM_{50} , FGM_{100} , and FGM_{150} , respectively. Phosphorus uptake was increased in all plant parts of sorghum as a result of soil incorporated with faba bean shell pod wastes (FGM). FGM increased P yield by 30, 92, and 101%, for the whole plant; and by 37, 156, and 205% for panicles in FGM_{50} , FGM_{100} , and FGM_{150} , respectively, as compared with the control (FGM).




Fig. (1). Dry matter yield (g pot⁻¹), nitrogen (mg N pot⁻¹) and phosphorus (mg P pot⁻¹) uptake in the whole plant of sorghum as affected by green manuring with pea (PGM) or faba bean (FGM) pod shell wastes. Columns followed by the same letter are not significantly different ($P < 0.05$). Small letters above columns: comparison among nitrogen rates for PGM or FGM. Capital letters inside columns: comparison between PGM and FGM for each nitrogen rate.

Table 4. Dry matter (g pot⁻¹), nitrogen (mg N pot⁻¹) and phosphorus (mg P pot⁻¹) yield in different plant parts of sorghum as affected by green manuring with pea pod shell wastes (PGM).

Treatment	Panicles	Roots	Leaves
	Dry Matter Yield (g pot ⁻¹)		
PGM	1.76 ± 0.03 ^c	2.29 ± 0.07 ^c	6.78 ± 0.10 ^c
PGM ₅₀	2.62 ± 0.08 ^b	4.43 ± 0.1 ^b	9.84 ± 0.32 ^b
PGM ₁₀₀	3.67 ± 0.08 ^a	5.26 ± 0.06 ^a	12.60 ± 0.01 ^a
PGM ₁₅₀	3.63 ± 0.09 ^a	5.32 ± 0.09 ^a	12.45 ± 0.09 ^a
LSD _{0.05}	0.22	0.25	0.55
	N-yield (mg N pot ⁻¹)		
PGM	34.0 ± 1.11 ^c	22.7 ± 0.34 ^c	112.2 ± 1.12 ^d
PGM ₅₀	53.1 ± 1.08 ^b	44.3 ± 1.50 ^b	187.8 ± 3.03 ^c

(Table 4) contd....

PGM ₁₀₀	76.3± 1.36 ^a	54.1±0.36 ^a	246.3 ±4.63 ^b
PGM ₁₅₀	77.1± 2.32 ^a	53.9± 0.74 ^a	261.6 ±5.07 ^a
LSD _{0.05}	4.78	2.63	11.69
P-yield (mg P pot⁻¹)			
PGM	3.95± 0.31 ^c	3.02±0.19 ^d	9.96 ±0.82 ^c
PGM ₅₀	7.00±0.70 ^b	5.39± 0.14 ^c	19.00± 1.22 ^b
PGM ₁₀₀	11.25 ±0.68 ^a	7.50± 0.26 ^a	23.16±1.15 ^a
PGM ₁₅₀	11.84± 0.32 ^a	6.71±0.22 ^b	22.27 ±1.77 ^{ab}
LSD _{0.05}	1.65	0.64	3.96

Note: Means within a column followed by the same letter (a,b,c and d) are not significantly different ($P<0.05$). PGM: without GM; PGM₅₀, PGM₁₀₀, and PGM₁₅₀: pea pod shells added as GM at rates of 50, 100, and 150 kg N ha⁻¹, respectively.

Table 5. Dry matter (g pot⁻¹), nitrogen (mg N pot⁻¹) and phosphorus (mg P pot⁻¹) yield in different plant parts of sorghum as affected by green manuring with faba bean pod shell wastes (FGM).

Treatment	Panicles	Roots	Leaves
	Dry Matter Yield (g pot ⁻¹)		
FGM	1.76±0.03 ^c	2.29±0.07 ^d	6.78±0.10 ^d
FGM ₅₀	2.10±0.08 ^b	2.85±0.07 ^c	7.40±0.22 ^c
FGM ₁₀₀	3.58±0.04 ^a	4.41±0.11 ^b	9.67±0.16 ^b
FGM ₁₅₀	3.61±0.07 ^a	5.37±0.13 ^a	12.80±0.22 ^a
LSD _{0.05}	0.23	0.29	0.57
N-yield (mg N pot⁻¹)			
FGM0	34.0 ±1.11 ^c	22.7±0.34 ^d	112.2±1.12 ^d
FGM50	44.0±2.48 ^b	28.3±0.89 ^c	131.9±3.50 ^c
FGM100	71.7±1.04 ^a	42.8±0.33 ^b	171.0±4.56 ^b
FGM150	70.9±1.80 ^a	54.4±1.05 ^a	218.8±2.19 ^a
LSD _{0.05}	5.26	2.24	9.63
P-yield (mg P pot⁻¹)			
FGM	3.95±0.31 ^c	3.02±0.19 ^c	9.96±0.82 ^c
FGM ₅₀	5.43±0.48 ^c	4.20±0.15 ^b	12.41±0.34 ^b
FGM ₁₀₀	10.10±0.27 ^b	5.87±0.51 ^a	16.44±0.96 ^a
FGM ₁₅₀	12.05±0.75 ^a	6.18±0.39 ^a	15.92±0.51 ^a
LSD _{0.05}	1.50	1.06	2.16

Note: Means within a column followed by the same letter (a,b,c and d) are not significantly different ($P<0.05$). FGM: control, no green manure; FGM₅₀, FGM₁₀₀ and FGM₁₅₀: faba bean pod shells added as GM at rates of 50, 100 and 150 kg N ha⁻¹, respectively.

3.2. Nitrogen Uptake from Various Sources

In comparison with the un-manured control plants, soil incorporated with both forms of GM (*i.e.* PGM or FGM), resulted in lower atom %¹⁵N excess values in the different plant parts of sorghum (Table 6). The decrease in ¹⁵N in sorghum plants was related to the amount of N in the applied GM. Regardless of plant parts, the highest dilution was obtained in the N₁₅₀ followed by N₁₀₀ and then by N₅₀ and N treatments, for both PGM and FGM (Table 6). Percentages and amounts of nitrogen derived from fertilizer (Ndff), soil (Ndfs), and green manure (Ndf_{gm}) in different plant parts of sorghum as affected by green manuring with PGM or with FGM are shown in Tables 7 and 8, respectively.

For pea pod shells (PGM) treatments, %Ndff and %Ndfs values in the different plant parts as well as in the whole sorghum plant decreased with increasing rates of the applied PGM (Table 7). However, the percentage of nitrogen derived from PGM (%Ndf_{gm}) increased with increasing rates of the

applied PGM. Values of %Ndf_{gm} in the whole plant of sorghum were 15.7, 48.7, and 53.7% in PGM₅₀, PGM₁₀₀, and PGM₁₅₀, respectively. In regard to the amounts of N derived from the available sources, it can be shown from (Table 7) that soil amendment with PGM generally increased amounts of Ndff and Ndfs, as compared with the control, particularly in PGM₁₀₀. The percent increments in the amounts of Ndff and Ndfs in the whole plant was about 15% higher than the control in the latter treatment. Moreover, significant amounts of nitrogen were derived from green manure (Ndf_{gm}) which were increased by increasing rates of added PGM. The Ndf_{gm} values in the whole plant of sorghum were 99, 183, and 211 mg N pot⁻¹ in PGM₅₀, PGM₁₀₀, and PGM₁₅₀, respectively (Table 7).

For faba bean pod shells (FGM) treatments, the addition of GM addition in the form of faba bean shell pod wastes (FGM) also decreased %Ndff and %Ndfs values. Whereas, %Ndf_{gm} increased with increasing rates of the applied FGM (Table 8). %Ndf_{gm} values in the whole plant were 37, 48, and 55% for

FGM₅₀, FGM₁₀₀, and FGM₁₅₀, respectively. Moreover, it can be shown from Table 8 that the addition of FGM generally decreased the amounts of Ndff and Ndfs in the leaves, panicles, and the whole plant as compared with the control. However, N

uptake from both sources (*i.e.* Ndff and Ndfs) increased in roots, particularly in FGM₁₀₀ and FGM₁₅₀, where the percent increments either in Ndff or Ndfs values were 24 and 40% for both treatments, respectively,

Table 6. Atom % ¹⁵N_{in excess} in different plant parts of sorghum plants manured with pea (PGM) or faba bean (FGM) shell pod wastes.

Treatment	Leaves	Roots	Panicles	Whole plant
% ¹⁵ N _{in excess}				
PGM				
PGM	0.796±0.005 ^a	0.535±0.005 ^a	0.764±0.005 ^a	0.754±0.004 ^a
PGM ₅₀	0.503±0.003 ^b	0.414±0.005 ^b	0.483±0.003 ^b	0.485±0.003 ^b
PGM ₁₀₀	0.391±0.003 ^c	0.380±0.004 ^c	0.381±0.002 ^c	0.387±0.002 ^c
PGM ₁₅₀	0.353±0.007 ^d	0.353±0.005 ^d	0.334±0.003 ^d	0.349±0.005 ^d
LSD _{0.05}	0.015	0.014	0.010	0.011
FGM				
FGM	0.754±0.004 ^a	0.796±0.005 ^a	0.535±0.005 ^a	0.764±0.005 ^a
FGM ₅₀	0.476±0.005 ^b	0.514±0.006 ^b	0.379±0.003 ^b	0.430±0.005 ^b
FGM ₁₀₀	0.391±0.006 ^c	0.413±0.002 ^c	0.351±0.007 ^c	0.363±0.004 ^c
FGM ₁₅₀	0.343±0.002 ^d	0.359±0.005 ^d	0.314±0.007 ^d	0.315±0.004 ^d
LSD _{0.05}	0.014	0.017	0.018	0.013

Note: For each green manure type, means within a column followed by the same letter (a,b,c and d) are not significantly different ($P<0.05$). PGM & FGM: control without green manure; PGM & FGM₅₀, PGM & FGM₁₀₀, and PGM & FGM₁₅₀: pod shells of pea (P) or faba bean (F) added as GM at rates of 50, 100, and 150 kg N ha⁻¹, respectively.

Table 7. Proportions (%) and amounts (mg N pot⁻¹) of nitrogen derived from fertilizer (Ndff), soil (Ndfs) and green manure (Ndf_{gm}) in different plant parts of sorghum as affected by green manuring with pea pod shell wastes (PGM).

Treatment	Panicles		Roots		Leaves		Whole Plant	
	Ndff							
	mg	%	mg	%	mg	%	mg	%
PGM	2.7±0.1 ^b	7.93±0.05 ^a	1.26±0.02 ^c	5.55±0.05 ^a	9.3±0.13 ^b	8.26±0.05 ^a	13.2±0.2 ^c	7.83±0.04 ^a
PGM ₅₀	2.7±0.06 ^b	5.01±0.03 ^b	1.90±0.05 ^b	4.29±0.05 ^b	9.3±0.19 ^b	5.22±0.03 ^b	13.9±0.22 ^b	5.03±0.03 ^b
PGM ₁₀₀	3.1±0.07 ^a	3.95±0.02 ^c	2.13±0.02 ^a	3.94±0.04 ^c	10.0±0.13 ^a	4.06±0.40 ^c	15.2±0.12 ^a	4.02±0.02 ^c
PGM ₁₅₀	2.7±0.1 ^b	3.47±0.03 ^d	1.96±0.04 ^b	3.65±0.05 ^d	9.6±0.33 ^b	3.66±0.08 ^d	14.2±0.25 ^b	3.63±0.05 ^d
LSD _{0.05}	0.25	0.11	0.10	0.15	0.65	0.16	0.62	0.11
Ndfs								
PGM	31.3±1.01 ^b	92.07±0.05 ^a	21.4±0.35 ^c	94.45±0.05 ^a	102.9±1.0 ^b	91.74±0.05 ^a	155.7±1.51 ^c	92.17±0.04 ^a
PGM ₅₀	31.0±0.72 ^b	58.27±0.36 ^b	32.4±0.78 ^b	73.11±0.87 ^b	103.6±2.1 ^b	57.95±0.34 ^b	163.6±2.55 ^b	59.23±0.31 ^b
PGM ₁₀₀	35.1±0.76 ^a	45.94±0.23 ^c	36.3±0.33 ^a	67.12±0.71 ^c	111.0±1.4 ^a	45.07±0.37 ^c	178.4±1.40 ^a	47.33±0.19 ^c
PGM ₁₅₀	31.1±1.13 ^b	40.29±0.37 ^d	33.5±0.67 ^b	62.11±0.81 ^d	106.5±3.7 ^{ab}	40.67±0.86 ^d	167.5±2.90 ^b	42.67±0.55 ^d
LSD _{0.05}	2.83	0.87	1.74	2.14	7.0	1.54	6.74	1.02
Ndf _{gm}								
PGM	-	-	-	-	-	-	-	-
PGM ₅₀	19.5±0.39 ^c	36.72±0.39 ^c	10.0±0.71 ^c	22.59±0.93 ^c	65.8±1.1 ^c	36.82±0.37 ^c	98.7±0.25 ^c	35.74±0.34 ^c
PGM ₁₀₀	38.2±0.55 ^b	50.10±0.24 ^b	15.7±0.47 ^b	28.94±0.75 ^b	125.4±3.3 ^b	50.87±0.40 ^b	183.3±2.76 ^b	48.65±0.20 ^b
PGM ₁₅₀	43.4±1.15 ^a	56.26±0.40 ^a	18.4±0.5 ^a	34.24±0.86 ^a	145.6±2.6 ^a	55.67±0.94 ^a	210.8±3.17 ^a	53.71±0.60 ^a
LSD _{0.05}	2.45	1.13	1.81	2.71	7.96	2.00	7.77	1.32

Note: Means within a column followed by the same letter (a,b,c and d) are not significantly different ($P<0.05$). PGM: without GM; PGM₅₀, PGM₁₀₀, and PGM₁₅₀: pea pod shells added as GM at rates of 50, 100, and 150 kg N ha⁻¹, respectively.

Table 8. Proportions (%) and amounts (mg N pot⁻¹) of nitrogen derived from fertilizer (Ndff), soil (Ndfs), and green manure (Ndf_{gm}) in different plant parts of sorghum as affected by green manuring with faba bean pod shell wastes (FGM).

Treatment	Panicles		Roots		Leaves		Whole Plant	
	Ndff							
	mg	%	mg	%	mg	%	mg	%
FGM	2.7±0.1 ^a	7.93±0.05 ^a	1.26±0.02 ^c	5.55±0.05 ^a	9.3±0.13 ^a	8.26±0.05 ^a	13.2±0.20 ^a	7.83±0.04 ^a
FGM ₅₀	2.0±0.1 ^c	4.43±0.05 ^b	1.11±0.03 ^d	3.93±0.04 ^b	7.0±0.26 ^c	5.34±0.06 ^b	10.1±0.32 ^c	4.94±0.05 ^b
FGM ₁₀₀	2.7±0.02 ^a	3.77±0.04 ^c	1.56±0.03 ^b	3.64±0.07 ^c	7.3±0.30 ^c	4.28±0.07 ^c	11.6±0.32 ^b	4.06±0.06 ^c
FGM ₁₅₀	2.3±0.06 ^b	3.27±0.04 ^d	1.77±0.07 ^a	3.25±0.08 ^d	8.2±0.07 ^b	3.72±0.05 ^d	12.26±0.07 ^b	3.56±0.02 ^d
LSD _{0.05}	0.24	0.13	0.13	0.19	0.65	0.17	0.78	0.14
Ndfs								
FGM	31.3±1.0 ^a	92.07±0.05 ^a	21.4±0.35 ^c	94.45±0.1 ^a	103.0±1.0 ^a	91.74±0.05 ^a	155.7±1.51 ^a	92.17±0.04 ^a
FGM ₅₀	22.7±1.30 ^c	51.49±0.55 ^b	19.0±0.46 ^d	66.89±0.6 ^b	78.1±2.9 ^c	59.22±0.66 ^b	118.9±3.81 ^c	58.20±0.58 ^b
FGM ₁₀₀	31.4±0.25 ^a	43.78±0.44 ^c	26.5±0.53 ^b	61.94±1.3 ^c	81.3±3.3 ^c	47.55±0.72 ^c	136.3±3.77 ^b	47.76±0.71 ^c
FGM ₁₅₀	26.9±0.64 ^b	37.99±0.48 ^d	30.1±1.14 ^a	55.4±1.27 ^d	90.4±0.7 ^b	41.33±0.53 ^d	144.0±0.82 ^b	41.85±0.23 ^d
LSD _{0.05}	2.75	1.32	2.12	2.89	6.99	1.71	8.67	1.48
Ndf _{gm}								
FGM	-	-	-	-	-	-	-	-
FGM ₅₀	19.4±1.12 ^c	44.07±0.59 ^c	8.3±0.41 ^c	29.2±0.60 ^c	46.7±0.62 ^c	35.45±0.72 ^c	75.2±0.42 ^c	36.86±0.63 ^c
FGM ₁₀₀	37.6±0.86 ^b	52.46±0.50 ^b	14.7±0.61 ^b	34.4±1.33 ^b	82.2±1.17 ^b	48.16±0.79 ^b	137.5±0.88 ^b	48.18±0.77 ^b
FGM ₁₅₀	41.6±1.23 ^a	58.74±0.53 ^a	22.5±0.63 ^a	41.4±1.34 ^a	120.3±2.33 ^a	54.95±0.58 ^a	187.8±1.87 ^a	54.59±0.31 ^a
LSD _{0.05}	3.46	1.71	1.80	3.66	4.95	2.24	3.89	1.93

Note: Means within a column followed by the same letter (a,b,c and d) are not significantly different ($P<0.05$). FGM: control, no green manure; FGM₅₀, FGM₁₀₀, and FGM₁₅₀: faba bean pod shells added as GM at rates of 50, 100, and 150 kg N ha⁻¹, respectively.

Table 9. Nitrogen use efficiency (% NUE_{gm}) of added green manure in the form of pea (PGM) or faba bean (FGM) pod shells in the different plant parts of sorghum.

Treatment	Panicles		Roots		Leaves		Whole Plant	
	% NUE _{gm}							
	PGM							
PGM ₅₀	8.97±0.18 ^a	4.61±0.37 ^a	30.27±0.49 ^a	45.38±0.12 ^a				
PGM ₁₀₀	8.79±0.13 ^a	3.60±0.11 ^b	28.83±0.75 ^a	42.15±0.63 ^b				
PGM ₁₅₀	6.65±0.18 ^b	2.83±0.08 ^c	22.32±0.40 ^b	32.33±0.48 ^c				
LSD _{0.05}	0.52	0.65	1.81	1.49				
FGM								
FGM ₅₀	8.93±0.32 ^a	3.81±0.19 ^a	21.48±0.29 ^a	34.60±0.19 ^a				
FGM ₁₀₀	8.65±0.12 ^a	3.39±0.14 ^a	18.91±0.27 ^b	31.61±0.20 ^b				
FGM ₁₅₀	6.38±0.19 ^b	3.44±0.10 ^a	18.44±0.36 ^b	28.80±0.29 ^c				
LSD _{0.05}	1.08	N.S	0.98	0.74				

Note: For each green manure type, means within a column followed by the same letter (a,b,c and d) are not significantly different ($P<0.05$). PGM & FGM: control without green manure; PGM & FGM₅₀, PGM & FGM₁₀₀, and PGM & FGM₁₅₀: pod shells of pea (P) or faba bean (F) added as GM at rates of 50, 100, and 150 kg N ha⁻¹, respectively.

3.3. Nitrogen Use Efficiency (%NUE_{gm}) of Added Green Manure

Nitrogen use efficiency (%NUE_{gm}) of added green manure in the form of pea (PGM) or faba bean (FGM) pod shells in the different plant parts of sorghum are shown in Table 9. Regardless of GM forms, % NUE_{gm} values were high in leaves, followed by panicles and roots. In the whole plants, nitrogen use efficiency of added green manure decreased with increasing rates of applied GM. For pea pod shells (PGM)

treatments, %NUE_{gm} values were 45.4, 42.2, and 32.2% in PGM₅₀, PGM₁₀₀, and PGM₁₅₀, respectively. This result indicated that sorghum utilized approximately half of the N applied as pea shell pod wastes in PGM₅₀ and PGM₁₀₀ treatments, and third of the N applied in the PGM₁₅₀ treatment.

In regard to faba bean pod shells (FGM) treatments, the highest nitrogen recovery was in FGM₅₀. There were slight but significant decreases in %Ndf_{gm} as a result of increasing rates of applied GM. Values of %NUE_{gm} in the whole plant were 35, 32, and 29%, in FGM₅₀, FGM₁₀₀, and FGM₁₅₀, respectively.

This indicates that sorghum utilized approximately a third of the N applied in the form of faba bean shell pod wastes.

4. DISCUSSION

4.1. Effects of Green Manuring with Pea and Faba Bean Pod Shell on Dry Matter Production and N Yield of Sorghum

Performance of sorghum plants grown on the soil amended with pea (PGM) or faba bean (FGM) pod shells wastes as green manure (GM) was examined using indirect ^{15}N isotope dilution. Sorghum plants responded positively and differently to the soil amendments with either GMs used. The results showed that dry weight and N yields of sorghum plants significantly increased by soil amendment with PGM or FGM as compared with un-manured treatments. The beneficial effects of leguminous green manures on the growth, yield, and N-uptake of the following crops, particularly, the sorghum plants, have been previously observed in different cropping systems [3, 21 - 24]. The increase in dry weight and that in N uptake caused by the application of pea (PGM) or faba bean (FGM) pod shells as green manures in sorghum plants may be attributed to the increase in available N released from GM. Moreover, other nutrient elements such as phosphorus could also promote plant growth. The beneficial effects following the green manure amendment on sorghum plants were affected by the mass of incorporated GM material in the soil. In comparison with the un-manured control treatment (N0), soil amendment with 3.5 t ha^{-1} of PGM (PGM₁₀₀) or with 6.5 t ha^{-1} of FGM (FGM₁₅₀) almost doubled the above-mentioned parameters in different plant parts of sorghum. Similarly, Fall *et al.* [17] reported that soil amendment with 4 to 6 t ha^{-1} of peanut shells significantly increased seedlings growth of three leguminous trees. On the other hand, our study showed that growth and N uptake by sorghum plants were affected by the form of added GM. The enhancement in dry-matter production and N yield of sorghum was more pronounced in plants amended with pea than with faba bean pod shell residues. As indicated by Giller and Wilson [4], such differences may have resulted from variations in decomposition and immobilization rates between the two forms of plant material incorporated in the soil.

4.2. Effects of Green Manuring with Pea and Faba Bean Pod Shell on P uptake in Sorghum

In addition to the effect green manure of pea and faba bean pod shell on N uptake, the incorporation of GMs improved P content in sorghum plants. Bah, *et al.* [9] reported that improved availability of soil P due to the incorporation of plant materials has been attributed to direct P release from the decomposing materials and the action of other decomposition products on native soil P. Several researchers reported that the phosphorus in green manure can potentially be delivered to the soil in a form that is readily available to plants and soil microorganisms [25, 26]. Baddeley *et al.* [27] showed that the incorporation of a legume green manure can enhance biological phosphorus cycling in soil and improve its availability. In this study, the increase of phosphorus content in sorghum plants was relatively related to the mass of applied GM. In

comparison with the control (N0), total phosphorus content in sorghum plants increased by 86, 140, and 148% for PGM and by 30, 92, and 101% for FGM, in N₅₀, N₁₀₀, and N₁₅₀ treatments, respectively. Gao *et al.* [28] showed that the positive effect of alfalfa green manure on increasing yield of rice can be attributed to its good functions on increasing soil available P, promoting P uptake, and enhancing interactive effect of N and P. Such an organic amendment could increase some soil enzyme activities, such as dehydrogenase, urease, acid phosphatase, and β -glucosidase. The changes in soil enzyme activities affect the progress of nitrogen and phosphorus mineralization and release [28]. Pypers *et al.* [29] suggested that green manure with leguminous in crop rotation system increases the yield and growth of the maize plants, possibly for microbiological reasons, and the enhancement of P acquisition by plants resulted from improved soil P availability. On the other hand, phosphorus recovery following green manure crops may be derived from the decomposition and mineralization of incorporated plant material in the soil, from native P in the soil and from mineral P fertilizer. In this study, we did not estimate these values, only the total P content of the biomass. It is not possible to separate these P sources without using the ^{32}P isotope labeling technique [9]. However, it is evident that the incorporation of GMs improved P content in sorghum plants and may have contributed to yield increases. Based on the P-uptake under the two GM treatments (*i.e.* pea and faba bean pod shell), the phosphorus accumulation in total biomass of sorghum plants was 43, 29, and 20% higher in pea than faba bean, for N₅₀, N₁₀₀, and N₁₅₀ treatments, respectively. Such increments may be partly attributed to differences in soil enzyme activities between the two types of GMs [28].

4.3. Effect of Green Manuring with Pea and Faba Bean Pod Shell on N Derived from Various Sources in Sorghum

Sorghum plants amended with GM (*i.e.* pea or faba bean pod shells), had lower atom % ^{15}N excess values as compared to the un-manured control treatment. This result indicated that a dilution effect had taken place and that some portions of sorghum N were derived from the incorporated GM in the soil [3, 10 - 12]. Regardless of GM used, estimated values of %Ndf_{gm} in sorghum plants, using the indirect ^{15}N isotope dilution method, ranged from 35% to 55%. This result indicated that the use of pod shells of leguminous as green manures provided substantial portions of N requirements for sorghum. Amounts of nitrogen derived from GM (Ndf_{gm}) in the whole plant of sorghum were 99, 183, and 211 mg N/pot for PGM and 75, 138, and 188 mg/N pot for FGM in N₅₀, N₁₀₀, and N₁₅₀ treatments, respectively. As indicated by Rees *et al.* [30], the amounts of N taken up by plants were proportional to those applied in the form of legume residues. Moreover, based on amounts of N uptake from the two GM residues, (*i.e.* pea or faba bean pod shells), nitrogen accumulation in total biomass of sorghum plants were 31, 33, and 12% higher in pea than faba bean, for N₅₀, N₁₀₀, and N₁₅₀ treatments, respectively. Such divergence may result from differences in the decomposition and mineralization rates of organic N between the two forms of plant material incorporated in the soil [4]. Consequently, our results showed a beneficial effect of using pea or faba bean pod shells as GMs to meet some of the N-requirement in sorghum

plants.

With regard to soil (Ndfs) or fertilizer (Ndff) N uptake values (*i.e.* proportions and amounts), sorghum plants grown in a soil amended with pea pod shells (PGM) had significantly higher values in the different plant parts than those in the unmanured plants. (Table 7). With the exception of roots, both Ndff and Ndfs values in shoots, panicles, and whole plant sorghum, amended with FGM were lower than those in the control, indicating that the enhancement of total N uptake of sorghum (Table 8) mainly resulted from N released from FGM. However, amounts of soil or fertilizer N uptake in roots of the FGM treated plants were higher than those of the control, particularly in FGM₁₀₀ and FGM₁₅₀ treatments. Moreover, a positive effect on sorghum root dry matter yield was also observed following the addition of both GMs (Tables 4 and 5). The increase in the amount of soil N uptake by sorghum following the addition of plant residues as GMs, particularly PGM, was demonstrated in this study, a phenomenon highlighted previously by several authors [3, 30 - 32]. Jenkinson *et al.* [33] introduced the term “Added Nitrogen Interaction” or “priming effect” to describe any effect that the addition of N (*i.e.* organic N in the green manure) may have on the native soil N. Azam [34] reported that the extra nitrogen comes from soil organic matter as a result of the interaction of the added nitrogen. An increase in N availability from sources like soil organic matter could be attributed to a priming effect of the added nitrogen [34]. In other words, the added nitrogen interacts with the N already present in the soil, in a way to increase the availability of the later. Such an effect may result from an increase in root growth enabling the plants to explore a greater soil volume thus increasing the uptake of nutrients ‘like soil N’ by the green manured crops [33]. In this study, a positive effect on sorghum root dry matter yield was observed following the addition of both GMs. On the other hand, as suggested by Azam [31 - 34], there is a reason to believe that a priming effect may occur in soils because the endogenous soil microorganisms will react to the addition of energy-rich materials, and the increased microbial activity will involve mineralization process of the organic N in the GM. In view of our results, it can be concluded that the beneficial effects of incorporating pod shells in soil on sorghum nitrogen accumulation, may be attributed not only to the additional N released from GMs to the plants but also to their effects on the enhancement of soil N uptake which was more evident in PGM than FGM treatments [3, 12, 23].

4.4. Effect of Green Manuring with Pea and Faba Bean Pod Shells on N Recoveries (%NUE_{gm}) by Sorghum

Regardless of green manure used, nitrogen recoveries (%NUE_{gm}) in sorghum plants (uptake by plants) following the addition of pea or faba bean pod shells ranged from 29 to 45% indicating that both GM forms were used effectively. Release of N from pea GM which added to the soil 15 days before planting, seemed to be rapid with nearly half of N being utilized by the sorghum in PGM₅₀ and PGM₁₀₀, and third of N in PGM₁₅₀ treatments. For faba bean GM, around a quarter to third of its N content being used by the sorghum plants. Nearly, the same range of %NUE_{gm} values (20-52%) has been reported for sorghum manured with sesbania [3], Russian olive [12], or

leucaena [23] leaves. However, the efficient use of green manure by sorghum plants was higher in treatment with pea than those with faba bean residues. It is well documented that organic nitrogen in the GM is released into the soil after incorporation through the process of mineralization by soil microorganisms [27]. The rate of this process is affected by many factors such as temperature, moisture, quantity, and quality (*i.e.* chemical composition) of the GM residue [4, 30, 35]. The ratio of carbon to nitrogen (C/N) is a useful guide for the decomposition and mineralization of the organic nitrogen in the added GM [4]. Abdelhamid *et al.* [36] reported that the lower the C/N the greater the mineralization rate. In this study, nitrogen recovery rates (uptake by sorghum) from pea pod shells were higher than those from faba bean probably because C/N of pea (15.2) was lower than that of faba bean (18). Moreover, Daimon [37] reported that not only the C/N ratio of the incorporated green manure involved in nitrogen release, but also other parameters such as lignin and polyphenols, ratio of lignin /N, ratio of polyphenols to total N, and ratio of (lignin + polyphenols) to total N are determining for GM quality regarding to N release after incorporation. Accordingly, GMs with a high ratio of these parameters are usually considered to compose more slowly than those of low ratio [37, 38]. In this study, all the above-mentioned parameters in PGM were lower than those in FGM (Table 2). Such results may interpret the beneficial effects of the former over the latter GM regarding growth and N uptake by sorghum plants. Moreover, it is worth mentioning that the difference in polyphenols and polyphenols /N values between the two forms of GM were much higher than those of the other analyzed parameters (Table 2). In other words, polyphenols and polyphenols /N mean values were 5 and 6 times higher in FGM than PGM, respectively. Whereas, other values were less high (*i.e.* 1.2, 1.2, 1.5, 1.4 times higher in FGM than PGM, for L, C/N, L+ph/N, and L/N, respectively), (Table 2). This result may indicate the importance of polyphenols and polyphenols /N criteria in N released from GM. Our observation agrees well with the results of Fan *et al.* [39] who found that the impact of leaf litter polyphenols concentration and polyphenols /N ratio on N release rate was stronger than that of the C/N or lignin/N ratio.

While green manuring has been demonstrated to have beneficial effects on many aspects of the cropping systems, the most obvious direct economic benefit from the utilization of GM in agriculture is N fertilizer saving. It is well known that legume GM can replace a portion of the fertilizer N requirements for the subsequent crops. For example, Kurdali [11] reported that 18-54 kg N ha⁻¹ of inorganic N fertilizer may be saved as a result of using dhaincha (*Sesbania aculeata* Pers.) plant residues as GM for sorghum growth. In this study, amounts of added nitrogen from GMs to sorghum were equivalent to 50, 100, and 150 kg N ha⁻¹ (Table 3). The use of PGM or FGM as green manures could substitute significant amounts of N fertilizer. For example, when PGM was used as a GM, recoveries of the added N by sorghum (%NUE_{gm}) were 45, 42, and 32% (Table 9), for the above-mentioned treatments, respectively. Thus, equivalent amounts to 23, 42, and 48 kg N ha⁻¹ of inorganic fertilizer may be saved using PGM as green manure for sorghum growth; whereas, 17, 32, and 43 kg N ha⁻¹ of inorganic N fertilizer may be saved as a result of using

FGM.

Overall, this study demonstrated that the organic wastes of pea and faba bean pod shells applied to soil as GMs increased its fertility, like inorganic N fertilizer, and improved the growth of sorghum by enhancing nutrients uptake from soil and from organic sources. The use of these organic wastes as GMs is

very useful to reduce chemical fertilizer application in cropping systems, maintain a sustainable N and P supply for crops and can be effectively used for soil rehabilitation and plant growth enhancement. In light of this study, the importance of using pea and faba bean pod shells wastes as green manures for plant growth enhancement is summarized in Fig. (2).




Fig. (2). Scheme representing the importance of using pea or faba bean pod shells ‘after seed removals’ as green manure in N fertilizer saving and in percent increments of dry matter, N and P uptake of different plant parts of sorghum, as compared with the un-manured control treatment. Data represents the lowest and the highest value of increments according to applied N in the form of GM. The lowest values belonged to N50 treatments that were higher in PGM than FGM. The highest values belonged either to N100 or to N150.

CONCLUSION

To the best of our knowledge, the present study is the first report on the use of organic wastes of pea and faba bean pod shells to be used as green manures for the growth of sorghum. Based on the data of all parameters in the research results, it can be concluded that soil amendment with these agricultural wastes has a significant influence on the productivity and nutrient uptake (*i.e.* N and P) in sorghum plants. Our results also indicated that PGM or FGM may substitute 17 to 48 kg N ha⁻¹ of inorganic fertilizer. The use of such agricultural by-products as green manure is an additional way of handling the agricultural wastes in an organic farming system.

LIST OF ABBREVIATIONS

GM = Green Manure

PGM = Pea pod shells used as Green Manure

FGM = Faba bean pod shells used as Green Manure

Ndff = Nitrogen derived from fertilizer

Ndfs = Nitrogen derived from soil

Ndf_{gm} = Nitrogen derived from green manure

NUE_{gm} = Nitrogen Use Efficiency of added green manure

DM = Dry Matter

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

HUMAN AND ANIMAL RIGHTS

No animals/humans were used for studies that are the base of this research.

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The authors would like to thank Professor Ibrahim Othman, Director General of the Atomic Energy Commission of Syria (AECS), for his support. Thanks are also extended to research staff members of the agriculture department of AECS for their technical assistance when this work was carried out.

REFERENCES

- [1] Kim SY, Lee CH, Gutierrez J, Kim PJ. Contribution of winter cover crop amendments on global warming potential in rice paddy soil during cultivation. *Plant Soil* 2013; 366(1-2): 273-86. [http://dx.doi.org/10.1007/s11104-012-1403-4]
- [2] Fageria NK. Green manuring in crop production. *J Plant Nutr* 2007; 30(5): 691-719. [http://dx.doi.org/10.1080/01904160701289529]
- [3] Kurdali F, Al-Ain F, Al-Shamma'a M, Razzouk AK. Performance of sorghum grown on a salt affected soil manured with Dhaincha plant residues using ¹⁵N isotopic dilution technique. *J Plant Nutr* 2007; 30: 1605-21. [http://dx.doi.org/10.1080/01904160701615491]
- [4] Giller KE, Wilson J. Nitrogen fixation in tropical cropping systems. Oxon, UK: CAB International 1993; p. 313.
- [5] Aggarwal RK, Kumar P, Power JF. Use of crop residue and manure to conserve water and to enhance nutrient availability of pearl millet yield in an arid tropical region. *Soil Tillage Res* 1997; 41: 43-51. [http://dx.doi.org/10.1016/S0167-1987(96)01082-3]
- [6] Turgut I, Bilgili U, Duman A, Acikgoz E. Effect of green manuring on the yield of sweet corn. *Agron Sustain Dev* 2005; 25(4): 433-8. [http://dx.doi.org/10.1051/agro:2005044]
- [7] Eusuf Zai AK, Horiuchi T, Matsui T. Effects of compost and green manure of pea and their combinations with chicken manure and rapeseed oil residue on soil fertility and nutrient uptake in wheat-rice cropping system. *Afr J Agric Res* 2008; 3(9): 633-9.
- [8] Lupwayi NZ, Haque I. Mineralization of N, P, K, Ca and Mg from *Sesbania* and *Leucaena* leaves varying in chemicals composition. *Soil Biol Biochem* 1998; 30(3): 337-43. [http://dx.doi.org/10.1016/S0038-0717(97)00132-6]
- [9] Bah AR, Zaharah AR, Hussin A. Phosphorus uptake from green manures and phosphate fertilizers applied in an acid tropical soil. *Commun Soil Sci Plant Anal* 2006; 37(13-14): 2077-93. [http://dx.doi.org/10.1080/00103620600770433]
- [10] Hood RC, Merckx R, Jensen ES, Powelson D, Matijevic M, Hardarson G. Estimating crop N uptake from organic residues using a new approach to the ¹⁵N isotope dilution technique. *Plant Soil* 2000; 223(1-2): 33-46. [http://dx.doi.org/10.1023/A:1004789103949]
- [11] Kurdali F. Estimates of dry matter yield and N uptake in sorghum grown on saline and non-saline soils manured with dhaincha plant residues. *J Plant Nutr* 2004; 27(9): 1611-33. [http://dx.doi.org/10.1081/PLN-200026004]
- [12] Al-Ain F, Al-Chamma'a M, Kurdali F. Growth and nitrogen uptake in sorghum plants manured with *Elaeagnus angustifolia* leaves as affected by alternate irrigation with saline and non-saline water using ¹⁵N. *Open Agric J* 2017; 11: 24-34. [http://dx.doi.org/10.2174/1874331501711010024]
- [13] Hawtin GC, Muehlbauer FJ, Slinkard AE, Singh KB. Current status of cool season food legume crop improvement: An assessment of critical needs. *World Crops: Cool Season Food Legumes*. Dordrecht, The Netherlands: Kluwer Academic Publishers 1988; pp. 67-80. [http://dx.doi.org/10.1007/978-94-009-2764-3_8]
- [14] Mateos-Aparicio I, Redondo-Cuenca A, Villanueva-Suárez M-J, Zapata-Revilla M-A, Tenorio-Sanz MD. Pea pod, broad bean pod and okara, potential sources of functional compounds. *Lebensm Wiss Technol* 2010; 43(9): 1467-70. [http://dx.doi.org/10.1016/j.lwt.2010.05.008]
- [15] Wadhwa M, Kaushal S, Bakshi MPS. Nutritive evaluation of vegetable wastes as complete feed for goat bucks. *Small Rumin Res* 2006; 64(3): 279-84. [http://dx.doi.org/10.1016/j.smallrumres.2005.05.017]
- [16] Bath B, Ekblad G, Ascard J, Olsson K, Andersson B. Yield and nitrogen uptake in organic potato production with green manures as pre-crop and the effect of supplementary fertilization with fermented slurry. *Biol Agric Hort* 2006; 24(2): 135-48. [http://dx.doi.org/10.1080/01448765.2006.9755015]
- [17] Fall D, Bakhoun N, Fall F, et al. Effect of peanut shells amendment on soil properties and growth of seedlings of *Senegalia senegal* (L.) Britton, *Vachellia seyal* (Delile) P. Hurter, and *Prosopis juliflora* (Swartz) DC in salt-affected soils. *Ann For Sci* 2018; 75(1) [http://dx.doi.org/10.1007/s13595-018-0714-x]
- [18] Constantinides M, Fownes JH. Nitrogen mineralization from leaves and litter of tropical plants; relationship to nitrogen, lignin and soluble polyphenol concentrations. *Soil Biol Biochem* 1994; 26(1): 49-55. [http://dx.doi.org/10.1016/0038-0717(94)90194-5]
- [19] Rowland AP, Roberts JD. Lignin and cellulose fractionation in decomposition studies using acid-detergent fibre methods. *Commun Soil Sci Plant Anal* 1994; 25(3-4): 269-77. [http://dx.doi.org/10.1080/00103629409369035]
- [20] Kumarasinghe KS, Eskew DL. Comparison of direct and indirect 15N methods for evaluation of N uptake by rice from azolla isotopic studies of azolla and nitrogen fertilization of rice. Dordrecht, The Netherlands: Kluwer Academic Publishers 1993; pp. 16-21. [http://dx.doi.org/10.1007/978-94-011-1681-7_3]
- [21] Kouyaté Z, Franzluebbers K, Juo ASR, Hossner LR. Tillage, crop residue, legume rotation, and green manure effects on sorghum and millet yields in the semiarid tropics of Mali. *Plant Soil* 2000; 225(1-2): 141-51. [http://dx.doi.org/10.1023/A:1026589528352]
- [22] Sweeney DW, Moyer JL. In-Season nitrogen uptake by grain sorghum following legume green manures in conservation tillage systems. *Agron J* 2004; 96(2): 510-5. [http://dx.doi.org/10.2134/agronj2004.5100]
- [23] Kurdali F, Al-Shamma'a M. Growth and N-uptake in sorghum plants manured with *Leucaena leucocephala* leaves as affected by nitrogen rate and time of application. *Commun Soil Sci Plant Anal* 2010; 41(3): 308-19. [http://dx.doi.org/10.1080/00103620903460823]
- [24] Getu A, Teshager A. Effect of adaptable green manure plants on sorghum yields and soil fertility in Eastern Amhara region of Ethiopia. *J Biol Agri Healthcare* 2015; 5(11): 223-31.
- [25] Azeez JO, Van Averbek W. Effect of manure types and period of incubation on phosphorus-sorption indices of a weathered tropical soil. *Commun Soil Sci Plant Anal* 2011; 42(8): 2200-18. [http://dx.doi.org/10.1080/00103624.2011.602452]
- [26] Requejo MI, Eichler-Lobermann B. Organic and inorganic phosphorus forms in soil as affected by long-term application of organic amendments. *Nutr Cycl Agroecosyst* 2014; 100(2): 245-55. [http://dx.doi.org/10.1007/s10705-014-9642-9]
- [27] Baddeley JA, Pappa VA, Pristeri A, et al. Legume-based green manure crops. In: Murphy-Bokern D, Stoddard FL, Watson CA, Eds. *Legumes in cropping systems*. CAB International 2017; pp. 125-38. [http://dx.doi.org/10.1079/9781780644981.0125]
- [28] Gao X, Shi D, Lv A, et al. Increase phosphorus availability from the use of alfalfa (*Medicago sativa* L.) green manure in rice (*Oryza sativa* L.) agroecosystem. *Sci Rep* 2016; 6: 36981. [http://dx.doi.org/10.1038/srep36981] [PMID: 27833163]
- [29] Pypers P, Huybrighs M, Diels J, Abaidoo R, Smolders E, Merckx R. Does the enhanced P acquisition by maize following legumes in a rotation result from improved soil P availability? *Soil Biol Biochem* 2007; 39(10): 2555-66. [http://dx.doi.org/10.1016/j.soilbio.2007.04.026]
- [30] Rees RM, Yan L, Ferguson M. The release and plant uptake of nitrogen from some plant and animal manures. *Biol Fertil Soils* 1993;

- 15(4): 285-93.
[<http://dx.doi.org/10.1007/BF00337214>]
- [31] Azam F. Comparative effects of organic and inorganic nitrogen sources applied to a flooded soil on rice yield and availability of N. *Plant Soil* 1990; 125(2): 255-62.
[<http://dx.doi.org/10.1007/BF00010664>]
- [32] Manguiat IJ, Singleton PW, Rocamora PM, Calo MU, Taleon EE. Effectiveness of *Sesbania rostrata* and *Phaseolus calcaratus* as green manure for upland rice grown in acidic soil. *Plant Soil* 1997; 192(2): 321-31.
[<http://dx.doi.org/10.1023/A:1004242803894>]
- [33] Jenkinson DS. Interactions between fertilizer nitrogen and soil nitrogen-so called priming effect. *J Soil Sci* 1985; 36: 425-44.
[<http://dx.doi.org/10.1111/j.1365-2389.1985.tb00348.x>]
- [34] Azam F. Added nitrogen interaction in the soil plant system. A review. *Pak J Agron* 2002; 1(1): 54-9.
[<http://dx.doi.org/10.3923/ja.2002.54.59>]
- [35] Cadisch G, Handayanto E, Malama C, Seyni F, Giller KE. N recovery from legume prunings and priming effects are governed by the residue quality. *Plant Soil* 1998; 205(2): 125-34.
[<http://dx.doi.org/10.1023/A:1004365217018>]
- [36] Abdelhamid M, Horiuchi T, Oba S. Nitrogen uptake by faba bean from ¹⁵N-labelled oil seed-rape residue and chicken manure with ryegrass as a reference crop. *Plant Prod Sci* 2004; 7(4): 371-6.
[<http://dx.doi.org/10.1626/pp.s.7.371>]
- [37] Daimon H. Traits of genus *Crotalaria* uses as a green manure legume on sustainable cropping system. *Jpn Agric Res Q* 2006; 40(4): 299-305.
[<http://dx.doi.org/10.6090/jarq.40.299>]
- [38] Palm CA, Sanchez PA. Nitrogen release from the leaves of some tropical legumes as affected by their lignin and polyphenolic contents. *Soil Biol Biochem* 1991; 23(1): 83-8.
[[http://dx.doi.org/10.1016/0038-0717\(91\)90166-H](http://dx.doi.org/10.1016/0038-0717(91)90166-H)]
- [39] Fan D, Fan K, Yu C, Lu Y, Wang X. Tea polyphenols dominate the short-term tea (*Camellia sinensis*) leaf litter decomposition. *J of Zhejiang Uni-Sci B (Biomedicine & Biotech)* 2017; 18(2): 99-108.

© 2019 Al-Chammaa *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: (<https://creativecommons.org/licenses/by/4.0/legalcode>). This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.