




# The Open Biology Journal

Content list available at: <https://openbiologyjournal.com/>



## REVIEW ARTICLE

### New Record of the Giant Timber Flies (Diptera: Pantophthalmidae) from Sartenejas Valley, Miranda, Venezuela

Humberto Carvajal-Chitty<sup>1,\*</sup> and Aivlé C. Cabrera G.<sup>2</sup>

<sup>1</sup>Departamento de Estudios Ambientales, Universidad Simón Bolívar, Valle de Sartenejas, Baruta, Edo. Miranda, Apartado 89000. Caracas, Venezuela

<sup>2</sup>Departamento de Química, Universidad Simón Bolívar, Valle de Sartenejas, Baruta, Edo. Miranda, Apartado 89000. Caracas, Venezuela

#### Abstract:

Giant timber fly *Pantophthalmus bellardii* (Austen 1923) have been collected at 1200 meters above sea level, at Sartenejas Valley, a campus of the Universidad Simón Bolívar, Miranda State, Venezuela. Giant timber flies are very difficult to find in nature and rarer yet in the urban environments. Their biology and interactions with environmental variables are virtually unknown. In Venezuela, these flies have been observed from 100 to 1100 meters above sea level. This record extends the geographical distribution of this rare timber fly within Venezuelan territory. An updated distribution map for *P. bellardii* in Venezuela is provided.

**Keywords:** Pantophthalmidae, Diptera, Giant, Flies, Venezuela, *Pantophthalmus*, Neotropical.

#### Article History

Received: March 14, 2019

Revised: June 22, 2019

Accepted: July 04, 2019

## 1. INTRODUCTION

**Pantophthalmidae** or giant timber flies are among the largest Diptera, with wingspan up to 10 cm [1 - 3]. Giant timber flies (Diptera: Pantophthalmidae) are very rare [1] and are endemic in Neotropical region [2]. Actually, only 20 valid species are recognized in two genera: *Opetiops* (Enderlei, 1921) and *Pantophthalmus* (Thunberg 1919) [2,3]. Traditionally, large size has been the most notable characteristic trait of the Pantophthalmidae.

Giant timber flies are living on dead or still alive trees [2] on which their larvae feed of accumulated organic matter or the wood fermentation products. The adults emerge for the reproduction only and are aphagus (do not feed past larval stage).

Their regional geographical distribution has been covered by Val [4] and recently Amat [5] has added important data about Pantophthalmidae distribution in Mexico and their biology in the Red Oak forest (for *Pantophthalmus roseni* (Enderlein 1931) within Mexico. Pujol-Luz and Souto Morgado [2] offer new data about *Pantophthalmus pictus* (Wiedemann 1821) records from central Brazil. Despite the

recent progress on giant timber flies studies, very little information is available regarding their distribution in Venezuela.

Here we are reporting a rare case of the Pantophthalmidae urban record in Venezuela. Such record was made for the first time for *Pantophthalmus bellardii* for Sartenejas Valley (Miranda, Venezuela) at 1.200 meters above sea level [3], at the campus of Universidad Simón Bolívar (Fig. 1).

### 1.1. Collected Material

The University is inside a valley surrounded by evergreen primary forest. The main campus is encircled by the plantations of *Pinus caribea*, *P. pseudostrobus* and *P.patula* [6]. Nowadays, it is a very stable conifer forest that keeps the climate constant and protects the original forest.

Two specimens of *Pantophthalmus bellardii* (Austen 1923) [3], a male and a female, (Fig. 2) were found dead on the roof of the Química y Procesos building (QYP) at the central area of the campus.


The genus identification was done by Dr. Torsten Dikow, (Research Entomologist for Diptera of Smithsonian Institution Natural Museum of Natural History) and Val [4] was used for the final identification of the species.

The female was a 4.5 cm long (from tip of head to tip ovipositor) (Fig. 3), curve ovipositor; 3.6 cm length (Fig. 4).

\* Address correspondence to this author at the Departamento de Estudios Ambientales, Universidad Simón Bolívar, Valle de Sartenejas, Baruta, Edo. Miranda, Apartado 89000. Caracas, Venezuela; Tel: +58-424-1731315; E-mail: [hchitty@usb.ve](mailto:hchitty@usb.ve)

Abdomen coloration varies from black (female) to pale orange (male). Both specimens have beige parallel lines at the mesonotum. As it was noted by Val [4], the female is larger than the male.

The presence of *P. bellardii* at Sartenejas Valley (Miranda Venezuela) at 1200 meters above sea level, is an important extension of the altitudinal profile of the Panthophthalmidae records in Venezuela. It is also the first record of this family for the Miranda State.



**Fig. (1).** Venezuela map with the location of the Universidad Simón Bolívar where the specimens were found.



**Fig. (2).** Dorsal view of the male (upper) and female (lower) of *P. bellardii* collected at Sartenejas Valley, Miranda Venezuela.



**Fig. (3).** Female coloration and characteristics of *P. bellardii*.



Fig. (4). Male coloration and characteristics of *P. bellardii*.



Fig. (5). Location of the current record of *Pantophthalmus bellardii* in Northern Venezuela (see legend inside the map).

## 1.2. Summary of the *Pantophthalmus bellardii* records in Venezuela

Pantophthalmidae are currently reported from only a handful of sites in Venezuela. Báez [1] recorded *P. bellardii* in Choroni (100 meters above sea level, Aragua State), El Limón (450 meters above sea level, Aragua), Ocumare de La Costa (400 meters above sea level, Aragua), Rancho Grande (1100 meters above sea level, Aragua, Henri Pittier National Park) and Sierra de Perijá (800 meters above sea level, Zulia). Fig. (5) shows the current updated location and distribution of this species along the country.

The current distribution records of the *Pantophthalmus bellardii* [2], [3] extends the current knowledge of the species distribution alongside the records from Colombia [5], Brazil [3] and Venezuela of the timber giant fly [1]. It is also the highest altitude above sea level at which the species have been encountered so far in Venezuela. This new record is important for the studies of the Giant Timber flies, which probably act as an important ecosystem engineers and forestry pests, facilitating organic matter flow in the Neotropical forests.

## CONSENT FOR PUBLICATION

Not applicable.

## FUNDING

None.

## CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

## ACKNOWLEDGEMENTS

We thank the identification to genus level to Dr. Torsten Dikow, (Research Entomologist for Diptera of Smithsonian Institution Natural Museum of Natural History) which kindly help us with this material. Also to Dr. Solange Issa for her comments and suggestions for the present note. To Dr. M. Báez (Departamento de Biología Animal, Edafología y Geología, Universidad de La Laguna, España) y Dr. José Clavijo (Facultad de Agronomía Universidad Central de Venezuela) for their help in the search for bibliography associated with the giant timber flies in Venezuela and Dr. Viktor Banarov for suggestions regarding the manuscript.

## REFERENCES

- [1] Báez M. Contribución al conocimiento de los Pantophthalmidae de Venezuela (Insecta, Diptera). Bol Ent Venez NS 1983; 2(18): 133-6.
- [2] Pujol-Luz JR, Souto Morgado G. 2019. New record of *Pantophthalmus pictus* (Wiedemann, 1821) (Diptera, Pantophthalmidae) in the Cerrado vegetation of central Brazil. Pap Avulsos Zool 2018; 58e20185828 [http://dx.doi.org/ 10.11606/1807- 0205 / 2018.58.28]
- [3] Austen EE. A revision of the family Pantophthalmidae (Diptera) with descriptions of new species and a new genus. Proc Zool Soc Lond 1923; 551-98. Available from: https:// www.ma p re ss.co m/j/ zt/article/view/zootaxa.4122.1.25
- [4] Val FC. Systematics and evolution of the Pantophthalmidae (Diptera, Brachycera). Arq Zool S Paulo 1976; 27(2): 51-164.

- [http://dx.doi.org/10.11606 / issn.2176 - 7793.v27i2p51-164]
- [5] Amat EC. New records of Timber flies (Diptera: Pantophthalmidae) from Colombia. *Entomotrópica* 2005; 20(2): 95-6. Available from: <http://www.bioline.org.br/pdf/em05017>
- [6] Baruch Z, Nozawa S. Abandoned coffee plantations: Biodiversity conservation or path for nonnative species? Case study in a neotropical montane forest. *Interciencia* 2014; 39(8): 554-61. Available from: <https://www.interciencia.net/wp-content/uploads/2017/11/554-Baruch-8.pdf>

---

© 2019 Carvajal-Chitty and Cabrera G.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: (<https://creativecommons.org/licenses/by/4.0/legalcode>). This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.