



# The Open Biotechnology Journal

Content list available at: [www.benthamopen.com/TOBIOTJ/](http://www.benthamopen.com/TOBIOTJ/)

DOI: 10.2174/1874070701812010123



## RESEARCH ARTICLE

# Experimental Study of Thermal Restraint in Bio-Protectant Disaccharides by FTIR Spectroscopy

S. Magazù<sup>1,2,\*</sup>, E. Calabrò<sup>1</sup> and M.T. Caccamo<sup>1,2</sup>

<sup>1</sup>Department of Mathematical and Informatics Sciences, Physical Sciences and Earth Sciences of Messina University, Viale Ferdinando Stagno D'Alcontres 31, 98166 Messina, Italy

<sup>2</sup>Istituto Nazionale di Alta Matematica "F. Severi" – INDAM - Gruppo Nazionale per la Fisica Matematica – GNFM, Messina, Italy

Received: December 14, 2017

Revised: June 5, 2018

Accepted: June 22, 2018

### Abstract:

#### Background:

In the present paper, InfraRed (IR) spectra on water mixtures of two homologous disaccharides, *i.e.* sucrose and trehalose, as a function of temperature have been collected.

#### Methods:

In particular, IR spectra were registered, in the spectral range from 4000 cm<sup>-1</sup> to 400 cm<sup>-1</sup>, to investigate the thermal response of the water mixtures of two homologous disaccharides, through positive thermal scans, *i.e.* by increasing the temperature from the value of 25°C to the value of 50°C. The OH-stretching region has been analyzed by means of two simple and straightforward procedures, *i.e.* by evaluating the shift of the intramolecular OH stretching center frequency and the Spectral Distance (SD).

#### Result and Conclusion:

Both the analyses indicate that trehalose water mixture have a higher thermal response than that of the sucrose-water mixture.

**Keywords:** Sucrose, Trehalose, Bio-protection, Temperature, FTIR spectroscopy, Infrared.


## 1. INTRODUCTION

It is well known that homologous disaccharides *i. e.* sucrose and trehalose although have the same chemical formula (C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>), they present different bio-protective properties. In particular, trehalose shows a higher bio-protective effectiveness in comparison with sucrose, playing a key role in cryptobiotic-activating substances [1 - 11]. Sucrose is constituted by a glucose ring in the  $\alpha$  configuration and a fructose ring in the  $\beta$  configuration; the  $\alpha$  and  $\beta$  structures of the same monosaccharide differ only in the orientation of the OH groups at some carbon atom in the ring itself [12 - 18]. Trehalose is a disaccharide of glucose constituted by two pyranose rings in the same configuration, linked by a glycosidic bond between the chiral carbon atoms C1 of the two rings [19 - 28].

In Fig. (1), the chemical structures of sucrose and trehalose are reported.

More precisely, trehalose, due to its bioprotective properties allows to many organisms, which synthesize it to undergo in a state of "suspended life" and to re-activate the vital functions when the external conditions come back favourable [29 - 42].

\* Address correspondence to this author at the Department of Mathematical and Informatics Sciences, Physical Sciences and Earth Sciences of Messina University, Viale Ferdinando Stagno D'Alcontres 31, 98166 Messina, Italy; Tel: +39 090 6765025; E-mail: [smagazu@unime.it](mailto:smagazu@unime.it)



**Fig. (1).** Chemical structure of sucrose (left) and chemical structure of trehalose (right).

Thus, such organisms can resist dehydration by means of the formation of a glass of intracellular carbohydrates, that, with its high viscosity and hydrogen bonding, is able to stabilize and protect the integrity of molecules. For these reasons, the trehalose can be employed to preserve red blood cells [43 - 55].

It should be stressed that trehalose possesses an unusual ability to stabilize biomolecules since it is able to stabilize the structure without interfering with the functionality of the protein [56 - 64].

Furthermore, the thermal properties of bioprotectant systems have been the subject of attention not only for the physical, biological and chemical research, but also for the food products and for the cultural heritage field [65 - 74]. Although a lot of studies have been addressed the anhydrobiosis and biopreservation phenomena; the molecular mechanisms are not yet clear [75 - 87]. As far as the employed spectroscopic technique is concerned, in our case, IR allows to characterize the system vibrational motions and hence the system structural arrangements and their changes with temperature. In this study, the Mid IR (MIR) region spanning from  $4000\text{ cm}^{-1}$  to  $400\text{ cm}^{-1}$  has been analyzed [88 - 92]. In particular, to highlight the thermal properties of these disaccharides mixtures, two straightforward approaches have been performed: the shift of the OH-stretching region ( $3700 - 3000\text{ cm}^{-1}$ ) as a function of temperature and the analysis of the temperature dependence of the Spectral Distance (SD). From the first approach, it is possible to see changes in the spectral band position while the SD reflects the structural changes in disaccharides mixtures. Both the approaches suggest that trehalose water mixture has a higher thermal restraint than sucrose-water mixture [93 - 97].

## 2. EXPERIMENTAL SETUP AND SAMPLE PREPARATION


Sucrose and trehalose were purchased from Aldrich-Chemie. Infrared spectra were collected in the  $25\div 50^\circ\text{C}$  temperature range, by means of the FTIR Vertex 70 V spectrometer (Bruker Optics) using Platinum diamond ATR. The investigated concentration value, expressed as weight fraction, *i.e.* (grams of disaccharide)/(grams of disaccharide+ grams of water), was: 0.30. For each spectrum, 32 scans were repeated at a resolution of  $4\text{ cm}^{-1}$  for the spectral range between  $4000$  and  $400\text{ cm}^{-1}$ . Such experimental conditions were kept constant for all the measurements. First to proceed with the spectra analysis, data pre-processing has been performed:

1. baseline treatment
2. smoothing treatment
3. first and second derivatives
4. normalization of spectra

Then, spectral analysis has been performed by means of the software MATLAB 2016a (Mathworks, Natick, USA) and the software Origin 9 (OriginLab Co., Northampton, USA).

## 3. EXPERIMENTAL DATA RESULTS AND DISCUSSION


Fig. (2a) reports, as an example, the IR spectra in the OH-stretching region ( $3700 < \Delta\omega < 3000\text{ cm}^{-1}$  spectral range) for different values of temperature, *i.e.*  $T=25^\circ\text{C}$ ,  $30^\circ\text{C}$ ,  $35^\circ\text{C}$ ,  $40^\circ\text{C}$ ,  $45^\circ\text{C}$  and  $50^\circ\text{C}$  for sucrose water mixture while Fig. (2b) shows, as an example, the IR spectra in the OH-stretching region ( $3700 < \Delta\omega < 3000\text{ cm}^{-1}$  spectral range) for different values of temperature, *i.e.*  $T=25^\circ\text{C}$ ,  $30^\circ\text{C}$ ,  $35^\circ\text{C}$ ,  $40^\circ\text{C}$ ,  $45^\circ\text{C}$  and  $50^\circ\text{C}$  for trehalose water mixture.



**Fig. (2).** a) OH-stretching region ( $3700 < \Delta\omega < 3000 \text{ cm}^{-1}$  spectral range) for different values of temperature, *i.e.* T=25°C, 30°C, 35°C, 40°C, 45°C and 50°C for sucrose water mixture and b) OH-stretching region ( $3700 < \Delta\omega < 3000 \text{ cm}^{-1}$  spectral range) for different values of temperature, *i.e.* T=25°C, 30°C, 35°C, 40°C, 45°C and 50°C for trehalose water mixture.

To quantitatively characterize the thermal response of the IR spectra, we have first evaluated the OH stretching center frequency shift and then the spectral distance.

In Fig. (3) the trend of the OH stretching center frequency shift for sucrose water mixture and for trehalose water mixture is reported. As it can be seen the sucrose in the temperature range  $25,0^\circ\text{C} < \Delta T < 50,0^\circ\text{C}$  changes the intramolecular band center frequency from  $3271 \text{ cm}^{-1}$  to  $3310 \text{ cm}^{-1}$ , while trehalose changes the intramolecular band center frequency from  $3264 \text{ cm}^{-1}$  to  $3285 \text{ cm}^{-1}$ .




**Fig. (3).** Trend of the OH stretching center frequency shift for sucrose water mixture and for trehalose water mixture.

Such a result suggests a higher thermal restraint for trehalose water mixture than sucrose water mixture. In this

framework, in order to extract a quantitative value and to perform a comparison between the two homologous disaccharides, an interpretative model making reference to the well known logistic function has been applied; such a procedure allows to characterize the thermal restraint of the system:


$$S(T) = A \left( 1 - \frac{1}{1 + e^{-B(T-T_0)}} \right) + (C - DT) \quad (1)$$

where A is the relaxation amplitude, B represents the relaxation stepness, T is the temperature value of the sigmoid inflection point and the contribution C-DT takes into account the low-temperature contribution. Furthermore, it is important to remember that the inverse of the relaxation amplitude is connected with the thermal restraint of the system. In Fig. (4), the fit, performed by means of equation (1), of OH-stretching band center frequency as a function of temperature for sucrose water mixture is shown.



**Fig. (4).** Intramolecular band center frequency versus temperature for sucrose water mixture. Blue diamonds: Experimental data; magenta continuous line: Fitting curve obtained by equation (1).

Fig. (5) reports the fit, performed by means of equation (1), of the OH-stretching band center frequency as a function of temperature for trehalose water mixture.



**Fig. (5).** Intramolecular band center frequency versus temperature for trehalose water mixture. Light blue circles: Experimental data; magenta continuous line: Fitting curve by equation (1).

What emerges from this analysis is that the data arrange themselves along an increasing sigmoidal curve providing a

higher thermal restraint, together with a higher  $T_0$  value for trehalose water mixture in respect to sucrose water mixture. More precisely, the value of thermal restraint for the sucrose-water mixture is equal to 0.024 that is lower than that of trehalose water solution whose thermal restraint value is 0.031. As far as the crossover temperature value is concerned, the value of trehalose water mixture, 37.32°C, is higher than that of sucrose water mixture, which is equal to 35.77°C. These results show that dynamics of trehalose water mixture is slower than sucrose water mixture; the trehalose has a low sensitivity to temperature changes.


In order to catch small changes in intramolecular band center frequency, the SD has been calculated by means of the following expression:

$$SD = (\sum_v [I(\omega T) - I(\omega, T = 25^\circ C)]^2 \Delta\omega)^{1/2} \quad (2)$$

where  $I(\omega, t)$  represents the absorbance as a function of frequency  $\omega$  and temperature  $T$ , while  $\Delta\omega$  is the frequency resolution instrument.

Such a method is able to evaluate the difference in the spectra collected at different temperatures from the spectrum measured at the lowest temperature, that in our case, equal to 25 °C.

In Fig. (6), the SD of sucrose water mixture (blue diamond) and trehalose water mixture (light blue circle) is shown.



**Fig. (6).** SD evaluated in the center frequency of sucrose water mixture (blue diamond) and of trehalose water mixture (light blue circle).

As it can be seen that the trend is linear; in order to obtain a quantitative value for the thermal response of the system, we have calculated the straight line slope. For the sucrose-water mixture, the straight line slope value is 0.02648 while for the trehalose water mixture, the slope value is equal to 0.02244.

This latter value suggests that trehalose is less sensitive to changes in temperature.

## CONCLUSION

The aim of this work was to characterize the thermal response for two homologous disaccharides, *i.e.* sucrose and trehalose in water mixtures by FTIR spectroscopy. The spectral range investigated spanned from 4000  $\text{cm}^{-1}$  to 400  $\text{cm}^{-1}$ , with the intramolecular OH vibrational contribution, covering the spectral range from 3000 to 3800  $\text{cm}^{-1}$ , in the

temperature range between 25°C and 50°C.

The registered spectra have been analyzed by means of two straightforward approaches, *i.e.* by evaluating both the shift of the intramolecular OH stretching center frequency and the Spectral Distance (SD). Although the two different approaches are simple, both the analyses furnish the same results confirming that trehalose water mixture has a higher thermal restraint with respect to the sucrose-water mixture.

The agreement between the results obtained by the two different approaches provides coherent information on the thermal stability of the disaccharide water mixtures investigated.

#### **ETHICS APPROVAL AND CONSENT TO PARTICIPATE**

Not applicable.

#### **HUMAN AND ANIMAL RIGHTS**

No animals/humans were used for studies that are the basis of this research.

#### **CONSENT FOR PUBLICATION**

Not applicable.

#### **CONFLICT OF INTEREST**

The authors declare no conflict of interest, financial or otherwise.

#### **ACKNOWLEDGEMENTS**

Declared none.

#### **REFERENCES**

- [1] Crowe JH, Carpenter JF, Crowe LM. The role of vitrification in anhydrobiosis. *Annu Rev Physiol* 1998; 60: 73-103. [<http://dx.doi.org/10.1146/annurev.physiol.60.1.73>] [PMID: 9558455]
- [2] Crowe JH, Crowe LM, Oliver AE, Tsvetkova N, Wolkers W, Tablin F. The trehalose myth revisited: Introduction to a symposium on stabilization of cells in the dry state. *Cryobiology* 2001; 43(2): 89-105. [<http://dx.doi.org/10.1006/cryo.2001.2353>] [PMID: 11846464]
- [3] Green JL, Angell CA. Phase relations and vitrification in saccharide-water solutions and the trehalose anomaly. *J Phys Chem* 1989; 93: 2880-2. [<http://dx.doi.org/10.1021/j100345a006>]
- [4] Magazù S, Migliardo F, Benedetto A, Calabrò E, La Torre R, Caccamo MT. Bioprotective effects of sucrose and trehalose on proteins in Sucrose: Properties. *Biosynthesis and Health Implications* 2013; pp. 35-46.
- [5] Chen T, Fowler A, Toner M. Literature review: Supplemented phase diagram of the trehalose-water binary mixture. *Cryobiology* 2000; 40(3): 277-82. [<http://dx.doi.org/10.1006/cryo.2000.2244>] [PMID: 10860627]
- [6] Kandror O, DeLeon A, Goldberg AL. Trehalose synthesis is induced upon exposure of *Escherichia coli* to cold and is essential for viability at low temperatures. *Proc Natl Acad Sci USA* 2002; 99(15): 9727-32. [<http://dx.doi.org/10.1073/pnas.142314099>] [PMID: 12105274]
- [7] Baptista RP, Pedersen S, Cabrita GJ, Otzen DE, Cabral JM, Melo EP. Thermodynamics and mechanism of cutinase stabilization by trehalose. *Biopolymers* 2008; 89(6): 538-47. [<http://dx.doi.org/10.1002/bip.20926>] [PMID: 18213692]
- [8] Sola-Penna M, Meyer-Fernandes JR. Stabilization against thermal inactivation promoted by sugars on enzyme structure and function: Why is trehalose more effective than other sugars? *Arch Biochem Biophys* 1998; 360(1): 10-4. [<http://dx.doi.org/10.1006/abbi.1998.0906>] [PMID: 9826423]
- [9] Magazù S, Migliardo F, Parker SF. Vibrational properties of bioprotectant mixtures of trehalose and glycerol. *J Phys Chem B* 2011; 115(37): 11004-9. [<http://dx.doi.org/10.1021/jp205599a>] [PMID: 21830802]
- [10] Iannelli E, Tettamanti E, Galantini L, Magazù S. An integrated quasi-elastic light-scattering, pulse-gradient-spin-echo study on the transport properties of  $\alpha,\alpha$ -trehalose, sucrose, and maltose deuterium oxide solutions. *J Phys Chem B* 2001; 105(48): 12143-9. [<http://dx.doi.org/10.1021/jp011275j>]
- [11] Tanaka M, Machida Y, Niu S, *et al.* Trehalose alleviates polyglutamine-mediated pathology in a mouse model of Huntington disease. *Nat Med* 2004; 10(2): 148-54.

- [http://dx.doi.org/10.1038/nm985] [PMID: 14730359]
- [12] Magazù S, Migliardo F, Mondelli C, Vadalà M. Correlation between bioprotective effectiveness and dynamic properties of trehalose-water, maltose-water and sucrose-water mixtures. *Carbohydr Res* 2005; 340(18): 2796-801. [http://dx.doi.org/10.1016/j.carres.2005.09.026] [PMID: 16246316]
- [13] Lebrét A, Bordat P, Affouard F, Descamps M, Migliardo F. How homogeneous are the trehalose, maltose, and sucrose water solutions? An insight from molecular dynamics simulations. *The Journal of Physical Chemistry B* 2005; 109(21): 11046-57.
- [14] Magazù S, Migliardo F, Mondelli C, Romeo G. Temperature dependence of mean square displacement by IN13: A comparison between trehalose and sucrose water mixtures. *Chem Phys* 2003; 292: 247-51. [http://dx.doi.org/10.1016/S0301-0104(03)00101-0]
- [15] O'Brien J. Stability of trehalose, sucrose and glucose to nonenzymatic browning in model systems. *J Food Sci* 1996; 61: 679-82. [http://dx.doi.org/10.1111/j.1365-2621.1996.tb12180.x]
- [16] Liu Q, Schmidt RK, Teo B, Karplus PA, Bray JW. Molecular dynamics studies of the hydration of  $\alpha,\alpha$ -trehalose. *J Am Chem Soc* 1997; 119: 7851-62. [http://dx.doi.org/10.1021/ja970798v]
- [17] Affouard F, Bordat P, Descamps M, *et al.* A combined neutron scattering and simulation study on bioprotectant systems. *Chem Phys* 2005; 317: 258-66. [http://dx.doi.org/10.1016/j.chemphys.2005.05.033]
- [18] Arai C, Kohguchi M, Akamatsu S, *et al.* Trehalose suppresses lipopolysaccharide-induced osteoclastogenesis bone marrow in mice. *Nutr Res* 2001; 21(7): 993-9. [http://dx.doi.org/10.1016/S0271-5317(01)00315-3] [PMID: 11446983]
- [19] Minutoli L, Altavilla D, Bitto A, *et al.* Trehalose: A biophysics approach to modulate the inflammatory response during endotoxic shock. *Eur J Pharmacol* 2008; 589(1-3): 272-80. [http://dx.doi.org/10.1016/j.ejphar.2008.04.005] [PMID: 18555988]
- [20] Pagnotta SE, Ricci MA, Bruni F, McLain S, Magazù S. Water structure around trehalose. *Chem Phys* 2008; 345(2-3): 159-63. [http://dx.doi.org/10.1016/j.chemphys.2007.07.052]
- [21] Magazù S, Migliardo F, Telling MTF. Structural and dynamical properties of water in sugar mixtures. *Food Chem* 2008; 106(4): 1460-6. [http://dx.doi.org/10.1016/j.foodchem.2007.05.097]
- [22] Magazù S, Migliardo F, Barreca D, Bellocco E, Laganà G. Aggregation processes of biomolecules in presence of trehalose. *J Mol Struct* 2007; 840: 114-8. [http://dx.doi.org/10.1016/j.molstruc.2006.11.028]
- [23] Calabrò E, Magazù S. On the bioprotective effectiveness of trehalose against man-made electromagnetic fields. *Adv Chem Res* 2015; 24: 25-50.
- [24] Magazù S, Migliardo F, Ramirez-Cuesta AJ. Concentration dependence of vibrational properties of bioprotectant/water mixtures by inelastic neutron scattering. *J R Soc Interface* 2007; 4(12): 167-73. [http://dx.doi.org/10.1098/rsif.2006.0162] [PMID: 17018423]
- [25] Magazù S, Migliardo F, Telling MTF. Study of the dynamical properties of water in disaccharide solutions. *Eur Biophys J* 2007; 36(2): 163-71. [http://dx.doi.org/10.1007/s00249-006-0108-0] [PMID: 17109123]
- [26] Magazù S, Migliardo F, Ramirez-Cuesta AJ. Changes in vibrational modes of water and bioprotectants in solution. *Biophys Chem* 2007; 125(1): 138-42. [http://dx.doi.org/10.1016/j.bpc.2006.07.003] [PMID: 16887256]
- [27] Minutoli L, Altavilla D, Bitto A, *et al.* The disaccharide trehalose inhibits proinflammatory phenotype activation in macrophages and prevents mortality in experimental septic shock. *Shock* 2007; 27(1): 91-6. [http://dx.doi.org/10.1097/01.shk.0000235092.76292.bc] [PMID: 17172986]
- [28] Magazù S, Migliardo F, Telling MTF.  $\alpha,\alpha$ -trehalose-water solutions. VIII. Study of the diffusive dynamics of water by high-resolution quasi elastic neutron scattering. *J Phys Chem B* 2006; 110(2): 1020-5. [http://dx.doi.org/10.1021/jp0536450] [PMID: 16471637]
- [29] Branca C, Maccarrone S, Magazù S, Maisano G, Bennington SM, Taylor J. Tetrahedral order in homologous disaccharide-water mixtures. *J Chem Phys* 2005; 122(17): 174513. [http://dx.doi.org/10.1063/1.1887167] [PMID: 15910051]
- [30] Magazù S, Migliardo F, Ramirez-Cuesta AJ. Inelastic neutron scattering study on bioprotectant systems. *J R Soc Interface* 2005; 2(5): 527-32. [http://dx.doi.org/10.1098/rsif.2005.0059] [PMID: 16849211]
- [31] Crowe LM. Lessons from nature: The role of sugars in anhydrobiosis. *Comp Biochem Physiol A Mol Integr Physiol* 2002; 131(3): 505-13. [http://dx.doi.org/10.1016/S1095-6433(01)00503-7] [PMID: 11867276]
- [32] Magazù S, Maisano G, Migliardo F, Mondelli C.  $\alpha,\alpha$ -trehalose/water solutions. VII: An elastic incoherent neutron scattering study on fragility. *J Phys Chem B* 2004; 108(36): 13580-5. [http://dx.doi.org/10.1021/jp035973a]

- [33] Magazù S, Maisano G, Migliardo F, Mondelli C. Mean-square displacement relationship in bioprotectant systems by elastic neutron scattering. *Biophys J* 2004; 86(5): 3241-9.  
[[http://dx.doi.org/10.1016/S0006-3495\(04\)74372-6](http://dx.doi.org/10.1016/S0006-3495(04)74372-6)] [PMID: 15111437]
- [34] Magazu S, Migliardo F, Mondelli C. Elastic incoherent neutron scattering from homologous disaccharides/H<sub>2</sub>O mixtures. *J Chem Phys* 2004; 119(24): 13033-8.  
[<http://dx.doi.org/10.1063/1.1625369>]
- [35] Abate L, Blanco I, Branca C, *et al.* Homologous disaccharide properties at low temperatures. *J Mol Liq* 2003; 103-104(SPEC.): 177-80.  
[[http://dx.doi.org/10.1016/S0167-7322\(02\)00137-X](http://dx.doi.org/10.1016/S0167-7322(02)00137-X)]
- [36] Magazù S, Migliardo F, Vertessy BG, Caccamo MT. Investigations of homologous disaccharides by elastic incoherent neutron scattering and wavelet multiresolution analysis. *Chem Phys* 2013; 424: 56-61.  
[<http://dx.doi.org/10.1016/j.chemphys.2013.05.004>]
- [37] Migliardo F, Caccamo MT, Magazù S. Elastic incoherent neutron scatterings wavevector and thermal analysis on glass-forming homologous disaccharides. *J Non-Cryst Solids* 2013; 378: 144-51.  
[<http://dx.doi.org/10.1016/j.jnoncrysol.2013.06.030>]
- [38] Marchese N, Cannuli A, Caccamo MT, Pace C. 2016 New generation non-stationary portable neutron generators for biophysical applications of neutron activation analysis *Biochimica et Biophysica Acta (BBA) BBAGEN-28495 1861 Issue 1 Part B 3661-3670*.  
[<http://dx.doi.org/10.1016/j.bbagen.2016.05.023>]
- [39] Magazù S, Villari V, Faraone A, Maisano G, Heenan RK, King S.  $\alpha$ , $\alpha$ -trehalose-water solutions VI. A view of the structural and dynamical properties of O $\beta$ G micelles in the presence of trehalose. *J Phys Chem B* 2002; 106(27): 6954-60.  
[<http://dx.doi.org/10.1021/jp020698y>]
- [40] Branca C, Magazù S, Maisano G, Migliardo F. Vibrational and relaxational contributions in disaccharide/H<sub>2</sub>O glass formers. *Phys Rev B* 2001; 64(22): 224204.  
[<http://dx.doi.org/10.1103/PhysRevB.64.224204>]
- [41] Elbein AD, Pan YT, Pastuszak I, Carroll D. New insights on trehalose: A multifunctional molecule. *Glycobiology* 2003; 13(4): 17R-27R.  
[<http://dx.doi.org/10.1093/glycob/cwg047>] [PMID: 12626396]
- [42] Garg AK, Kim J-K, Owens TG, *et al.* Trehalose accumulation in rice plants confers high tolerance levels to different abiotic stresses. *Proc Natl Acad Sci USA* 2002; 99(25): 15898-903.  
[<http://dx.doi.org/10.1073/pnas.252637799>] [PMID: 12456878]
- [43] Higashiyama T. Novel functions and applications of trehalose. *Pure Appl Chem* 2002; 74: 1263-9.  
[<http://dx.doi.org/10.1351/pac200274071263>]
- [44] Kilburn D, Townrow S, Meunier V, Richardson R, Alam A, Ubbink J. Organization and mobility of water in amorphous and crystalline trehalose. *Nat Mater* 2006; 5(8): 632-5.  
[<http://dx.doi.org/10.1038/nmat1681>] [PMID: 16845422]
- [45] Chen Q, Haddad GG. Role of trehalose phosphate synthase and trehalose during hypoxia: From flies to mammals. *J Exp Biol* 2004; 207(Pt 18): 3125-9.  
[<http://dx.doi.org/10.1242/jeb.01133>] [PMID: 15299033]
- [46] Magazù S, Migliardo F, Benedetto A, Mondelli C, Gonzalez MA. Thermal behaviour of hydrated lysozyme in the presence of sucrose and trehalose by EINS. *J Non-Cryst Solids* 2011; 357(2): 664-70.  
[<http://dx.doi.org/10.1016/j.jnoncrysol.2010.06.075>]
- [47] Migliardo F, Caccamo MT, Magazù S. Thermal analysis on bioprotectant disaccharides by elastic incoherent neutron scattering. *Food Biophys* 2014; 9(2): 99-104.  
[<http://dx.doi.org/10.1007/s11483-013-9322-3>]
- [48] Caccamo MT, Cannuli A, Calabrò E, Magazù S. Acoustic levitator power device: Study of ethylene-glycol water mixtures IOP conference series: Materials science and engineering. 2017; 199.(1) Article number: 012119  
[<http://dx.doi.org/10.1088/1757-899X/199/1/012119>]
- [49] Magazu S, Villari V, Migliardo P, Maisano G, Telling MTF. Diffusive dynamics of water in the presence of homologous disaccharides: A comparative study by quasi elastic neutron scattering. IV *J Phys Chem B* 2001; 105(9): 1851-5.  
[<http://dx.doi.org/10.1021/jp002155z>]
- [50] Branca C, Magazù S, Maisano G, Migliardo P. Experimental study of the hydration properties of homologous disaccharides. *J Biol Phys* 2000; 26(4): 295-306.  
[<http://dx.doi.org/10.1023/A:1010344122760>] [PMID: 23345728]
- [51] Ballone P, Marchi M, Branca C, Magazù S. Structural and vibrational properties of trehalose: A density functional study. *J Phys Chem B* 2000; 104(26): 6313-7.  
[<http://dx.doi.org/10.1021/jp994346b>]
- [52] Lokotosh TV, Magazù S, Maisano G, Malomuzh NP. Nature of self-diffusion and viscosity in supercooled liquid water. *Phys Rev E Stat Phys Plasmas Fluids Relat Interdiscip Topics* 2000; 62(3 Pt A): 3572-80.  
[<http://dx.doi.org/10.1103/PhysRevE.62.3572>] [PMID: 11088858]



- [53] Varga B, Migliardo F, Takacs E, Vertessy B, Magazù S, Mondelli C. Neutron scattering studies on dUTPase complex in the presence of bioprotectant systems. *Chem Phys* 2008; 345(2-3): 250-8. [http://dx.doi.org/10.1016/j.chemphys.2007.07.004]
- [54] Hennet L, Cristiglio V, Kozaily J, *et al.* Aerodynamic levitation and laser heating: Applications at synchrotron and neutron sources. *Eur Phys J Spec Top* 2011; 196(1): 151-65. [http://dx.doi.org/10.1140/epjst/e2011-01425-0]
- [55] Branca C, Magazù S, Migliardo F, Migliardo P. Destructuring effect of trehalose on the tetrahedral network of water: A Raman and neutron diffraction comparison. *Physica A* 2002; 304(1-2): 314-8. [http://dx.doi.org/10.1016/S0378-4371(01)00517-9]
- [56] Magazù S, Migliardo F, Benedetto A. Mean square displacements from elastic incoherent neutron scattering evaluated by spectrometers working with different energy resolution on dry and hydrated (H<sub>2</sub>O and D<sub>2</sub>O) lysozyme. *J Phys Chem B* 2010; 114(28): 9268-74. [http://dx.doi.org/10.1021/jp102436y] [PMID: 20575549]
- [57] Magazù S, Calabrò E, Caccamo MT, Cannuli A. The shielding action of disaccharides for typical proteins in aqueous solution against static, 50 Hz and 1800 MHz frequencies electromagnetic fields. *Curr Chem Biol* 2016; 10(1) [http://dx.doi.org/10.2174/2212796810666160419153722]
- [58] Jovanović N, Bouchard A, Hofland GW, Witkamp GJ, Crommelin DJ, Jiskoot W. Distinct effects of sucrose and trehalose on protein stability during supercritical fluid drying and freeze-drying. *Eur J Pharm Sci* 2006; 27(4): 336-45. [http://dx.doi.org/10.1016/j.ejps.2005.11.003] [PMID: 16338123]
- [59] Calabrò E, Condello S, Currò M, *et al.* Effects of low intensity static magnetic field on FTIR spectra and ROS production in SH-SY5Y neuronal-like cells. *Bioelectromagnetics* 2013; 34(8): 618-29. [http://dx.doi.org/10.1002/bem.21815] [PMID: 24217848]
- [60] Barreca D, Laganà G, Ficarra S, *et al.* Spectroscopic determination of lysozyme conformational changes in the presence of trehalose and guanidine. *Cell Biochem Biophys* 2013; 66(2): 297-307. [http://dx.doi.org/10.1007/s12013-012-9485-4] [PMID: 23184705]
- [61] Barreca D, Laganà G, Ficarra S, *et al.* Anti-aggregation properties of trehalose on heat-induced secondary structure and conformation changes of bovine serum albumin. *Biophys Chem* 2010; 147(3): 146-52. [http://dx.doi.org/10.1016/j.bpc.2010.01.010] [PMID: 20171005]
- [62] Barreca D, Laganà G, Bruno G, Magazù S, Bellocco E. Diosmin binding to human serum albumin and its preventive action against degradation due to oxidative injuries. *Biochimie* 2013; 95(11): 2042-9. [http://dx.doi.org/10.1016/j.biochi.2013.07.014] [PMID: 23886889]
- [63] Magazù S, Calabrò E, Campo S, Interdonato S. New insights into bioprotective effectiveness of disaccharides: an FTIR study of human haemoglobin aqueous solutions exposed to static magnetic fields. *J Biol Phys* 2012; 38(1): 61-74. [http://dx.doi.org/10.1007/s10867-010-9209-1] [PMID: 23277670]
- [64] Branca C, Magazù S, Maisano G, Migliardo P, Villari V, Sokolov AP. The fragile character and structure-breaker role of  $\alpha$ ,  $\alpha$ -trehalose: Viscosity and Raman scattering findings. *J Phys Condens Matter* 1999; 11(19): 3823-32. [http://dx.doi.org/10.1088/0953-8984/11/19/305]
- [65] Branca C, Magazù S, Maisano G, Migliardo P, Tettamanti E. On the bioprotective effectiveness of trehalose: Ultrasonic technique, Raman scattering and NMR investigations. *J Mol Struct* 1999; 480-481: 133-40. [http://dx.doi.org/10.1016/S0022-2860(98)00626-7]
- [66] Branca C, Magazù S, Maisano G, Migliardo P.  $\alpha$ ,  $\alpha$ -trehalose-water solutions. 3. Vibrational dynamics studies by inelastic light scattering. *J Phys Chem B* 1999; 103(8): 1347-53. [http://dx.doi.org/10.1021/jp983470c]
- [67] Magazù S, Maisano G, Migliardo P, Middendorf HD, Villari V. Hydration and transport properties of aqueous solutions of  $\alpha$ - $\alpha$ -trehalose. *J Chem Phys* 1998; 109(3): 1170-4. [http://dx.doi.org/10.1063/1.476662]
- [68] Magazù S, Maisano G, Middendorf HD, Migliardo P, Musolino M, Villari V.  $\alpha$ ,  $\alpha$ -trehalose-water solutions. II. Influence of hydrogen bond connectivity on transport properties. *J Phys Chem B* 1998; 102(11): 2060-3. [http://dx.doi.org/10.1021/jp980235l]
- [69] Duong T, Barrangou R, Russell WM, Klaenhammer TR. Characterization of the tre locus and analysis of trehalose cryoprotection in *Lactobacillus acidophilus* NCFM. *Appl Environ Microbiol* 2006; 72(2): 1218-25. [http://dx.doi.org/10.1128/AEM.72.2.1218-1225.2006] [PMID: 16461669]
- [70] Magazù S, Maisano G, Migliardo F, *et al.* Characterization of molecular motions in biomolecular systems by elastic incoherent neutron scattering. *J Chem Phys* 2008; 129(15): 155103. [http://dx.doi.org/10.1063/1.2989804] [PMID: 19045233]
- [71] Blazhnov IV, Magazù S, Maisano G, Malomuzh NP, Migliardo F. Macro- and microdefinitions of fragility of hydrogen-bonded glass-forming liquids. *Phys Rev E Stat Nonlin Soft Matter Phys* 2006; 73(3 Pt 1): 031201. [http://dx.doi.org/10.1103/PhysRevE.73.031201] [PMID: 16605509]

- [72] Magazù S, Migliardo F, Malomuzh NP, Blazhnov IV. Theoretical and experimental models on viscosity: I. Glycerol. *J Phys Chem B* 2007; 111(32): 9563-70. [<http://dx.doi.org/10.1021/jp071949b>] [PMID: 17655214]
- [73] Leszek B. Dimensional changes of waterlogged archaeological hardwoods pre-treated with aqueous mixtures of lactitol/trehalose and mannitol/trehalose before freeze-drying. *J Cult Herit* 2015; 16(6): 876-82. [<http://dx.doi.org/10.1016/j.culher.2015.03.010>]
- [74] Magazù S, Maisano G, Migliardo F, Benedetto A. Elastic incoherent neutron scattering on systems of biophysical interest: mean square displacement evaluation from self-distribution function. *J Phys Chem B* 2008; 112(30): 8936-42. [<http://dx.doi.org/10.1021/jp711930b>] [PMID: 18610954]
- [75] Lombardo D, Kiselev MA, Magazù S, Calandra P. Amphiphiles self-assembly: Basic concepts and future perspectives of supramolecular approaches. *Adv Condens Matter Phys* 2015; 2015: 151683. [<http://dx.doi.org/10.1155/2015/151683>]
- [76] Nagase H, Endo T, Ueda H, Nakagaki M. An anhydrous polymorphic form of trehalose. *Carbohydr Res* 2002; 337(2): 167-73. [[http://dx.doi.org/10.1016/S0008-6215\(01\)00294-4](http://dx.doi.org/10.1016/S0008-6215(01)00294-4)] [PMID: 11814449]
- [77] Magazù S, Calabrò E, Campo S, Interdonato S. New insights into bioprotective effectiveness of disaccharides: an FTIR study of human haemoglobin aqueous solutions exposed to static magnetic fields. *J Biol Phys* 2012; 38(1): 61-74. [<http://dx.doi.org/10.1007/s10867-010-9209-1>] [PMID: 23277670]
- [78] Crowe JH, Crowe LM, Chapman D. Preservation of membranes in anhydrobiotic organisms: The role of trehalose. *Science* 1984; 223(4637): 701-3. [<http://dx.doi.org/10.1126/science.223.4637.701>] [PMID: 17841031]
- [79] Magazù S, Migliardo F, Affouard F, Descamps M, Telling MTF. Study of the relaxational and vibrational dynamics of bioprotectant glass-forming mixtures by neutron scattering and molecular dynamics simulation. *J Chem Phys* 2010; 132(18): 184512. [<http://dx.doi.org/10.1063/1.3407428>]
- [80] Magazù S, Maisano G, Migliardo F, Benedetto A. Mean square displacement evaluation by elastic neutron scattering self-distribution function. *Phys Rev E Stat Nonlin Soft Matter Phys* 2008; 77(6 Pt 1): 061802. [<http://dx.doi.org/10.1103/PhysRevE.77.061802>] [PMID: 18643290]
- [81] Magazù S, Maisano G, Migliardo F, *et al.* Characterization of molecular motions in biomolecular systems by elastic incoherent neutron scattering. *J Chem Phys* 2008; 129(15): 155103. [<http://dx.doi.org/10.1063/1.2989804>] [PMID: 19045233]
- [82] Magazù S. IQENS - Dynamic light scattering complementarity on hydrogenous systems. *Phys B* 1996; 226(1-3): 92-106. [[http://dx.doi.org/10.1016/0921-4526\(96\)00255-4](http://dx.doi.org/10.1016/0921-4526(96)00255-4)]
- [83] Lupi L, Comez L, Paolantoni M, Fioretto D, Ladanyi BM. Dynamics of biological water: insights from molecular modeling of light scattering in aqueous trehalose solutions. *J Phys Chem B* 2012; 116(25): 7499-508. [<http://dx.doi.org/10.1021/jp301988f>] [PMID: 22651571]
- [84] Branca C, Magazù S, Maisano G, Migliardo P, Villari V. Conformational distribution of poly(ethylene oxide) in molten phase and in aqueous solution by quasi-elastic and inelastic light scattering. *J Phys Condens Matter* 1998; 10(45): 10141-57. [<http://dx.doi.org/10.1088/0953-8984/10/45/004>]
- [85] Faraone A, Magazù S, Maisano G, Ponterio R, Villari V. Experimental evidence of slow dynamics in semidilute polymer solutions. *Macromol* 1999; 32(4): 1128-33. [<http://dx.doi.org/10.1021/ma9809684>]
- [86] Magazù S, Maisano G, Mallamace F, Micali N. Growth of fractal aggregates in water solutions of macromolecules by light scattering. *Phys Rev A Gen Phys* 1989; 39(8): 4195-200. [<http://dx.doi.org/10.1103/PhysRevA.39.4195>] [PMID: 9901749]
- [87] Jannelli MP, Magazù S, Migliardo P, Aliotta F, Tettamanti E. Transport properties of liquid alcohols investigated by IQENS, NMR and DLS studies. *J Phys Cond Matter. J Phys Condens Matter* 1996; 8(43): 8157-71. [<http://dx.doi.org/10.1088/0953-8984/8/43/012>]
- [88] Caccamo MT, Magazù S. Tagging the oligomer-to-polymer crossover on EG and PEGs by infrared and Raman spectroscopies and by wavelet cross-correlation spectral analysis. *Vib Spectrosc* 2016; 2016(85): 222-7. [<http://dx.doi.org/10.1016/j.vibspec.2016.04.017>]
- [89] Magazù S, Migliardo F, Caccamo MT. Innovative wavelet protocols in analyzing elastic incoherent neutron scattering. *J Phys Chem B* 2012; 116(31): 9417-23. [<http://dx.doi.org/10.1021/jp3060087>] [PMID: 22793379]
- [90] Cannuli A, Caccamo M T, Castorina G, Colombo F, Magazù S. Laser Techniques on Acoustically Levitated Droplets EPJ Web of Conferences 167, article number 05010, 2018. [<http://dx.doi.org/10.1051/epjconf/201816705010>]
- [91] Caccamo MT, Magazù S. Ethylene Glycol - Polyethylene Glycol (EG-PEG) mixtures: Infrared spectra wavelet cross-correlation analysis. *Appl Spectrosc* 2017; 71(3): 401-9.

- [http://dx.doi.org/10.1177/0003702816662882] [PMID: 27558367]
- [92] Caccamo MT, Magazù S. Multiscaling wavelet analysis of infrared and raman data on polyethylene glycol 1000 aqueous solutions. *Spectrosc Lett* 2017; 50(3): 130-6.  
[http://dx.doi.org/10.1080/00387010.2017.1291524]
- [93] Migliardo F, Magazù S, Caccamo MT. Infrared, raman and INS studies of poly-ethylene oxide oligomers. *J Mol Struct* 2013; 1048: 261-6.  
[http://dx.doi.org/10.1016/j.molstruc.2013.05.060]
- [94] Magazù S, Calabrò E, Campo S. FTIR spectroscopy studies on the bioprotective effectiveness of trehalose on human hemoglobin aqueous solutions under 50 Hz electromagnetic field exposure. *J Phys Chem B* 2010; 114(37): 12144-9.
- [95] Branca C, Magazù S, Maisano G. S. M. Bennington A, Fåk B. Vibrational studies on disaccharide/H<sub>2</sub>O systems by inelastic neutron scattering, raman, and IR spectroscopy. *J Phys Chem B* 2003; 107(6): 1444-51.  
[http://dx.doi.org/10.1021/jp026255b]
- [96] Caccamo MT, Zammuto V, Gugliandolo C, Madeleine-Perdrillat C, Spanò A, Magazù S. Thermal restraint of a bacterial exopolysaccharide of shallow vent origin. *Int J Biol Macromol* 2018; 114: 649-55.  
[http://dx.doi.org/10.1016/j.ijbiomac.2018.03.160] [PMID: 29601879]
- [97] Caccamo MT, Magazù S. Thermal restraint on PEG-EG mixtures by FTIR investigations and wavelet cross-correlation analysis. *Polym Test* 2017; 62: 311-8.  
[http://dx.doi.org/10.1016/j.polymertesting.2017.07.008]

---

© 2018 Magazù *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: (<https://creativecommons.org/licenses/by/4.0/legalcode>). This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.