

The Open Cardiovascular Medicine Journal

Content list available at: <https://opencardiovascularmedicinejournal.com>



REVIEW ARTICLE

Role of Warburg Effect in Cardiovascular Diseases: A Potential Treatment Option

Niken Puspa Kuspriyanti^{1,2}, Eko Fuji Ariyanto³ and Mas Rizky A. A. Syamsunarno^{3,*}

¹Biomedical Science Master Program, Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia

²Department of Biomedical Sciences, Faculty of Medicine, Universitas Pasundan, Bandung, Indonesia

³Department of Biomedical Sciences, Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia

Abstract:

Background:

Under normal conditions, the heart obtains ATP through the oxidation of fatty acids, glucose, and ketones. While fatty acids are the main source of energy in the heart, under certain conditions, the main source of energy shifts to glucose where pyruvate converts into lactate, to meet the energy demand. The Warburg effect is the energy shift from oxidative phosphorylation to glycolysis in the presence of oxygen. This effect is observed in tumors as well as in diseases, including cardiovascular diseases. If glycolysis is more dominant than glucose oxidation, the two pathways uncouple, contributing to the severity of the heart condition. Recently, several studies have documented changes in metabolism in several cardiovascular diseases; however, the specific mechanisms remain unclear.

Methods:

This literature review was conducted by an electronic database of Pub Med, Google Scholar, and Scopus published until 2020. Relevant papers are selected based on inclusion and exclusion criteria.

Results:

A total of 162 potentially relevant articles after the title and abstract screening were screened for full-text. Finally, 135 papers were included for the review article.

Discussion:

This review discusses the effects of alterations in glucose metabolism, particularly the Warburg effect, on cardiovascular diseases, including heart failure, atrial fibrillation, and cardiac hypertrophy.

Conclusion:

Reversing the Warburg effect could become a potential treatment option for cardiovascular diseases.

Keywords: Warburg effect, Metabolism, Cardiovascular disease, Atrial fibrillation, Heart failure, Cardiac hypertrophy.

Article History

Received: September 14, 2020

Revised: January 05, 2021

Accepted: January 06, 2021

1. INTRODUCTION

The heart pumps blood throughout the body *via* the circulatory system [1]. This process requires energy intake from carbohydrates, lipids, and proteins. Cardiomyocytes are heart muscle cells that can convert glucose or ketone bodies into mechanical energy [2]. While cardiomyocytes predominantly use fatty acids (FAs) as their energy source, glucose provides approximately 25% of the total energy required by the heart [3, 4]. This is different from other cells where almost 70% of glucose is used as the energy source *via*

oxidative phosphorylation; the rest is obtained *via* glycolysis. In the human heart, approximately 60-70% of the ATP produced in the mitochondria is used as an energy source, whereas the remainder is used for calcium homeostasis, ion pumping into the sarcoplasmic reticulum *via* calcium ATPase, and cell death signaling [5]. The requirement for ATP varies depending on the cell type, cellular condition, growth status, and microenvironment [6].

In conditions of impaired myocardial energy, such as cardiac hypertrophy, a major metabolic shift occurs from FA oxidation to glucose utilization, affecting glucose uptake and glycolysis rate [7]. This is accompanied by a decrease in

* Address correspondence to this author at the Department of Biomedical Science, Faculty of Medicine, Universitas Padjadjaran, Jalan Raya, Bandung, Indonesia; Tel: 6281380036560; E-mail: rizky@unpad.ac.id

cardiac mitochondrial oxidative metabolism and an increase in glycolysis (pyruvate conversion into lactate) [8]. The energy produced *via* mitochondrial glucose oxidation is greater than that produced *via* glycolysis [9, 10]. However, cardiac hypertrophy and Heart Failure (HF) rat models decrease myocardial glucose oxidation. Several studies have shown that changes in energy metabolism in the heart increase cardiovascular disease severity, including cardiac hypertrophy and HF [11 - 13].

In HF, glucose metabolism increases glucose uptake and glycolysis, ultimately leading to uncoupling of glycolysis and glucose oxidation. This may also contribute to the development of HF by increasing cardiac hypertrophy [14]. In proliferating cells, such as cancer cells, uncoupling of glycolysis and glucose oxidation occurs, which plays an important role in promoting cell growth [6, 15]. In 1920, Otto Warburg reported an increase in glycolysis and lactate production in cancer cells compared with normal cells, despite the presence of oxygen. This is known as the “Warburg effect” [16]. Although glycolysis only produces two ATP molecules compared with oxidative phosphorylation, which produces 36 ATP molecules, glycolysis is a relatively quick process [17], and the total amount of ATP produced by glycolysis is higher than that by oxidative phosphorylation in the same amount of time [18, 19]. This “Warburg” phenomenon may also occur in other conditions, such as cardiac hypertrophy, which can, in turn, result in HF because increased pyruvate dehydrogenase kinase (PDK) activity is observed, suggesting an increased rate of glycolysis [20, 21].

Recent studies have proposed the existence of the Warburg effect in cardiovascular diseases. The metabolites produced by glycolysis play an important role in regulating extracellular matrix formation, cell proliferation, apoptosis, and autophagy [22]. In addition, glycolytic enzymes are found in many diseases, including pulmonary fibrosis, hypertension, HF, atrial fibrillation (AF), and cardiac hypertrophy [23 - 25]. The next section focuses on the metabolic shift involved in the signaling pathways and the Warburg effect in cardiovascular disease, and discusses their potential roles in cardiovascular disease therapy.

2. MATERIALS AND METHODS

Our aim was to review the Warburg effect in cardiovascular diseases and its molecular interactions as targeted therapy. We conducted a literature search using PubMed and Google Scholar for articles published up until 2020. The keywords used were: “cardiovascular disease” or “cardiac hypertrophy” or “heart failure” or “failing heart” and “Warburg effect,” “heart metabolism” or “heart failure metabolism” or “cardiac hypertrophy metabolism” or “atrial fibrillation metabolism,” and “heart failure treatment” or “cardiac hypertrophy treatment” or “atrial fibrillation treatment.” The inclusion criteria were articles published up until 2020; those focusing on the Warburg effect in cardiovascular disease; and those regarding metabolism and treatment in cardiac hypertrophy, heart failure, and AF that were related to the mechanisms of the Warburg effect.

3. RESULTS AND DISCUSSION

The initial search found 388 article sources. After duplicates were removed, 351 records remained. A further 184 records relevant to the inclusion criteria were articles published up until 2020; those focusing on the Warburg effect in cardiovascular disease and those regarding metabolism and treatment in cardiac hypertrophy, heart failure, and AF that were related to the mechanisms of the Warburg effect. A total of 162 potentially relevant articles after the title and abstract screening were screened for full-text. Finally, 135 papers were included for the review article.

3.1. Energy Metabolism in Cardiomyocytes

The adult heart pumping 10 tons of blood a day requires approximately 6 kg of ATP. The heart obtains energy from various metabolites, including glucose, FAs, lactate, and ketones [26]. Under normal heart conditions, >95% of the ATP is produced *via* oxidative phosphorylation; the rest is obtained *via* glycolysis. Approximately 50-70% of the ATP is produced *via* the oxidation of free FAs [27, 28].

The transport of FAs from the plasma to cardiomyocytes is facilitated by fatty acid translocase (FAT/CD36), fatty acid transport protein, plasma membrane fatty acid-binding protein (FABPpm), and heart-type fatty acid-binding protein [29]. FAs are esterified to fatty acyl-CoA, which enters the mitochondrial membrane with the help of carnitine palmitoyltransferase I and II (CPT I and II). Fatty acyl-CoA is converted to acetyl-CoA, which then enters the tricarboxylic acid (TCA) cycle to produce nicotinamide adenine dinucleotide (NADH) and flavin adenine dinucleotide (FADH₂). These cofactors are used in the electron transport chain to produce ATP [30]. Malonyl-CoA negatively regulates CPT I activity and is formed *via* acetyl-CoA carboxylase (ACC) and degraded by malonyl-CoA decarboxylase. AMP-activated protein kinase (AMPK) phosphorylates ACC, which inhibits malonyl-CoA formation and increases FA oxidation. Under normal heart conditions, FA oxidation, esterification, and uptake are activated by peroxisome proliferator-activated receptor coactivator-1 α (PGC1 α), which stimulates peroxisome proliferator-activated receptor α (PPAR α). PPAR α can modify glucose oxidation by inhibiting pyruvate dehydrogenase (PDH) *via* pyruvate dehydrogenase kinase 4 (PDK4) [31]. PDK is activated by NADH and acetyl-CoA, which are produced *via* FA oxidation, and directly inhibits PDH. PDK is inhibited by pyruvate and reduced the acetyl-CoA/free CoA and NADH/NAD⁺ ratios [32].

The main sources of glucose used by the heart are *via* the uptake of exogenously obtained glucose and glycogen stored in the liver. Glycolysis plays a critical role in cell metabolism and produces pyruvate, NADH, and ATP. Pyruvate is converted to lactate by lactate dehydrogenase (LDH) and mostly undergoes mitochondrial glucose oxidation [33]. Glucose from the circulation is transported to cells *via* glucose transporters, including GLUT1 and GLUT4. In the fetal heart, GLUT1 is the major glucose transporter, and its level increases in response to myocardial stress or injury. In contrast, GLUT4 is the predominant isoform in the adult heart and plays a role in basal myocardial glucose uptake [34]. The deletion of GLUT4 in

mice hearts leads to metabolic adaptations [35]. GLUT 4-knockout mice (G4H^{-/-}) show changes in glycolysis as a defense mechanism against ischemic injury [36]. After being absorbed in the intestine, glucose is phosphorylated to form glucose-6-phosphate (G6P), which then enters the metabolic pathway. In the mitochondria, pyruvate is decarboxylated by pyruvate dehydrogenase to form acetyl-CoA, which then enters the TCA cycle to produce NADH and FADH₂. Pyruvate oxidation is regulated by PDH. ATP plays an important role in calcium absorption in the sarcoplasm and relaxation during diastole [37]. Glucose is catalyzed by glycolysis and intermediate glycolysis. Glucose is converted to G6P by hexokinase (HK). G6P then enters the pentose phosphate pathway (PPP) [38]. PPP produces NADPH, which maintains levels of reduced glutathione and ribulose 5-phosphate used for nucleotide biosynthesis or for the generation of intermediates for glycolysis [39, 40]. The polyol pathway is another glycolytic pathway in which glucose is converted to sorbitol by aldose reductase (AR). Sorbitol dehydrogenase then oxidizes sorbitol to fructose and advanced glycation end-products [41]. A study has reported that AR activity increased in the heart and kidneys of a hyperglycemic mouse model [42]. The hexosamine biosynthetic pathway generates the end products hyaluronan, glycosylphosphatidylinositol anchor, glycolipid, proteoglycan, N-glycan, and O-linked β-N-acetylglucosamine (O-GlcNAc), which are derived from uridine diphosphate N-acetylglucosamine [33]. In cardiac hypertrophy, inhibition of O-GlcNAc can reduce the progression of the development of hypertrophy [43, 44].

Ketone bodies, such as acetoacetate and β-hydroxybutyrate, are energy sources produced *via* FA β-oxidation in the liver [45]. Ketones are oxidized in the heart, brain, and skeletal muscle [30]. During terminal oxidation, ketones are converted back to acetyl-CoA in a reaction catalyzed by succinyl-CoA-3-oxoacid-CoA transferase, β-hydroxybutyrate dehydrogenase 1, and acetyl-CoA acetyltransferase 1-or through the lipogenic pathway *via* fatty acid synthase [46]. The heart uses ketones to maintain myocardial function in HF. In HF, the decrease in FA oxidation makes ketone bodies an efficient fuel for ATP production [47] (Fig. 1).

3.2. Warburg Effect in Cardiovascular Diseases

Under normal heart conditions, FAs are the main source of ATP production. In contrast, under various pathological conditions, including cardiac hypertrophy, there is a shift in energy metabolism from mitochondrial oxidative phosphorylation and the TCA cycle to glycolysis [48]. The mass changes in cardiac hypertrophy are associated with metabolic changes, resulting in switching to more ATP production from glucose than from FAs. Glycolysis is a biochemical process that converts glucose to pyruvate and then to lactate and CO₂ to produce ATP. Carbon atoms obtained from glucose are either derived exogenously or endogenously

by glycogen breakdown [49]. The change in the rates of glycolysis and oxidative phosphorylation results in the uncoupling of glycolysis and glucose oxidation, which interferes with cardiac function [8]. In HF, there is an increase in glycolysis with or without a decrease in mitochondrial oxidation [20, 50]. Impaired myocardial energy metabolism in cardiac hypertrophy is characterized by increased glucose demand and decreased FA oxidation. These alterations are associated with increased glycolysis in cardiac hypertrophy [51 - 54]. Previous studies have shown that an increase in glucose uptake and the activation of phosphofructokinase (PFK), play a role in increasing the rate of glycolysis in rats with hypertrophied hearts. In left ventricular hypertrophy (LVH), there is an increase in adenosine monophosphate (AMP) and adenosine diphosphate (ADP), triggering AMPK, which in turn increases glucose transport and PFK activation [54].

Dahl salt-sensitive rats with HF and preserved ejection fraction showed decreased FA metabolism and increased glucose metabolism. Uncoupling between glycolysis and glucose oxidation, resulting from increased glycolysis, occurs without changes in glucose oxidation; this results in an acidic environment that hampers cardiac function [8]. The Warburg effect is known to occur in cancer cells. These cells have high proliferation rates and exhibit high glycolysis rates in the presence of oxygen [55, 56]. In addition to cancer cells, the Warburg effect may occur in other cells with high energy demand. This effect has been demonstrated in cardiac hypertrophy, which causes HF. Inhibition of anaerobic glycolysis can reduce hypertrophy and increase the flexibility of cardiac metabolism to maintain the need for cardiac output to meet the high metabolic demand [57]. HF and hypertrophy mouse models showed decreased myocardial glucose oxidation levels. Metabolic changes are believed to increase the severity of HF [30].

Waleed et al. developed a post-infarction mouse model and revealed an imbalance in glucose metabolism and increased proton production, resulting in the deterioration and disruption in mechanical left ventricular function [14]. Systolic dysfunction and glucose oxidation damage in HF may be caused by mitochondrial dysfunction, decrease in the expression of the genes involved in glucose oxidation, or an increase in the expression of the PDH complex [11, 58]. Increased oxidative stress may be a cause of mitochondrial dysfunction in cardiac hypertrophy, HF, and AF. Reactive oxygen species (ROS) play a role in cardiac remodeling in the stress response *via* increased mitochondrial biogenesis, cardiomyocyte volume, and capillary density [59]. During chronic stress, ATP demand increases in the heart, leading to increased activity of the electron transport chain in the mitochondria and ROS generation. This results in a shift in the energy metabolism to glycolysis, even in the conditions of adequate oxygen levels, as compensation for the high energy demand. This could be the underlying mechanism of the Warburg effect in cardiovascular diseases, particularly HF.




Fig. (1). Glucose metabolism in the heart. During glucose metabolism (yellow arrows), glucose enters the cells *via* the glucose transporters GLUT1 and GLUT4. Glycolysis begins with the conversion of glucose to glucose-6-phosphate, which is, in turn, is converted to pyruvate by pyruvate dehydrogenase. Pyruvate either enters the tricarboxylic acid (TCA) cycle in the mitochondria to produce ATP *via* the electron transport chain or is converted to lactate by lactate dehydrogenase (LDH). In addition to glycolysis, glucose undergoes intermediate glycolysis *via* the hexosamine biosynthetic pathway (HBP), pentose phosphate pathway (PPP), and polyol pathway. Circulating fatty acids (FAs) cross the cardiomyocyte membrane (red arrows) *via* the transporters fatty acid translocase (FAT/CD36), plasma membrane fatty acid-binding protein (FABP), and fatty acid transport protein (FATP). Fatty acyl-CoA enters the mitochondria *via* carnitine palmitoyltransferase (CPT) I and CPT II and undergoes β -oxidation. Ketones are also converted to acetyl-CoA (blue arrows) by succinyl-CoA: 3-oxoacid-CoA transferase (SCOT) and β -hydroxybutyrate dehydrogenase 1 (BHD1).

Other studies have shown that FA oxidation stimulation and oxidative phosphorylation improve HF [11, 14, 60]. We previously reported that mice with CD36 genetic deletions showed decreased FA uptake and increased glycolysis flux; this, in turn, increased the progression from compensated hypertrophy to HF. This phenomenon was also observed in mice models of heart pressure overload with a double knockout of FABP4 and FABP5, with increased glycolysis and decreased FA uptake, resulting in severe hypertrophy, fibrosis, and cardiac dysfunction. Furthermore, the use of glucose was demonstrated not only for catabolic pathways but also for anabolic pathways to synthesize biomass [61, 62]. CD36 increased the expression of Forkhead Box O1 and PDK4 *via* the activation of PPAR δ / β to inhibit glucose oxidation [63]. This could support the association between the Warburg effect and cardiovascular diseases due to increased glycolysis, even accompanied by glucose oxidation. The increase in glycolysis in this study [61 - 63] may have involved the Warburg effect, although it remains unclear. Further research and clinical studies in humans are needed.

Structural and electrical remodeling also occurs in AF [64]. In acute AF, contraction and electrical activity increase four- to six-fold compared with normal atrial conditions. Tu et al. reported the downregulation of FA metabolism in the atria of permanent AF due to the downregulation of the enzymes regulating FA oxidation [65]. Tsuboi et al. reported that the deletion of mitochondrial DNA in patients with AF led to the

disruption of ATP synthesis [66]. Decreased oxidation of pyruvate and FAs increases glycolysis in AF, leading to the upregulation of glycolytic enzymes, lactate production, and the downregulation of the PDH complex [25, 37]. Myocardial fibrosis and metabolic stress are related to the Warburg effect and play a role in the development of AF. An increase in PDK4 is associated with an increase in myocardial fibrosis.

The Warburg theory suggests an ATP production shift from the TCA cycle and oxidative phosphorylation to glycolysis or lactate production, even in the presence of oxygen, and is thus termed aerobic glycolysis [55, 67, 68]. Under normal conditions, most of the glucose taken up by cells is metabolized *via* glycolysis to pyruvate, which then enters the TCA cycle and electron transport chain to generate ATP *via* oxidative phosphorylation [69]. Studies have shown that the increase in oxidative phosphorylation in cancer cells can be achieved by inhibiting glycolysis. Activation of glycolysis in the proliferative cells supports the cells to grow [15, 70]. Glycolysis can be induced by hypoxia *via* activation of hypoxia-inducible transcription factor 1 α (HIF-1 α). HIF-1 is a heterodimeric transcription factor that changes hypoxic gene expression and consists of HIF-1 α and HIF-1 β subunits. It is thought to be adaptive for cells exposed to an oxygen-deficient environment. PDK1 is the target of the HIF-1 α gene, which increases PDK activation and leads to the inhibition of PDH, which allows pyruvate to enter the mitochondria to perform oxidative phosphorylation. PDK1 regulates mitochondrial

function in hypoxic states by inhibiting the entry of pyruvate into the TCA cycle, indirectly inhibiting PDH [71 - 74].

In a monocrotaline-induced model of pulmonary hypertension, the glycolysis rate of right ventricular hypertrophy (RVH) was increased compared with that of the control group, and there was no increase in glucose oxidation. An increase in GLUT1 was observed as a compensatory mechanism of low ATP levels. Decreased glucose oxidation is thought to occur due to PDH phosphorylation [20]. Decreased glucose oxidation rate in fawn-hooded rats with pulmonary hypertension and RVH occurs due to increased PDK4 levels. Therefore, inhibition of PDK can increase PDH activity to prevent HF [21]. Kato et al. reported metabolic remodeling in compensated left ventricle hypertrophy and HF and showed an increase in glycolysis, characterized by an increase in lactate and a decrease in PDH [11]. Studies have reported that an increase in LDH, which converts pyruvate to lactate, results in increased lactic efflux in the hypertrophied heart muscle [75]. PH alteration and increase in protons can impair the sensitivity of troponin I to calcium, which interferes with cardiac contractility [76]. The upregulation of glycolysis rates results in uncoupling between glycolysis and glucose oxidation, leading to an increase in proton and lactate production [77, 78]. This suggests that an increase in lactate levels is due to a higher glycolysis rate compared with mitochondrial oxidation and that increased PDK inhibits the activation of the PDH complex and the biosynthesis of the precursors, which are needed to be required for the development of hypertrophy [23, 79]. Decreased oxidation of pyruvate and FAs increases glycolysis in AF, leading to the upregulation of glycolytic enzymes, lactate production, and the downregulation of the PDH complex. The Warburg effect increased glucose use and fibrosis levels in a canine model of p-AF cardiomyopathy, which is characterized by the increased expression of PDK1, PDK4, LDHA, and lactic acid and decreased expression of isocitrate dehydrogenase and PDH [80]. Increased glycolytic

enzymes in cardiovascular diseases, as seen in cancer cells, indicate the involvement of the Warburg effect.

HIF-1 α plays an important role in energy changes from oxidative phosphorylation to glycolysis. HIF-1 α is the main hallmark and regulator of the Warburg effect in cancer. It is upregulated under cardiac hypertrophy conditions [79, 81, 82]. Increased HIF-1 α expression in AF is an early response of atrial cardiomyocytes. In AF models, there was a decrease in PPAR α /PGC1 α activity that was inhibited by HIF-1 α , which caused a decrease in FA metabolism [81, 83]. HIF-1 α plays a role in metabolic reprogramming in end-stage HF. Furthermore, it regulates the gene expression of PDK1 and LDHA, which play a role in reprogramming glucose metabolism [84]. Interactions between HIF-1 α and pyruvate kinase M2 (PKM2) in the nucleus activate the transcription of genes involved in glycolysis, such as GLUT 1, LDHA, and PDK1 [85]. PKM2 is considered the hallmark of the Warburg effect in cancer cells [86]. HIF-1 α activates the conversion of adult pyruvate kinase M1 (PKM1) to fetal PKM2. PKM1 is normally expressed continuously in normal cells, whereas PKM2 is widely expressed in cancer cells [87]. Metabolic changes that occur in heart diseases cause a metabolic switch back to fetal metabolism, in which the glycolysis rate exceeds that of oxidative phosphorylation [88, 89]. The fetal heart uses glucose as the main energy source. A few days after birth, the heart switches to using FA oxidation as its main energy source. In the heart of a newborn rabbit, approximately 44% of the ATP was derived *via* glycolysis; however, after 7 days, only approximately 7% of the ATP was produced *via* glycolysis [87, 90, 91]. In cancer cells, HIF-1 suppresses the TCA cycle *via* PDK activation and functions to maintain ATP and ROS production [72]. The increase of PDK regulation leads to reduced levels of active PDH *via* PDH phosphorylation [73]. Suppression of HIF-1 α prevents hypertrophy-induced cardiac dysfunction [92]. The Warburg effect and HIF-1 α are thought to result in the metabolic shift to glycolysis in both cancer cells and cardiovascular diseases (Table 1).

Table 1. Role of the Warburg effect in cardiovascular diseases.

No.	Disease	Model	Event	Reference
1	LVH	Weanling male Wistar rats	Glycolysis \uparrow Lactate \uparrow	[54]
2	LVH HF	Inbred male Dahl Salt-sensitive rats	Glycolysis $\uparrow\uparrow$ Lactate \uparrow	[11]
3	LVH	Male Wistar rats	PDH \downarrow PDK \uparrow	[93]
4	HF	Male C57BL/6 mice with coronary artery ligation or sham operation procedure	Glycolysis \uparrow Glucose oxidation \uparrow Lactate \uparrow	[14]
5	HF	Male Dahl salt-sensitive rats with induced HfpEF	Glycolysis \uparrow Glucose oxidation \leftrightarrow FA oxidation \downarrow Lactate \uparrow Proton \uparrow	[8]
6	LVH	Mb transgenic mice with transverse aortic constriction surgery	Glycolysis \uparrow F-2,6-P2 \uparrow	[94]
7	RVH	Adult male Sprague–Dawley rats injected monocrotaline	Glycolysis \uparrow Glucose oxidation \leftrightarrow PDH \downarrow PDK \uparrow	[20]

No.	Disease	Model	Event	Reference
8	RVH	Fawn-hooded rats	Glycolysis ↑ Glucose oxidation ↓ PDK ↑	-
9	Chronic cardiac hypertrophy	Mouse transaortic-constriction	Glycolysis ↑ p-PDH ↑	[95]
10	AF	Beagle Canine model of paroxysmal atrial fibrillation	Glycolysis ↑ LDHA ↑ PDK1 ↑ PDK4 ↑	[80]

AF, atrial fibrillation; FA, fatty acid; HF, heart failure; F-2,6-P2, fructose 2,6-bisphosphate; HFpEF, heart failure with preserved ejection fraction; LVH, left ventricular hypertrophy; PDH, pyruvate dehydrogenase; PDK, pyruvate dehydrogenase kinase; RVH, right ventricular hypertrophy

3.3. Warburg Effect: Targeted Treatment in Cardiovascular Diseases?

In cardiac hypertrophy, cell growth and proliferation are accompanied by decreased FA β -oxidation, which is characterized by changes in the expression of enzymes involved in glycolysis and FA β -oxidation [33]. In HF, ATP is decreased to 30-40% compared with that in the normal heart due to a switch to the use of glucose as the primary energy source instead of FAs. In HF, a decrease in myocardial ATP up to 60-70% is accompanied by a decrease in mitochondrial glucose oxidation, increased glycolysis, and decreased creatine kinase activity [11, 30, 96]. Therefore, promoting the coupling of glucose oxidation and glycolysis is one approach to treat HF because it enhances oxidation [97]. Correcting cardiac function *via* the reverse Warburg effect can be achieved by stimulating glucose oxidation [8].

Dichloroacetate (DCA) can induce oxidative phosphorylation and the activation of the PDH complex and has been widely studied. Inhibition of aerobic glycolysis increases cardiac metabolic flexibility and reduces hypertrophy *via* PDH inhibition. In ischemia-reperfusion models, DCA has a cardioprotective mechanism [98, 99]. In an MCT-RVF, RVH leads to an increase in PDK activity. The administration of DCA, a pyruvate dehydrogenase kinase inhibitor, decreased the expression of PDK in MCT-RVF mice [20]. DCA decreased LDHA and upregulated superoxide dismutase 2 [100]. In Dahl-sensitive rats, the use of DCA treatment prevented the progression of LVH to HF. DCA was shown to improve cardiac function and survival in the rat model [11]. Interestingly, DCA treatment in mice with hyperthyroidism showed an increase in PDH activity and increased myocardial flexibility and reduced hypertrophy [93]. Clinical data on the use of DCA remains sparse. A clinical study of 10 patients with HF (New York Heart Association class III and IV) infused with DCA intravenously showed improved LV stroke volume and myocardial oxygen consumption [101]. The effect of DCA might inhibit the Warburg effect by improving the coupling of glycolysis to glucose oxidation by inhibiting PDK to activate PDH, thereby increasing pyruvate oxidation and mechanical efficiency.

Increased AMPK activity is associated with cardiac hypertrophy and HF. Targeting AMPK activity as a treatment in HF models improved cardiac function and reduced ventricular remodeling [102, 103]. AMPK phosphorylation is increased in patients with paroxysmal AF as an adaptive response to metabolic stress in AF for Ca^{2+} handling and

contractility [57]. Furthermore, AMPK increased levels of FA transport proteins FAT/CD36 and FABPpm [104]. In a mouse AF model, AMPK was associated with an increase in FAT/CD36, leading to excess uptake of FAs and decreased expression of GLUT4 in the membrane. Therefore, AMPK has cardioprotective functions and prevents the AF progression by increasing FA oxidation compared with glucose oxidation [51, 105]. Some drugs, such as metformin, statins, and resveratrol, activate AMPK. Metformin increases AMPK activity by directly increasing its phosphorylation or binding to AMPK subunits, thereby increasing the heterotrimeric complex. Metformin indirectly changes the ratio of AMP or ADP to ATP [106, 107]. In animal models, metformin phosphorylates AMPK and increases nitric oxide levels, thereby improving cardiac function [108]. The use of metformin as an AMPK activator in a Beagle dog AF model showed an increase in AMPK, PGC1 α , and PPAR1 α activity, and a decrease in HIF1 α , GLUT1, PDK, HK, and LDH. Metformin treatment also revealed that an increase in PDH is the key to the Warburg effect [109]. A clinical study has shown that the use of metformin reduces the mortality rate in ambulatory HF patients with diabetes [110]. AMPK activates FA metabolism *via* PGC1 α /PPAR1 α and inhibits HIF-1 α to reverse the Warburg effect [25]. Resveratrol is a natural polyphenol found in plants, including peanuts, cranberries, and grapes. It has a cardioprotective effect *via* the activation of AMPK/sirtuin 1, which is involved in signaling and modulating cardiac metabolism. The beneficial effects of resveratrol in HF have been shown in ischemic and non-ischemic HF animal models [111 - 113].

The activation of PPAR α and PGC1 α increases FA flux in cells as well as increases the activity of enzymes that play a role in FA β -oxidation [114]. Mice lacking PPAR α were shown to have increased glycolysis and decreased FA oxidation, characterized by a decrease in protein transport and in enzymes playing a role in β -oxidation (such as malonyl-CoA decarboxylase, which inhibits CPT I) [115]. In mouse HF models, treatment with fenofibrate, a PPAR α agonist, led to increased FA metabolism [116, 117] as well as the upregulation of PDH, GLUT4, PPAR α , PGC1 α , and p-sirtuin 1 and downregulation of PDK4 [118]. In dogs with high-frequency left ventricular pacing, fenofibrate administration prevented a decrease in FA metabolism and improved cardiac function [119]. The reverse Warburg effect can be generated by promoting the coupling of glycolysis and glucose oxidation and can be achieved by enhancing glucose and FA oxidation. Metabolic therapy by disrupting the Warburg effect represents

a potential treatment for cardiovascular diseases, although more extensive human studies are warranted to demonstrate this effect as well as the clinical implications of the Warburg effect on cardiovascular diseases (Fig. 2).

3.4. Dietary and Exercise Modifications in Cardiovascular Diseases

Behavioral modifications, such as a healthy lifestyle, including healthy dietary patterns and appropriate exercise, are important strategies recommended for the prevention of cardiovascular disease [120]. In recent years, epigenetic studies have shown that environmental factors, including diet and physical exercise, may change the epigenome.

Meta-analyses and randomized controlled trials have shown that a Mediterranean diet consisting of fruits, grains, vegetables, nuts, fish, unsaturated fats, and legumes reduces mortality and morbidity in patients with cardiovascular diseases [121, 122]. This healthy diet pattern should be low in trans fats, refined grains, red and processed meat, sugary drinks, and added sugars [123]. Cohort studies have shown an association between metabolite profiles and high adherence to dietary recommendations proposed by the Alternative Healthy Eating Index (AHEI) in patients with cardiovascular diseases [124]. High levels of monounsaturated and saturated FAs are associated with a high risk of cardiovascular diseases. Furthermore, a high ratio of omega-3 polyunsaturated fatty acids (PUFAs), such as docosahexaenoic acid, to total FAs is associated with a reduced risk of cardiovascular diseases [124]. A hereditary cardiomyopathy model demonstrated that

omega-3 PUFAs can reactivate the expression of genes damaged in cardiomyocytes and also prevent plasma membrane degradation, which maintains signal transmission from the membrane surface to the nucleus [125]. Nutrition can modulate myocardial genes by activating the transcription factor PPAR α [126, 127]. A population-based prospective cohort study has found that dietary intake of different PUFAs alters the association between genetic variation in fatty acid desaturase and high-density lipoprotein and low-density lipoprotein (LDL) [128]. Reduced risk of cardiovascular diseases is associated with the intake of polyphenol-rich diets related to DNA methylation. Polyphenols are present in fruits and vegetables, including cocoa and green tea. In vitro studies have shown that cocoa extract can inhibit the expression of genes encoding DNA methyltransferases and methylenetetrahydrofolate reductase, which are found in several disease conditions, including heart diseases [129, 130]. Cocoa polyphenol is cardioprotective and is associated with decreased LDL levels. Decreased LDL levels can lead to a reduced risk of atherosclerosis leading to cardiovascular disease [131, 132]. Dietary folic acid and vitamin B may reduce the risk of cardiovascular disease *via* the methylation of homocysteine to methionine, and folic acid deficiency increases cardiovascular diseases *via* DNA methylation [133].

Mouse model studies have shown increased expression of regulators of lipid metabolism, such as CPT I, PPAR α , and sterol regulatory element-binding protein 1c, and the up-regulation of CD36 gene expression [134, 135]. Dietary modifications and appropriate exercise can contribute to genetic changes in cardiovascular metabolism *via* epigenetics.




Fig. (2). Treatment targeting the Warburg effect in cardiovascular diseases. Agents that activate AMPK, such as metformin and resveratrol, inhibit hypoxia-inducible transcription factor 1 α (HIF1 α), CD36/FAT, and acetyl-CoA carboxylase (ACC). Other drugs, such as fenofibrate and dichloroacetate, inhibit the phosphorylation of pyruvate dehydrogenase (p-PDH) *via* the inhibition of pyruvate dehydrogenase (PDK).

CONCLUSION

The Warburg effect is an energy shift in glucose metabolism from oxidative phosphorylation to aerobic glycolysis that occurs in cancer as well as other non-cancer conditions, including cardiovascular diseases. The Warburg effect may also occur in highly proliferative cells involved in cardiovascular diseases such as cardiac hypertrophy, HF, and AF. The increased activity of glycolytic enzymes in cardiovascular diseases, as observed in cancer cells, indicates the involvement of the Warburg effect. The mechanism of the Warburg effect in the heart is not completely understood, although HIF-1 α is thought to play a role in its regulation. The Warburg effect can be used as a therapeutic target for improving heart conditions by inhibiting the shift of glucose oxidation to aerobic glycolysis. This can be achieved by increasing glucose oxidation *via* glycolysis inhibitors and increasing FA oxidation. Diet and exercise can also be used to prevent cardiovascular diseases. Further clinical studies are warranted to determine the involvement of the Warburg effect in cardiovascular diseases, identify more therapeutic targets, and assess the effect of epigenetics on the Warburg effect in cardiovascular diseases.

CONSENT FOR PUBLICATION

Not applicable.

FUNDING

This study was funded by the internal research grant of Universitas Padjadjaran and Universitas Pasundan Indonesia with grant number 16/Unpas.R/G.1/1/2020.

CONFLICT OF INTEREST

The authors confirm they have no conflicts of interest, financially or otherwise.

ACKNOWLEDGEMENTS

This study was supported by Universitas Padjadjaran and Universitas Pasundan.

REFERENCES

- Taegtmeier H, Lubrano G. Rethinking cardiac metabolism: metabolic cycles to refuel and rebuild the failing heart. *F1000Prime Rep* 2014; 6: 90. [http://dx.doi.org/10.12703/P6-90] [PMID: 25374668]
- Doenst T, Nguyen TD, Abel ED. Cardiac metabolism in heart failure: implications beyond ATP production. *Circ Res* 2013; 113(6): 709-24. [http://dx.doi.org/10.1161/CIRCRESAHA.113.300376] [PMID: 23989714]
- Fukushima A, Lopaschuk GD. Cardiac fatty acid oxidation in heart failure associated with obesity and diabetes. *Biochim Biophys Acta* 2016; 1861(10): 1525-34. [http://dx.doi.org/10.1016/j.bbali.2016.03.020] [PMID: 26996746]
- Szablewski L. Glucose transporters in healthy heart and in cardiac disease. *Int J Cardiol* 2017; 230: 70-5. [http://dx.doi.org/10.1016/j.ijcard.2016.12.083] [PMID: 28034463]
- Kolwicz SC Jr, Purohit S, Tian R. Cardiac metabolism and its interactions with contraction, growth, and survival of cardiomyocytes. *Circ Res* 2013; 113(5): 603-16. [http://dx.doi.org/10.1161/CIRCRESAHA.113.302095] [PMID: 23948585]
- Jang M, Kim SS, Lee J. Cancer cell metabolism: implications for therapeutic targets. *Exp Mol Med* 2013; 45(10): e45-8. [http://dx.doi.org/10.1038/emm.2013.85] [PMID: 24091747]
- Kaur M, Tappia PS. Metabolic shifts during cardiac hypertrophy. *Clin Lipidol* 2010; 4(6): 725-9. [http://dx.doi.org/10.2217/clp.09.73]
- Fillmore N, Levasseur JL, Fukushima A, *et al.* Uncoupling of glycolysis from glucose oxidation accompanies the development of heart failure with preserved ejection fraction. *Mol Med* 2018; 24(1): 3. [http://dx.doi.org/10.1186/s10020-018-0005-x] [PMID: 30134787]
- Zhabyeyev P, Gandhi M, Mori J, *et al.* Pressure-overload-induced heart failure induces a selective reduction in glucose oxidation at physiological afterload. *Cardiovasc Res* 2013; 97(4): 676-85. [http://dx.doi.org/10.1093/cvr/cvs424] [PMID: 23257023]
- Zhang L, Jaswal JS, Ussher JR, *et al.* Cardiac insulin-resistance and decreased mitochondrial energy production precede the development of systolic heart failure after pressure-overload hypertrophy. *Circ Heart Fail* 2013; 6(5): 1039-48. [http://dx.doi.org/10.1161/CIRCHEARTFAILURE.112.000228] [PMID: 23861485]
- Kato T, Niizuma S, Inuzuka Y, *et al.* Analysis of metabolic remodeling in compensated left ventricular hypertrophy and heart failure. *Circ Heart Fail* 2010; 3(3): 420-30. [http://dx.doi.org/10.1161/CIRCHEARTFAILURE.109.888479] [PMID: 20176713]
- Degens H, de Brouwer KF, Gilde AJ, *et al.* Cardiac fatty acid metabolism is preserved in the compensated hypertrophic rat heart. *Basic Res Cardiol* 2006; 101(1): 17-26. [http://dx.doi.org/10.1007/s00395-005-0549-0] [PMID: 16136293]
- Lei B, Lionetti V, Young ME, *et al.* Paradoxical downregulation of the glucose oxidation pathway despite enhanced flux in severe heart failure. *J Mol Cell Cardiol* 2004; 36(4): 567-76. [http://dx.doi.org/10.1016/j.yjmcc.2004.02.004] [PMID: 15081316]
- Masoud WGT, Ussher JR, Wang W, *et al.* Failing mouse hearts utilize energy inefficiently and benefit from improved coupling of glycolysis and glucose oxidation. *Cardiovasc Res* 2014; 101(1): 30-8. [http://dx.doi.org/10.1093/cvr/cvt216] [PMID: 24048945]
- Gill KS, Fernandes P, O'Donovan TR, *et al.* Glycolysis inhibition as a cancer treatment and its role in an anti-tumour immune response. *Biochim Biophys Acta* 2016; 1866(1): 87-105. [PMID: 27373814]
- Zhang Y, Yang JM. Altered energy metabolism in cancer: a unique opportunity for therapeutic intervention. *Cancer Biol Ther* 2013; 14(2): 81-9. [http://dx.doi.org/10.4161/cbt.22958] [PMID: 23192270]
- Yetkin-Arik B, Vogels IMC, Nowak-Sliwinska P, *et al.* The role of glycolysis and mitochondrial respiration in the formation and functioning of endothelial tip cells during angiogenesis. *Sci Rep* 2019; 9(1): 12608. [http://dx.doi.org/10.1038/s41598-019-48676-2] [PMID: 31471554]
- Epstein T, Gatenby RA, Brown JS. The Warburg effect as an adaptation of cancer cells to rapid fluctuations in energy demand. *PLoS One* 2017; 12(9): e0185085. [http://dx.doi.org/10.1371/journal.pone.0185085] [PMID: 28922380]
- Pfeiffer T, Schuster S, Bonhoeffer S. Cooperation and Competition in the 568 Evolution of ATP-Producing Pathways. *Science* 2001; 292(80): 504-7.
- Piao L, Fang YH, Cadete VJ, *et al.* The inhibition of pyruvate dehydrogenase kinase improves impaired cardiac function and electrical remodeling in two models of right ventricular hypertrophy: resuscitating the hibernating right ventricle. *J Mol Med (Berl)* 2010; 88(1): 47-60. [http://dx.doi.org/10.1007/s00109-009-0524-6] [PMID: 19949938]
- Piao L, Sidhu VK, Fang YH, *et al.* FOXO1-mediated upregulation of pyruvate dehydrogenase kinase-4 (PDK4) decreases glucose oxidation and impairs right ventricular function in pulmonary hypertension: therapeutic benefits of dichloroacetate. *J Mol Med (Berl)* 2013; 91(3): 333-46. [http://dx.doi.org/10.1007/s00109-012-0982-0] [PMID: 23247844]
- Lunt SY, Vander Heiden MG. Aerobic glycolysis: meeting the metabolic requirements of cell proliferation. *Annu Rev Cell Dev Biol* 2011; 27(1): 441-64. [http://dx.doi.org/10.1146/annurev-cellbio-092910-154237] [PMID: 21985671]
- Davogusto G, Dillon W, Salazar R, Vasquez H, Taegtmeier H. The Warburg Effect in the Heart: Increased Glucose Metabolism Drives Cardiomyocyte Hypertrophy in Response To Adrenergic Stimulation. *J Am Coll Cardiol* 2018; 71(11): A802. [http://dx.doi.org/10.1016/S0735-1097(18)31343-3]
- Goodwin J, Choi H, Hsieh MH, *et al.* Targeting hypoxia-inducible factor-1 α /pyruvate dehydrogenase kinase 1 axis by dichloroacetate

- suppresses bleomycin-induced pulmonary fibrosis. *Am J Respir Cell Mol Biol* 2018; 58(2): 216-31. [http://dx.doi.org/10.1165/rcmb.2016-0186OC] [PMID: 28915065]
- [25] Liu Y, Bai F, Liu N, Ouyang F, Liu Q. The Warburg effect: A new insight into atrial fibrillation. *Clin Chim Acta* 2019; 499: 4-12. [http://dx.doi.org/10.1016/j.cca.2019.08.029] [PMID: 31473195]
- [26] Stanley WC, Recchia FA, Lopaschuk GD. Myocardial substrate metabolism in the normal and failing heart. *Physiol Rev* 2005; 85(3): 1093-129. [http://dx.doi.org/10.1152/physrev.00006.2004] [PMID: 15987803]
- [27] Diane E. Handy Rita Castro Joseph Loscalzo. Cardiac Metabolism in Heart Failure - Implications beyond ATP production. *Bone* 2011; 23(1): 1-7.
- [28] Wang ZV, Li DL, Hill JA. Heart failure and loss of metabolic control. *J Cardiovasc Pharmacol* 2014; 63(4): 302-13. [http://dx.doi.org/10.1097/FJC.000000000000054] [PMID: 24336014]
- [29] van der Vusse GJ, van Bilsen M, Glatz JFC. Cardiac fatty acid uptake and transport in health and disease. *Cardiovasc Res* 2000; 45(2): 279-93. [http://dx.doi.org/10.1016/S0008-6363(99)00263-1] [PMID: 10728348]
- [30] Lopaschuk GD, Ussher JR, Folmes CDL, Jaswal JS, Stanley WC. Myocardial fatty acid metabolism in health and disease. *Physiol Rev* 2010; 90(1): 207-58. [http://dx.doi.org/10.1152/physrev.00015.2009] [PMID: 20086077]
- [31] Lionetti V, Stanley WC, Recchia FA. Modulating fatty acid oxidation in heart failure. *Cardiovasc Res* 2011; 90(2): 202-9. [http://dx.doi.org/10.1093/cvr/cvr038] [PMID: 21289012]
- [32] Nagoshi T, Yoshimura M, Rosano GM, Lopaschuk GD, Mochizuki S. Optimization of cardiac metabolism in heart failure. *Curr Pharm Des* 2011; 17(35): 3846-53. [http://dx.doi.org/10.2174/138161211798357773] [PMID: 21933140]
- [33] Tran DH, Wang ZV. Glucose Metabolism in Cardiac Hypertrophy and Heart Failure. *J Am Heart Assoc* 2019; 8(12): e012673. [http://dx.doi.org/10.1161/JAHA.119.012673] [PMID: 31185774]
- [34] Aerni-Flessner L, Abi-Jaoude M, Koenig A, Payne M, Hruz PW. GLUT4, GLUT1, and GLUT8 are the dominant GLUT transcripts expressed in the murine left ventricle. *Cardiovasc Diabetol* 2012; 11(63): 63. [http://dx.doi.org/10.1186/1475-2840-11-63] [PMID: 22681646]
- [35] Abel ED. Glucose transport in the heart. *Front Biosci* 2004; 9(1): 201-15. [http://dx.doi.org/10.2741/1216] [PMID: 14766360]
- [36] Tian R, Abel ED. Responses of GLUT4-deficient hearts to ischemia underscore the importance of glycolysis. *Circulation* 2001; 103(24): 2961-6. [http://dx.doi.org/10.1161/01.CIR.103.24.2961] [PMID: 11413087]
- [37] Azevedo PS, Minicucci MF, Santos PP, Paiva SAR, Zornoff LAM. Energy metabolism in cardiac remodeling and heart failure. *Cardiol Rev* 2013; 21(3): 135-40. [http://dx.doi.org/10.1097/CRD.0b013e318274956d] [PMID: 22990373]
- [38] Zimmer HG. Regulation of and intervention into the oxidative pentose phosphate pathway and adenine nucleotide metabolism in the heart. *Mol Cell Biochem* 1996; 160-161: 101-9. [http://dx.doi.org/10.1007/BF00240038] [PMID: 8901462]
- [39] Jain M, Brenner DA, Cui L, *et al.* Glucose-6-phosphate dehydrogenase modulates cytosolic redox status and contractile phenotype in adult cardiomyocytes. *Circ Res* 2003; 93(2): e9-e16. [http://dx.doi.org/10.1161/01.RES.0000083489.83704.76] [PMID: 12829617]
- [40] Wu G, Fang Y-Z, Yang S, Lupton JR, Turner ND. Glutathione metabolism and its implications for health. *J Nutr* 2004; 134(3): 489-92. [http://dx.doi.org/10.1093/jn/134.3.489] [PMID: 14988435]
- [41] Srivastava SK, Ramana KV, Bhatnagar A. Role of aldose reductase and oxidative damage in diabetes and the consequent potential for therapeutic options. *Endocr Rev* 2005; 26(3): 380-92. [http://dx.doi.org/10.1210/er.2004-0028] [PMID: 15814847]
- [42] Sharavana G, Joseph GS, Baskaran V. Lutein attenuates oxidative stress markers and ameliorates glucose homeostasis through polyol pathway in heart and kidney of STZ-induced hyperglycemic rat model. *Eur J Nutr* 2017; 56(8): 2475-85. [http://dx.doi.org/10.1007/s00394-016-1283-0] [PMID: 27488609]
- [43] Banerjee PS, Lagerlöf O, Hart GW. Roles of O-GlcNAc in chronic diseases of aging. *Mol Aspects Med* 2016; 51: 1-15. [http://dx.doi.org/10.1016/j.mam.2016.05.005] [PMID: 27259471]
- [44] Lunde IG, Aronsen JM, Kvaløy H, *et al.* Cardiac O-GlcNAc signaling is increased in hypertrophy and heart failure. *Physiol Genomics* 2012; 44(2): 162-72. [http://dx.doi.org/10.1152/physiolgenomics.00016.2011] [PMID: 22128088]
- [45] Cotter DG, Schugar RC, Crawford PA. Ketone body metabolism and cardiovascular disease. *Am J Physiol - Hear Circ Physiol* 2013; 304(8): 1060-76. [http://dx.doi.org/10.1152/ajpheart.00646.2012]
- [46] Mohan M, Al-Talabany S, McKinnie A, *et al.* A randomized controlled trial of metformin on left ventricular hypertrophy in patients with coronary artery disease without diabetes: the MET-REMODEL trial. *Eur Heart J* 2019; 40(41): 3409-17. [http://dx.doi.org/10.1093/eurheartj/ehz203] [PMID: 30993313]
- [47] Janardhan A, Chen J, Crawford PA. Altered systemic ketone body metabolism in advanced heart failure. *Tex Heart Inst J* 2011; 38(5): 533-8. [PMID: 22163128]
- [48] Chen Z, Liu M, Li L, Chen L. Involvement of the Warburg effect in non-tumor diseases processes. *J Cell Physiol* 2018; 233(4): 2839-49. [http://dx.doi.org/10.1002/jcp.25998] [PMID: 28488732]
- [49] Depré C, Rider MH, Hue L. Mechanisms of control of heart glycolysis. *Eur J Biochem* 1998; 258(2): 277-90. [http://dx.doi.org/10.1046/j.1432-1327.1998.2580277.x] [PMID: 9874192]
- [50] Leong HS, Grist M, Parsons H, Wambolt RB, Lopaschuk GD, Brownsey R, *et al.* Accelerated rates of glycolysis in the hypertrophied heart: Are they 657 a methodological artifact? *Am J Physiol - Endocrinol Metab* 2002; 282(545-5): 1039-45.
- [51] Toyama EQ, Herzig S, Courchet J, Jr TLL, Kranias EG, Prevention C. AMPK in Cardiovascular Diseases. *Science* (80-) 2016; 107(6270): 275-82.
- [52] John P, Konhilas YL. AMP-Activated Protein Kinase Signalling in Cancer and Cardiac Hypertrophy. *Cardiovasc Pharmacol Open Access* 2015; 04(03)
- [53] Frey N, Olson EN. Cardiac hypertrophy: the good, the bad, and the ugly. *Annu Rev Physiol* 2003; 65(1): 45-79. [http://dx.doi.org/10.1146/annurev.physiol.65.092101.142243] [PMID: 12524460]
- [54] Nascimben L, Ingwall JS, Lorell BH, *et al.* Mechanisms for increased glycolysis in the hypertrophied rat heart. *Hypertension* 2004; 44(5): 662-7. [http://dx.doi.org/10.1161/01.HYP.0000144292.69599.0c] [PMID: 15466668]
- [55] Warburg O. On the origin of cancer cells. *Science* (80-) 1956; 123(3191): 309-14. [http://dx.doi.org/10.1126/science.123.3191.309]
- [56] Hodge CJ, Apkarian AV. Understanding the Warburg Effect: The Metabolic Requirements of Cell Proliferation. *Science* (80-) 2009; 5(4): 363-97.
- [57] Harada M, Tadevosyan A, Qi X, *et al.* Atrial Fibrillation Activates AMP-Dependent Protein Kinase and its Regulation of Cellular Calcium Handling: Potential Role in Metabolic Adaptation and Prevention of Progression. *J Am Coll Cardiol* 2015; 66(1): 47-58. [http://dx.doi.org/10.1016/j.jacc.2015.04.056] [PMID: 26139058]
- [58] Dai DF, Hsieh EJ, Liu Y, *et al.* Mitochondrial proteome remodelling in pressure overload-induced heart failure: the role of mitochondrial oxidative stress. *Cardiovasc Res* 2012; 93(1): 79-88. [http://dx.doi.org/10.1093/cvr/cvr274] [PMID: 22012956]
- [59] Ozcan C, Li Z, Kim G, Jeevanandam V, Uriel N. Molecular Mechanism of the Association Between Atrial Fibrillation and Heart Failure Includes Energy Metabolic Dysregulation Due to Mitochondrial Dysfunction. *J Card Fail* 2019; 25(11): 911-20. [http://dx.doi.org/10.1016/j.cardfail.2019.08.005] [PMID: 31415862]
- [60] Ussher JR, Koves TR, Jaswal JS, *et al.* Insulin-stimulated cardiac glucose oxidation is increased in high-fat diet-induced obese mice lacking malonyl CoA decarboxylase. *Diabetes* 2009; 58(8): 1766-75. [http://dx.doi.org/10.2337/db09-0011] [PMID: 19478144]
- [61] Umbarawan Y, Syamsunarno MRAA, Koitabashi N, *et al.* Myocardial fatty acid uptake through CD36 is indispensable for sufficient bioenergetic metabolism to prevent progression of pressure overload-induced heart failure. *Sci Rep* 2018; 8(1): 12035. [http://dx.doi.org/10.1038/s41598-018-30616-1] [PMID: 30104639]
- [62] Umbarawan Y, Syamsunarno MRAA, Koitabashi N, *et al.* Glucose is preferentially utilized for biomass synthesis in pressure-overloaded hearts: evidence from fatty acid-binding protein-4 and -5 knockout

- mice. *Cardiovasc Res* 2018; 114(8): 1132-44. [http://dx.doi.org/10.1093/cvr/cvy063] [PMID: 29554241]
- [63] Nahle Z, Hsieh M, Pietka T, *et al.* CD36-dependent regulation of muscle FoxO1 and PDK4 in the PPAR δ/β -mediated adaptation to metabolic stress. *J Biol Chem* 2008; 283(21): 14317-26. [http://dx.doi.org/10.1074/jbc.M706478200] [PMID: 18308721]
- [64] Wakili R, Voigt N, Kääh S, Dobrev D, Nattel S. Recent advances in the molecular pathophysiology of atrial fibrillation. *J Clin Invest* 2011; 121(8): 2955-68. [http://dx.doi.org/10.1172/JCI46315] [PMID: 21804195]
- [65] Tu T, Zhou S, Liu Z, Li X, Liu Q. Quantitative proteomics of changes in energy metabolism-related proteins in atrial tissue from valvular disease patients with permanent atrial fibrillation. *Circ J* 2014; 78(4): 993-1001. [http://dx.doi.org/10.1253/circj.CJ-13-1365] [PMID: 24553264]
- [66] Tsuboi M, Hisatome I, Morisaki T, *et al.* Mitochondrial DNA deletion associated with the reduction of adenine nucleotides in human atrium and atrial fibrillation. *Eur J Clin Invest* 2001; 31(6): 489-96. [http://dx.doi.org/10.1046/j.1365-2362.2001.00844.x] [PMID: 11422398]
- [67] Woolf EC, Scheck AC. The Ketogenic Diet for the Adjuvant Treatment of Malignant Brain Tumors. *Bioactive Nutraceuticals and Dietary Supplements in Neurological and Brain Disease: Prevention and Therapy*. Elsevier Inc. 2015; pp. 125-35. [http://dx.doi.org/10.1016/B978-0-12-411462-3.00013-8]
- [68] Vazquez A, Liu J, Zhou Y, Oltvai ZN. Catabolic efficiency of aerobic glycolysis: the Warburg effect revisited. *BMC Syst Biol* 2010; 4: 58. [http://dx.doi.org/10.1186/1752-0509-4-58] [PMID: 20459610]
- [69] Vazquez A, Kamphorst JJ, Markert EK, Schug ZT, Tardito S, Gottlieb E. Cancer metabolism at a glance. *J Cell Sci* 2016; 129(18): 3367-73. [http://dx.doi.org/10.1242/jcs.181016] [PMID: 27635066]
- [70] Fantin VR, St-Pierre J, Leder P. Attenuation of LDH-A expression uncovers a link between glycolysis, mitochondrial physiology, and tumor maintenance. *Cancer Cell* 2006; 9(6): 425-34. [http://dx.doi.org/10.1016/j.ccr.2006.04.023] [PMID: 16766262]
- [71] Papandreou I, Cairns RA, Fontana L, Lim AL, Denko NC. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab* 2006; 3(3): 187-97. [http://dx.doi.org/10.1016/j.cmet.2006.01.012] [PMID: 16517406]
- [72] Kim JW, Tchernyshyov I, Semenza GL, Dang CV. HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia. *Cell Metab* 2006; 3(3): 177-85. [http://dx.doi.org/10.1016/j.cmet.2006.02.002] [PMID: 16517405]
- [73] Holness MJ, Sugden MC. Regulation of pyruvate dehydrogenase complex activity by reversible phosphorylation. *Biochem Soc Trans* 2003; 31(Pt 6): 1143-51. [http://dx.doi.org/10.1042/bst0311143] [PMID: 14641014]
- [74] Elstrom RL, Bauer DE, Buzzai M, *et al.* Akt stimulates aerobic glycolysis in cancer cells. *Cancer Res* 2004; 64(11): 3892-9. [http://dx.doi.org/10.1158/0008-5472.CAN-03-2904] [PMID: 15172999]
- [75] Kolwicz SC Jr, Tian R. Glucose metabolism and cardiac hypertrophy. *Cardiovasc Res* 2011; 90(2): 194-201. [http://dx.doi.org/10.1093/cvr/cvr071] [PMID: 21502371]
- [76] Morimoto S, Goto T. Role of troponin I isoform switching in determining the pH sensitivity of Ca(2+) regulation in developing rabbit cardiac muscle. *Biochem Biophys Res Commun* 2000; 267(3): 912-7. [http://dx.doi.org/10.1006/bbrc.1999.2068] [PMID: 10673390]
- [77] Sansbury BE, Riggs DW, Brainard RE, Salabei JK, Jones SP, Hill BG. Responses of hypertrophied myocytes to reactive species: implications for glycolysis and electrophile metabolism. *Biochem J* 2011; 435(2): 519-28. [http://dx.doi.org/10.1042/BJ20101390] [PMID: 21275902]
- [78] Jaswal JS, Keung W, Wang W, Ussher JR, Lopaschuk GD. Targeting fatty acid and carbohydrate oxidation--a novel therapeutic intervention in the ischemic and failing heart. *Biochim Biophys Acta* 2011; 1813(7): 1333-50. [http://dx.doi.org/10.1016/j.bbamer.2011.01.015] [PMID: 21256164]
- [79] Bauer DE, Harris MH, Plas DR, *et al.* Cytokine stimulation of aerobic glycolysis in hematopoietic cells exceeds proliferative demand. *FASEB J* 2004; 18(11): 1303-5. [http://dx.doi.org/10.1096/fj.03-1001fje] [PMID: 15180958]
- [80] Hu HJ, Zhang C, Tang ZH, Qu SL, Jiang ZS. Regulating the Warburg effect on metabolic stress and myocardial fibrosis remodeling and atrial intracardiac waveform activity induced by atrial fibrillation. *Biochem Biophys Res Commun* 2019; 516(3): 653-60. [http://dx.doi.org/10.1016/j.bbrc.2019.06.055] [PMID: 31242971]
- [81] Lopaschuk GD, Jaswal JS. Energy metabolic phenotype of the cardiomyocyte during development, differentiation, and postnatal maturation. *J Cardiovasc Pharmacol* 2010; 56(2): 130-40. [http://dx.doi.org/10.1097/FJC.0b013e3181e74a14] [PMID: 20505524]
- [82] Balamurugan K. HIF-1 at the crossroads of hypoxia, inflammation, and cancer. *Int J Cancer* 2016; 138(5): 1058-66. [http://dx.doi.org/10.1002/ijc.29519] [PMID: 25784597]
- [83] Liu Y, Shi Q, Ma Y, Liu Q. The role of immune cells in atrial fibrillation. *J Mol Cell Cardiol* 2018; 123: 198-208. [http://dx.doi.org/10.1016/j.yjmcc.2018.09.007] [PMID: 30267749]
- [84] Semenza GL. Hypoxia-inducible factor 1 and cardiovascular disease. *Annu Rev Physiol* 2014; 76(1): 39-56. [http://dx.doi.org/10.1146/annurev-physiol-021113-170322] [PMID: 23988176]
- [85] Das C, Hansen KC and Tyler JK LMS. PKM2 Regulates the Warburg Effect and Promotes HMGB1 Release in Sepsis. *Nat Commun* 2017; 176(3): 139-48.
- [86] Li Y huan, Li X feng, Liu J tao, Wang H, Fan L lu, Li J, *et al.* PKM2, a potential target for regulating cancer. *Gene* 2018; 668(2017): 48-53.
- [87] Rees ML, Subramaniam J, Li Y, Hamilton DJ, Frazier OH, Taegtmeier H. A PKM2 signature in the failing heart. *Biochem Biophys Res Commun* 2015; 459(3): 430-6. [http://dx.doi.org/10.1016/j.bbrc.2015.02.122] [PMID: 25735978]
- [88] Fillmore N, Mori J, Lopaschuk GD. Mitochondrial fatty acid oxidation alterations in heart failure, ischaemic heart disease and diabetic cardiomyopathy. *Br J Pharmacol* 2014; 171(8): 2080-90. [http://dx.doi.org/10.1111/bph.12475] [PMID: 24147975]
- [89] Taegtmeier H, Sen S, Vela D. Return to the fetal gene program: a suggested metabolic link to gene expression in the heart. *Ann N Y Acad Sci* 2010; 1188: 191-8. [http://dx.doi.org/10.1111/j.1749-6632.2009.05100.x] [PMID: 20201903]
- [90] Lopaschuk GD, Spafford MA. Energy substrate utilization by isolated working hearts from newborn rabbits. *Am J Physiol - Hear Circ Physiol* 1990; 258(5): H1275-80. [http://dx.doi.org/10.1152/ajpheart.1990.258.5.H1274]
- [91] Lopaschuk GD, Spafford MA, Marsh DR. Glycolysis is predominant source of myocardial ATP production immediately after birth. *Am J Physiol - Hear Circ Physiol* 1991; 261(6 30-6): 1698-705. [http://dx.doi.org/10.1152/ajpheart.1991.261.6.H1698]
- [92] Krishnan J, Suter M, Windak R, *et al.* Activation of a HIF1 α -PPARgamma axis underlies the integration of glycolytic and lipid anabolic pathways in pathologic cardiac hypertrophy. *Cell Metab* 2009; 9(6): 512-24. [http://dx.doi.org/10.1016/j.cmet.2009.05.005] [PMID: 19490906]
- [93] Atherton HJ, Dodd MS, Heather LC, *et al.* Role of pyruvate dehydrogenase inhibition in the development of hypertrophy in the hyperthyroid rat heart: a combined magnetic resonance imaging and hyperpolarized magnetic resonance spectroscopy study. *Circulation* 2011; 123(22): 2552-61. [http://dx.doi.org/10.1161/CIRCULATIONAHA.110.011387] [PMID: 21606392]
- [94] Wang J, Xu J, Wang Q, *et al.* Reduced cardiac fructose 2,6 biphosphate increases hypertrophy and decreases glycolysis following aortic constriction. *PLoS One* 2013; 8(1): e53951. [http://dx.doi.org/10.1371/journal.pone.0053951] [PMID: 23308291]
- [95] Zhu H-L, Gu C-H, Tao C, Xiong H-Y, Sun G-C. The role of "glycolysis- uncoupling of glucose oxidation" in the progression of chronic cardiac hypertrophy *Med J Chin PLA* 2013; 38
- [96] Beer M, Seyfarth T, Sandstede J, *et al.* Absolute concentrations of high-energy phosphate metabolites in normal, hypertrophied, and failing human myocardium measured noninvasively with (31)P-SLOOP magnetic resonance spectroscopy. *J Am Coll Cardiol* 2002; 40(7): 1267-74. [http://dx.doi.org/10.1016/S0735-1097(02)02160-5] [PMID: 12383574]
- [97] Sankaralingam S, Lopaschuk GD. Cardiac energy metabolic alterations in pressure overload-induced left and right heart failure (2013 Grover Conference Series). *Pulm Circ* 2015; 5(1): 15-28. [http://dx.doi.org/10.1086/679608] [PMID: 25992268]
- [98] Liu Q, Docherty JC, Rendell JCT, Clanachan AS, Lopaschuk GD. High levels of fatty acids delay the recovery of intracellular pH and cardiac efficiency in post-ischemic hearts by inhibiting glucose oxidation. *J Am Coll Cardiol* 2002; 39(4): 718-25.

- [http://dx.doi.org/10.1016/S0735-1097(01)01803-4] [PMID: 11849874]
- [99] Ussher JR, Wang W, Gandhi M, *et al.* Stimulation of glucose oxidation protects against acute myocardial infarction and reperfusion injury. *Cardiovasc Res* 2012; 94(2): 359-69. [http://dx.doi.org/10.1093/cvr/cvs129] [PMID: 22436846]
- [100] Sun XQ, Zhang R, Zhang HD, *et al.* Reversal of right ventricular remodeling by dichloroacetate is related to inhibition of mitochondria-dependent apoptosis. *Hypertens Res* 2016; 39(5): 302-11. [http://dx.doi.org/10.1038/hr.2015.153] [PMID: 26763846]
- [101] Bersin RM, Wolfe C, Kwasman M, *et al.* Improved hemodynamic function and mechanical efficiency in congestive heart failure with sodium dichloroacetate. *J Am Coll Cardiol* 1994; 23(7): 1617-24. [http://dx.doi.org/10.1016/0735-1097(94)90665-3] [PMID: 8195522]
- [102] Beauloye C, Bertrand L, Horman S, Hue L. AMPK activation, a preventive therapeutic target in the transition from cardiac injury to heart failure. *Cardiovasc Res* 2011; 90(2): 224-33. [http://dx.doi.org/10.1093/cvr/cvr034] [PMID: 21285292]
- [103] Gundewar S, Calvert JW, Jha S, *et al.* Activation of AMP-activated protein kinase by metformin improves left ventricular function and survival in heart failure. *Circ Res* 2009; 104(3): 403-11. [http://dx.doi.org/10.1161/CIRCRESAHA.108.190918] [PMID: 19096023]
- [104] Arad M, Seidman CE, Seidman JG. AMP-activated protein kinase in the heart: role during health and disease. *Circ Res* 2007; 100(4): 474-88. [http://dx.doi.org/10.1161/01.RES.0000258446.23525.37] [PMID: 17332438]
- [105] Maria Z, Campolo AR, Lacombe VA. Diabetes alters the expression and translocation of the insulin-sensitive glucose transporters 4 and 8 in the atria. *PLoS One* 2015; 10(12): e0146033 [http://dx.doi.org/10.1371/journal.pone.0146033] [PMID: 26720696]
- [106] Owen MR, Doran E, Halestrap AP. Evidence that metformin exerts its anti-diabetic effects through inhibition of complex I of the mitochondrial respiratory chain. *Biochem J* 2000; 348(Pt 3): 607-14. [http://dx.doi.org/10.1042/bj3480607] [PMID: 10839993]
- [107] Hawley SA, Gadalla AE, Olsen GS, Hardie DG. The antidiabetic drug metformin activates the AMP-activated protein kinase cascade via an adenine nucleotide-independent mechanism. *Diabetes* 2002; 51(8): 2420-5. [http://dx.doi.org/10.2337/diabetes.51.8.2420] [PMID: 12145153]
- [108] Sasaki H, Asanuma H, Fujita M, *et al.* Metformin prevents progression of heart failure in dogs: role of AMP-activated protein kinase. *Circulation* 2009; 119(19): 2568-77. [http://dx.doi.org/10.1161/CIRCULATIONAHA.108.798561] [PMID: 19414638]
- [109] Liu Y, Bai F, Liu N, *et al.* Metformin improves lipid metabolism and reverses the Warburg effect in a canine model of chronic atrial fibrillation. *BMC Cardiovasc Disord* 2020; 20(1): 50. [http://dx.doi.org/10.1186/s12872-020-01359-7] [PMID: 32013885]
- [110] Aguilar D, Chan W, Bozkurt B, Ramasubbu K, Deswal A. Metformin Use and Mortality in Ambulatory Patients With Diabetes and Heart Failure. *J Card Fail* 2010; 16(8): S83-4. [http://dx.doi.org/10.1016/j.cardfail.2010.06.292] [PMID: 20952583]
- [111] Kanamori H, Takemura G, Goto K, *et al.* Resveratrol reverses remodeling in hearts with large, old myocardial infarctions through enhanced autophagy-activating AMP kinase pathway. *Am J Pathol* 2013; 182(3): 701-13. [http://dx.doi.org/10.1016/j.ajpath.2012.11.009] [PMID: 23274061]
- [112] Baczko I, Liknes D, Yang W, *et al.* Characterization of a novel multifunctional resveratrol derivative for the treatment of atrial fibrillation. *Br J Pharmacol* 2014; 171(1): 92-106. [http://dx.doi.org/10.1111/bph.12409] [PMID: 24102184]
- [113] Raj P, Louis XL, Thandapilly SJ, Movahed A, Zieroth S, Nettiadan T. Potential of resveratrol in the treatment of heart failure. *Life Sci* 2014; 95(2): 63-71. [http://dx.doi.org/10.1016/j.lfs.2013.12.011] [PMID: 24361400]
- [114] Sarma S, Ardehali H, Gheorghide M. Enhancing the metabolic substrate: PPAR-alpha agonists in heart failure. *Heart Fail Rev* 2012; 17(1): 35-43. [http://dx.doi.org/10.1007/s10741-010-9208-0] [PMID: 21104312]
- [115] Campbell FM, Kozak R, Wagner A, Altarejos JY, Dyck JRB, Belke DD, *et al.* A role for peroxisome proliferator-activated receptor α (PPAR α) in the control of cardiac malonyl-CoA levels: Reduced fatty acid oxidation rates and increased glucose oxidation rates in the hearts of mice lacking PPAR α are associated with higher concentra. *J Biol Chem* 2002; 277(6): 4098-103. [http://dx.doi.org/10.1074/jbc.M106054200] [PMID: 11734553]
- [116] Hafstad AD, Khalid AM, Hagve M, *et al.* Cardiac peroxisome proliferator-activated receptor- α activation causes increased fatty acid oxidation, reducing efficiency and post-ischaemic functional loss. *Cardiovasc Res* 2009; 83(3): 519-26. [http://dx.doi.org/10.1093/cvr/cvp132] [PMID: 19398469]
- [117] Li P, Luo S, Pan C, Cheng X. Modulation of fatty acid metabolism is involved in the alleviation of isoproterenol-induced rat heart failure by fenofibrate. *Mol Med Rep* 2015; 12(6): 7899-906. [http://dx.doi.org/10.3892/mmr.2015.4466] [PMID: 26497978]
- [118] Liu GZ, Hou TT, Yuan Y, *et al.* Fenofibrate inhibits atrial metabolic remodelling in atrial fibrillation through PPAR- α /sirtuin 1/PGC-1 α pathway. *Br J Pharmacol* 2016; 173(6): 1095-109. [http://dx.doi.org/10.1111/bph.13438] [PMID: 26787506]
- [119] Labinskyy V, Bellomo M, Chandler MP, *et al.* Chronic activation of peroxisome proliferator-activated receptor- α with fenofibrate prevents alterations in cardiac metabolic phenotype without changing the onset of decompensation in pacing-induced heart failure. *J Pharmacol Exp Ther* 2007; 321(1): 165-71. [http://dx.doi.org/10.1124/jpet.106.116871] [PMID: 17215446]
- [120] Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, *et al.* 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. 2019; 140: pp. Circulation. 596-646. [http://dx.doi.org/10.1056/NEJMoa1200303] [PMID: 23432189]
- [121] Estruch R, Ros E, Salas-Salvadó J, *et al.* Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* 2013; 368(14): 1279-90. [http://dx.doi.org/10.1056/NEJMoa1200303] [PMID: 23432189]
- [122] Estruch R, Martínez-González MA, Corella D, *et al.* Effects of a Mediterranean-style diet on cardiovascular risk factors: a randomized trial. *Ann Intern Med* 2006; 145(1): 1-11. [http://dx.doi.org/10.7326/0003-4819-145-1-200607040-00004] [PMID: 16818923]
- [123] Yu E, Malik VS, Hu FB. Cardiovascular Disease Prevention by Diet Modification: JACC Health Promotion Series. *J Am Coll Cardiol* 2018; 72(8): 914-26. [http://dx.doi.org/10.1016/j.jacc.2018.02.085] [PMID: 30115231]
- [124] Akbaraly T, Würtz P, Singh-Manoux A, *et al.* Association of circulating metabolites with healthy diet and risk of cardiovascular disease: analysis of two cohort studies. *Sci Rep* 2018; 8(1): 8620. [http://dx.doi.org/10.1038/s41598-018-26441-1] [PMID: 29872056]
- [125] Fiaccavento R, Carotenuto F, Minieri M, *et al.* Alpha-linolenic acid-enriched diet prevents myocardial damage and expands longevity in cardiomyopathic hamsters. *Am J Pathol* 2006; 169(6): 1913-24. [http://dx.doi.org/10.2353/ajpath.2006.051320] [PMID: 17148657]
- [126] Ordovas JM. Nutrigenetics, plasma lipids, and cardiovascular risk. *J Am Diet Assoc* 2006; 106(7): 1074-81. [http://dx.doi.org/10.1016/j.jada.2006.04.016] [PMID: 16815124]
- [127] Haro D, Marrero PF, Relat J. Nutritional regulation of gene expression: Carbohydrate-, fat- and amino acid-dependent modulation of transcriptional activity. *Int J Mol Sci* 2019; 20(6): E1386. [http://dx.doi.org/10.3390/ijms20061386] [PMID: 30893897]
- [128] Hellstrand S, Sonestedt E, Ericson U, *et al.* Intake levels of dietary long-chain PUFAs modify the association between genetic variation in FADS and LDL-C. *J Lipid Res* 2012; 53(6): 1183-9. [http://dx.doi.org/10.1194/jlr.P023721] [PMID: 22451038]
- [129] Crescenti A, Solà R, Valls RM, *et al.* Cocoa Consumption Alters the Global DNA Methylation of Peripheral Leukocytes in Humans with Cardiovascular Disease Risk Factors: A Randomized Controlled Trial. *PLoS One* 2013; 8(6): e65744 [http://dx.doi.org/10.1371/journal.pone.0065744] [PMID: 23840361]
- [130] Feil R, Fraga MF. Epigenetics and the environment: emerging patterns and implications. *Nat Rev Genet* 2012; 13(2): 97-109. [http://dx.doi.org/10.1038/nrg3142] [PMID: 22215131]
- [131] Khan N, Monagas M, Andres-Lacueva C, *et al.* Regular consumption of cocoa powder with milk increases HDL cholesterol and reduces oxidized LDL levels in subjects at high-risk of cardiovascular disease. *Nutr Metab Cardiovasc Dis* 2012; 22(12): 1046-53. [http://dx.doi.org/10.1016/j.numecd.2011.02.001] [PMID: 21550218]
- [132] Sarriá B, Martínez-López S, Sierra-Cinos JL, García-Diz L, Mateos R, Bravo L. Regular consumption of a cocoa product improves the cardiometabolic profile in healthy and moderately hypercholesterolaemic adults. *Br J Nutr* 2014; 111(1): 122-34. [http://dx.doi.org/10.1017/S000711451300202X] [PMID: 23823716]
- [133] Yang HT, Lee M, Hong KS, Ovbiagele B, Saver JL. Efficacy of folic acid supplementation in cardiovascular disease prevention: an updated

- meta-analysis of randomized controlled trials. *Eur J Intern Med* 2012; 23(8): 745-54.
[http://dx.doi.org/10.1016/j.ejim.2012.07.004] [PMID: 22884409]
- [134] Dobrzyn P, Pyrkowska A, Duda MK, *et al.* Expression of lipogenic genes is upregulated in the heart with exercise training-induced but not pressure overload-induced left ventricular hypertrophy. *Am J Physiol Endocrinol Metab* 2013; 304(12): E1348-58.
[http://dx.doi.org/10.1152/ajpendo.00603.2012] [PMID: 23632628]
- [135] Strøm CC, Aplin M, Ploug T, *et al.* Expression profiling reveals differences in metabolic gene expression between exercise-induced cardiac effects and maladaptive cardiac hypertrophy. *FEBS J* 2005; 272(11): 2684-95.
[http://dx.doi.org/10.1111/j.1742-4658.2005.04684.x] [PMID: 15943803]

© 2021 Kuspriyanti *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.