



# The Open Orthopaedics Journal

Content list available at: <https://openorthopaedicsjournal.com>



## RESEARCH ARTICLE

### Patellar Tendon Rupture Following Total Knee Arthroplasty

Steven T. Heer<sup>1</sup>, James O'Dowd<sup>2</sup>, Rebecca R. Butler<sup>3</sup>, David O. Dewitt<sup>4</sup>, Gaurav Khanna<sup>4</sup> and Raffy Mirzayan<sup>4,\*</sup>

<sup>1</sup>Tulane School of Medicine, New Orleans, LA, USA

<sup>2</sup>Department of Orthopaedic Surgery, USC Keck School of Medicine, Los Angeles, CA, USA

<sup>3</sup>Department of Research and Evaluation, Kaiser Permanente, Pasadena, CA, USA

<sup>4</sup>Department of Orthopaedic Surgery, Kaiser Permanente Southern California, Baldwin Park, CA, USA

#### Abstract:

#### Background:

Patellar tendon rupture following Total Knee Arthroplasty (TKA) is rare. There is no consensus on optimal treatment.

#### Methods:

All patients who underwent a primary repair of a traumatic patellar tendon rupture following a TKA between 2008 and 2016, were retrospectively reviewed. Patient information, implant, repair type (anchor vs. bone tunnel), graft use, and complications were recorded.

#### Results:

Twenty-six patients met our inclusion criteria. The average age was 69.7±11 years. There were 19 females (73.1%). The average time from TKA to PT rupture was 13.6 months (range: 0- 135 months). The average incidence was 62.32 per 100,000 TKA. PT was repaired with anchors (A) in 9 (4 with a graft) and trans-osseous tunnels (TO) in 12 (5 required graft), and 5 with other methods. There was a significant improvement in KSS from 61 to 83 (P=0.023). There was a significant difference in time from PT tear to surgery in patients with grafts (42 days) and those without grafts (6 days) (P<0.001). Compared to A repair, TO had 2.39 times odds of re-tear (95% CI: 0.38,15.4; P=0.354) and 1.37 times odds of infection (95% CI:0.074,25.6; P=0.83). Repairs with a graft had a 1.90 times odds of re-tear (95% CI: 0.29, 12.19; P=0.49) and 6.3 time odds of infection (95% CI 0.26, 166.7; P=0.25).

#### Conclusion:

Surgical repair of PT tears following TKA leads to significant clinical improvement, regardless of the fixation method or graft use. We found no difference in outcomes between A and TO repairs and or with graft use.

**Keywords:** Total knee arthroplasty, Patellar tendon rupture, Anchor, Bone tunnel, Retrospectively, Fixation method.

#### Article History

Received: August 09, 2019

Revised: November 16, 2019

Accepted: November 21, 2019

## 1. INTRODUCTION

Patella Tendon (PT) rupture is a rare complication following Total Knee Arthroplasty (TKA) and is estimated to occur in 0.17-1.0% of patients [1 - 3]. Most of these injuries are a result of avulsion from the tibial tubercle insertion, with a minority being mid-substance or infra-patellar avulsions [2]. This complication is thought to be due to compromised blood supply to the extensor mechanism that occurs during TKA [4]. Studies exploring outcomes of patellar tendon rupture repair have shown mixed results [1, 2, 5, 6]. Presently, the two main repair types include suture anchors (A) and transosseous (TO)

repairs and both can be augmented by an allograft in the setting of poor tissue quality. Due to the low incidence of this complication, it is still not known which technique is superior. The purpose of this study to compare the outcomes and complications of these techniques.

## 2. MATERIALS AND METHODS

After obtaining approval from our Institutional Review Board, we searched through surgical database for patellar repair surgeries performed between January 1, 2008, and December 31, 2016. Our integrated health care system includes 14 medical centers with over 100 orthopaedic surgeons. Our system does not catalogue surgical cases by CPT codes, rather by interfacility codes. We, therefore, searched for "patellar

\* Address correspondence to this author at the Department of Orthopaedic Surgery, Kaiser Permanente Southern California, Baldwin Park, CA, USA; E-mails: [lakersdoc@gmail.com](mailto:lakersdoc@gmail.com), [raffy.mirzayan@kp.org](mailto:raffy.mirzayan@kp.org)

tendon repair” and “open repair of tendon, knee” in our database to identify patients who had undergone a patellar tendon repair. A retrospective chart review was then performed to identify those patients who had a prior total knee arthroplasty (TKA) who underwent surgical repair of a patellar tendon tear. The operative reports, progress reports, and history and physical examination, physical therapy notes, and emergency room notes were reviewed and relevant data were collected.

Data gathered included age, gender, side of surgery, BMI, length of time between TKA and patellar tendon rupture (months). In addition, pre- and post-operative Knee Society Scores (KSS) [7, 8], range of motion including extensor lag were also recorded. We were unable to collect pre-operative KSS for 10 patients and post-operative KSS were only used for patients with a minimum of 3 months post-PT repair, leaving 12 patients with both scores.

The information recorded from the operative report included tourniquet time, repair type, patellar thickness after resection (mm) (measured using the internal digital software (Philips iSite Radiology, Version 3.6.120), patellar polyethylene thickness (mm), implant manufacturer, and use of auto- or allograft. Complications such as re-rupture, infection, and thromboembolic events were also recorded.

Means and medians of continuous variables were

compared using t-tests and Kruskal-Wallis tests as deemed appropriate. Unadjusted odd ratios were calculated using logistic regression. All P values were 2-sided with an alpha = 0.05. Statistical analyses were performed with SAS 9.3 (SAS Institute, Cary, NC).

### 3. RESULTS

#### 3.1. Patient Demographics

Four hundred PT repairs were identified during our database search and from this, 26 patients were identified who sustained a PT disruption after a TKA. During our study period, 41,722 total knee arthroplasties were performed in our institution. The average incidence was 62.32 per 100,000 TKA. The average annual incidence of PT tears with TKA can be seen in Fig. (1).

The average age at the time of surgery was 69.7 years (range, 53 to 88) and there were 19 females (73.1%). Seventeen patients were Caucasian, three Asian, three Hispanic, and two African American. Twenty-five patients underwent a TKA for osteoarthritis and one patient for posttraumatic degenerative reasons. The average time between TKA to PT tear was 13.6 months (range 0 to 135 months). Patients with predisposing factors included fourteen taking statins, six with diabetes, and one with stage three chronic kidney disease (Table 1).




Fig. (1). Annual Incidence of PT tears.

Table 1. Chronic kidney disease.

Age	Gender	Mechanism of Injury	Time from Injury to Surgery (days)	Prosthesis	Notes
59	F	Fall	0	Zimmer Natural Knee	Diabetic, Taking statins, had re-tear
88	M	Progressive	18	J&J Depuy Sigma	Had re-tear
67	F	Fall	25	J&J Depuy Sigma	-
80	F	Fall	31	J&J Depuy Sigma	Had re-tear
57	F	During TKA	0	Zimmer Natural Knee (Revision)	Had infection
81	F	Fall	16	J&J Depuy Sigma (Revision)	Taking statins

(Table 1) contd.....

Age	Gender	Mechanism of Injury	Time from Injury to Surgery (days)	Prosthesis	Notes
64	F	Spontaneous	117	Zimmer Natural Knee (Revision)	Taking statins, had re-tear, had DVT
81	M	Getting up from sitting	65	Zimmer Natural Knee	Taking statins
53	F	Fall	21	J&J Depuy Sigma (Revision)	Diabetic, Taking statins
83	F	Spontaneous	15	J&J Depuy Sigma	Taking statins, had infection
65	F	Getting up from sitting	13	Zimmer Natural Knee	Had e-tear
80	F	Fall	0	Zimmer Natural Knee	Stage 3 CKD
67	F	Getting up from sitting	1	Zimmer Natural Knee	Diabetic, Taking statins
60	F	Fall	23	Zimmer Natural Knee	Taking statins
86	M	Spontaneous	6	J&J Depuy Sigma	Diabetic, Taking statins
75	F	Getting up from sitting	2	Zimmer Natural Knee	Taking statins
64	F	Fall	41	J&J Depuy Sigma	Diabetic, Taking statins, had DVT
53	F	Progressive	n/a	J&J Depuy Sigma	Taking statins
53	M	Fall	n/a	J&J Depuy Sigma	Had infection
68	F	During TKA	112	J&J Depuy Sigma	Had re-tear, Had infection
78	M	Playing sports	43	J&J Depuy Sigma	Had re-tear
62	F	Fall	7	J&J Depuy Sigma	-
65	F	Fall	6	Smith & Nephew Genesis II	-
83	M	Spontaneous	0	Unknown	-
60	M	Walking	1	J&J Depuy Sigma	Diabetic, Taking statins
80	F	Progressive	21	J&J Depuy Sigma (Revision)	Taking statins

Eleven injuries occurred due to a fall, four from getting up from a seated position, four were progressive, three had a spontaneous or unknown cause, two during TKA surgery, one occurred while playing sports, and one occurred while walking. Twenty disruptions occurred following primary total knee arthroplasty, five after a revision total knee arthroplasty, and one during a primary TKA. The average time between surgery and tear was 0.8 months (range, 0 to 3.9 months).

**3.2. Surgical Findings**

Twenty-one patients had complete tears and five had partial tears. Of the complete tears, eleven were avulsions off the tubercle, five from the patella, four were midsubstance

tears, and one was a midsubstance and patella combined tear. Of the five partial tears, three were midsubstance tears and two from the tubercle (Table 2). Five patients underwent concomitant operations; two patients had a polyethylene liner exchange, one had a contralateral TKA, one had a knee arthroscopy revision, and one had a revision of the femoral component.

Twenty-one PT ruptures were repaired by two main methods, suture anchor (A)(N=9) and transosseous (TO)(N=12) repair with allografts being used in four and five patients respectively. The types of allografts used are listed in Table 2. The remaining 5 patients underwent other types of surgeries detailed in Table 2.

**Table 2. Complete partial tears of 21 patients.**

Age	Gender	Partial/Complete tear	Location of Tear	Type of Repair	Retear	Notes
59	F	Complete	Tibial Tubercle	Suture Anchor	Y	-
88	M	Complete	Tibial Tubercle	Suture Anchor	Y	-
67	F	Complete	Tibial Tubercle	Suture Anchor	N	-
80	F	Complete	Inferior pole of Patella	Suture Anchor	Y	-
57	F	Complete	Tibial Tubercle	Suture Anchor	N	Concomitant Knee arthroscopy revision
81	F	Complete	Tibial Tubercle	Suture Anchor	N	-
64	F	Complete	Tibial Tubercle	Suture Anchor	Y	Polyethylene liner exchange
81	M	Partial	Tibial Tubercle	Suture Anchor	N	-
53	F	Complete	Tibial Tubercle	Suture Anchor	N	-
83	F	Complete	Inferior pole of Patella	Transosseous	N	-
65	F	Complete	Midsubstance and patella combined tear	Transosseous	Y	-
80	F	Complete	Inferior pole of Patella	Transosseous	N	-
67	F	Partial	Midsubstance	Transosseous	N	-

(Table 2) *contd....*

Age	Gender	Partial/ Complete tear	Location of Tear	Type of Repair	Retear	Notes
60	F	Partial	Midsubstance	Transosseous	N	Revision of femoral component
86	M	Complete	Inferior pole of Patella	Transosseous	Y	-
75	F	Complete	Midsubstance	Transosseous	N	Polyethylene exchange
64	F	Complete	Inferior pole of Patella	Transosseous	Y	-
53	F	Complete	Tibial Tubercle	Transosseous	N	Ipsilateral TKA
53	M	Complete	Tibial Tubercle	Transosseous	N	-
68	F	Complete	Tibial Tubercle	Transosseous	N	-
78	M	Complete	Midsubstance	Transosseous	N	-
62	F	Partial	Midsubstance	End to End	N	-
65	F	Complete	Tibial Tubercle	End to End	N	-
83	M	Partial	Midsubstance	End to End	N	-
60	M	Complete	Midsubstance	End to End	Y	-
80	F	Complete	Tibial Tubercle	Repaired with polypropylene mesh	N	-

### 3.3. Implant Characteristics

In the 26 patients, 14 Sigma (Johnson and Johnson, Warsaw, IN), 10 Natural Knee (Zimmer, Warsaw, IN), 1 Genesis II (Smith and Nephew, London, UK), and 1 unknown system were implanted during the TKA. Following the procedure, the average patella thickness was 17.3mm (range, 14.3-22.4mm).

### 3.4. Clinical Findings

There was a significant difference in time from PT tear to surgery in patients with grafts (42 days) and those without grafts (6 days) ( $p < 0.001$ ). Overall, there was a significant improvement of KSS from 61 to 83 ( $p = 0.023$ ) after PT repair surgery. There was no difference between A and TO in pre- ( $p = 0.38$ ) and post- ( $p = 0.78$ ) operative KSS scores (TO =  $20 \pm 21.6$ , A =  $15.3 \pm 15.3$ ), post-op extension lag (TO =  $9.8 \pm 10.82$ , A =  $10.1 \pm 10.82$ ,  $p = 0.91$ ) or ROM (TO =  $9.69 \pm 7.69$ , A =  $10.22 \pm 7.69$ ,  $p = 0.98$ ). In the A group, there was no difference in pre and post KSS ( $15.3 \pm 14.6$ ,  $p = 0.054$ ) nor in the TO group ( $20.3 \pm 20.6$ ,  $p = 0.06$ ). There was no difference in post-op KSS in graft vs. no graft ( $20.1 \pm 17.6$ ,  $p = 0.46$ ) or extensor lag ( $45.6 \pm 34.2$ ,  $p = 0.72$ ). Five of the TO patients had extensor lags greater than or equal to 30 degrees as compared to just one in the A cohort. The average postoperative flexion for the TO group was 111 degrees (range, 90-120 degrees) and 109 degrees for the A group (range, 70-125), with nine of the twenty-one patients being able to flex their knee 120 degrees or greater.

There were 2 (7.69%) deep vein thromboses and 4 infections (15.38%). Eight patients had re-tears, 4 in the SA group (3 non-allografts, 1 non-allografts), 3 in the TO group (2 non-allograft, 1 allograft) and one in the other group. Compared to A repair, TO repair had 2.39 times odds of re-tear (95% CI: 0.38,15.4;  $p = 0.354$ ), 1.37 times odds of infection (95% CI: 0.074,25.6;  $p = 0.83$ ). Compared to repairs without a graft, graft repairs had 1.90 times odds of re-tear (95% CI: 0.29, 12.19;  $p = 0.49$ ) and 6.3 times odds of infection (95% CI 0.26, 166.7;  $p = 0.25$ ).

## 4. DISCUSSION

The results of this study show that patellar tendon repair following a TKA can cause significant improvement to functional outcomes of patients. Furthermore, there was no significant difference between functional outcomes, ROM, or post-op extension lags when comparing the SA group to the TO group or in patients who received a graft *versus* those who did not receive a graft. There was, however, a significant difference in time from PT tear to surgery in patients with and without grafts, with grafts more often being used in patients who had longer duration between injury and treatment.

The current gold standard for direct repair is with transosseous drill holes in the patella, but suture anchors have gained popularity [1, 3, 6]. Augmentation techniques for patients with poor tissue quality have also been described [3, 9]. Augmentations usually involve hamstring autograft, fresh-frozen or freeze-dried Achilles tendon with a bone block, extensor mechanism allograft, or synthetic meshes [2, 9]. These recommendations are consistent with our findings of non-allograft techniques performed on an average of 6 days following injury *versus* allografts being used in surgeries performed on an average of 42 days post-injury.

Although functional outcomes as a whole improved for patients following the intervention, PT repair continues to have high rates of complications consistent with the current literature. A study by Rand *et al.* [1] reported on the results of patellar tendon repair following TKA found that 10 out of 18 attempted repairs sustained a re-rupture (55.5%). The re-tear rate reported in our study (30.8%), although high, was lower as compared to the findings in the study reported by Rand *et al.* [1]. This is likely due to improvements in surgical technology and techniques over the past three decades. Conversely, the mean extensor lag in patients with successful and unsuccessful repairs was similar (-1 and -22 degrees in Rand *et al.* [1] respectively, compared to -7 and -20 respectively in our study). Schoderbek *et al.* [2] compared pre and post-op KSS scores among patients who had evidence of an extensor mechanism injury who underwent a revision TKA. They found that there was a significant improvement in the KSS score, but improvements still lagged behind patients who had no extensor mechanism injury prior to TKA revision [2]. This supports our

conclusion that surgery can improve functional outcomes in patients with patellar tendon injuries following TKA, but that the results remain inferior to patients without patellar tendon disruptions. Our study reports an infection rate of 15.38% and DVT findings in 7.69% of patients, which is much higher than primary TKA surgery [10]. Furthermore, the failure rate of 30.8%, while lower than other studies, remains high [1, 5]. Eleven of the 26 patients in this study continued to have an extensor lag following surgery, indicating high levels of residual deficits.

Overall, this paper has shown some improvement as compared to past studies, but the outcomes nevertheless remain poor. More research is needed to assess differences in outcomes between transosseous *versus* suture anchor repair and augmentation *versus* no augmentation.

Our study has several limitations. The small number of patients that qualified for inclusion limits significant conclusions, and even more so once they were separated into the TO and A groups. Furthermore, due to the variable nature of patellar tendon injury presentations and surgeon dependent repairs, outcomes may be difficult to reproduce. Outcomes of patellar tendon repairs following TKA have steadily improved though overall success has remained low.

## CONCLUSION

In conclusion, surgical repair of PT tears following TKA leads to significant clinical improvement, regardless of the fixation method or graft use. We found no difference in outcomes between A and TO repairs and or with graft use.

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study was approved by the ethics committee of Tulane School of Medicine, USA.

## HUMAN AND ANIMAL RIGHTS

Not applicable.

## CONSENT FOR PUBLICATION

Not applicable.

## FUNDING

None.

## CONFLICTS OF INTEREST

The authors declare no conflicts of interest, financial or otherwise.

## ACKNOWLEDGEMENTS

Declared none.

## REFERENCES

- [1] Rand JA, Morrey BF, Bryan RS. Patellar tendon rupture after total knee arthroplasty. *Clin Orthop Relat Res* 1989; (244): 233-8. [<http://dx.doi.org/10.1097/00003086-198907000-00023>] [PMID: 2743664]
- [2] Nam D, Abdel MP, Cross MB, *et al.* The management of extensor mechanism complications in total knee arthroplasty. AAOS exhibit selection. *J Bone Joint Surg Am* 2014; 96(6)e47 [<http://dx.doi.org/10.2106/JBJS.M.00949>] [PMID: 24647514]
- [3] Schoderbek RJ Jr, Brown TE, Mulhall KJ, *et al.* Extensor mechanism disruption after total knee arthroplasty. *Clin Orthop Relat Res* 2006; 446: 176-85. [<http://dx.doi.org/10.1097/01.blo.0000218726.06473.26>] [PMID: 16672886]
- [4] Bonnin M, Lustig S, Hutten D. Extensor tendon ruptures after total knee arthroplasty. *Orthop Traumatol Surg Res* 2016; 102(1)(Suppl.): S21-31. [<http://dx.doi.org/10.1016/j.otsr.2015.06.025>] [PMID: 26797000]
- [5] Rosenberg AG. Management of extensor mechanism rupture after TKA. *J Bone Jt Surg - Br* 2012; 94-B(11\_Supple\_A): 116-9.
- [6] Parker DA, Dunbar MJ, Rorabeck CH. Extensor mechanism failure associated with total knee arthroplasty: prevention and management. *J Am Acad Orthop Surg* 2003; 11(4): 238-47. [<http://dx.doi.org/10.5435/00124635-200307000-00003>] [PMID: 12889862]
- [7] Caplan N, Kader DF. Rationale of the knee society clinical rating system. *Classic Papers in Orthopaedics*. 2014; pp. 197-9. [[http://dx.doi.org/10.1007/978-1-4471-5451-8\\_48](http://dx.doi.org/10.1007/978-1-4471-5451-8_48)]
- [8] Insall JN, Dorr LD, Scott RD, Scott WN. Rationale of the Knee Society clinical rating system. *Clin Orthop Relat Res* 1989; (248): 13-4. [<http://dx.doi.org/10.1097/00003086-198911000-00004>] [PMID: 2805470]
- [9] Browne JA, Hanssen AD. Reconstruction of patellar tendon disruption after total knee arthroplasty: Results of a new technique utilizing synthetic mesh. *J Bone Jt Surg - Ser A* 2011; 93(12): 1137-43.
- [10] Chun KC, Kim KM, Chun CH. Infection following total knee arthroplasty. *Knee Surg Relat Res* 2013; 25(3): 93-9. [<http://dx.doi.org/10.5792/ksrr.2013.25.3.93>] [PMID: 24032096]