

A New Strategy for the Synthesis of Nucleosides: One-Pot Enzymatic Transformation of D-Pentoses into Nucleosides

Anatoly I. Miroshnikov¹, Roman S. Esipov¹, Tat'yana I. Muravyova¹, Irina D. Konstantinova¹, Ilja V. Fateev¹ and Igor A. Mikhailopulo^{*2}

¹Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, Moscow B-437, Russian Federation,

²Institute of Bioorganic Chemistry, National Academy of Sciences of Belarus, 220141 Minsk, Acad. Kuprevicha 5/2, Republic of Belarus

Abstract: A possibility of the one-pot synthesis of purine and pyrimidine nucleosides employing pure recombinant ribokinase, phosphopentomutase and nucleoside phosphorylases in a cascade transformation of D-pentoses into nucleosides is demonstrated. Preliminary results of this study point to reliability to develop practical methods for the preparation of a number of biologically important nucleosides.

Keywords: Nucleosides, one-pot transformation of pentoses into nucleosides.

INTRODUCTION

During recent years, a chemo-enzymatic approach to the synthesis of nucleosides attracts continuously growing interest (for recent reviews, see [1]). There are three lines of investigation in this field of research, viz., (i) transglycosylation reaction consisting of the transfer of a pentofuranosyl moiety of commercially available nucleosides or prepared by chemical methods to purine or pyrimidine bases catalyzed by nucleoside phosphorylases (NP's) or *N*-deoxyribosyltransferases (DRT's) [1, 2]; (ii) biochemical (microbial, enzymatic) retro-synthesis of 2'-deoxyribonucleosides [3], and (iii) chemical or chemo-enzymatic synthesis of α -D-pentofuranose 1-phosphates (PF- α 1P) followed by the enzymatic condensation with heterocyclic bases [4]. Transglycosylation reaction was demonstrated to be a very efficient methodology for the synthesis of plenty of analogues of natural nucleosides of biological and medicinal importance; however, some limitations are well documented too [1, 2a,f]. The retro-synthesis can be employed only for the preparation of 2'-deoxy- β -D-nucleosides of natural or modified heterobases. Chemical [4a-d] and chemo-enzymatic syntheses [4e] of PF- α 1P has attracted much attention during recent years; however, the laborious preparation of PF- α 1P is a serious bottleneck of this approach to the synthesis of nucleosides.

Recently, we have suggested a new strategy for the synthesis of nucleosides [5]. In the present communication, we report on the preparation of the recombinant *E. coli* phosphopentomutase (PPM) and preliminary results of one-pot enzymatic transformation of D-ribose or 2-deoxy-D-ribose into nucleosides employing pure recombinant *E. coli*


ribokinase (RK) [D-pentose \rightarrow pentose-5-phosphates (D-PF-5P)], PPM [D-PF-5P \rightarrow α -D-pentofuranose 1-phosphates (α -D-PF-1P)], and nucleoside phosphorylases (NP's) (α -D-PF-1P + heterobase \rightarrow β -D-nucleosides) coupled with the appropriate pyrimidine or purine heterobases (Scheme 1). The preparation of pure recombinant RK as well as uridine (UP), thymidine (TP) and purine nucleoside (PNP) phosphorylases was described by us earlier [5, 6].

We have earlier prepared recombinant *E. coli* RK and investigated its properties [5]. It is noteworthy that the chemical [7a-9] and chemo-enzymatic [4e, 7] transformations of pentoses into 5-phosphates are rather laborious and low yielding. Ribokinase from *E. coli* was employed for the transformation of D-ribose into D-ribofuranose 5-phosphate in the chemical and enzymatic synthesis of nucleoside-5'-triphosphates with deuterium labels on the 3', 4', and 5' carbons through intermediate formation of 5-phospho-D-ribosyl α -1-pyrophosphate (PRPP) [7b]. We have found that under optimum conditions RK catalyzes the phosphorylation of the primary hydroxyl group not only of D-ribose and 2-deoxy-D-ribose, but also of D-arabinose and D-xylose [5]. These data prompted us to consider this reaction as the first stage in a cascade transformation of pentoses into nucleosides.

RESULTS AND DISCUSSION

The stereospecific C5 \rightarrow C1 translocation of phosphate catalyzed by PPM [3b, 10, 11] is a reliable bridge within the strategy under investigation. With this aim in view, we have prepared recombinant PPM and studied its properties. In brief, the PPM gene *DeoB* from *E. coli* was amplified by PCR with synthetic primers Deo-1b and Deo-2. The primers contained restriction sites *NcoI* (5'-end of the gene) and *EcoRI* (3'-end), respectively. PCR products were analyzed by agarose gel electrophoresis. Then, the amplified fragment was digested by restrictases *NcoI* and *EcoRI* and cloned into the corresponding sites of plasmid vector pET-23d

*Address correspondence to these authors at the Institute of Bioorganic Chemistry, National Academy of Sciences of Belarus, 220141 Minsk, Acad. Kuprevicha 5/2, Republic of Belarus; Tel: + 375 17 267 81 48; Fax: +375 17 267 87 61; E-mail: igor_mikhailo@yahoo.de



UP and TP - uridine and thymidine phosphorylase, respectively
 PNP - purine nucleoside phosphorylase
 RK - ribokinase
 PPM - phosphoenolpyruvate mutase
 Base - pyrimidine or purine heterocyclic base

Scheme 1. One-pot transformation of D-pentoses into β -D-nucleosides through intermediate formation of respective α -D-pentofuranose 1-phosphates (α -D-PF-1P).

(Novagen). The resulting plasmid pER-PPM1 was transformed into *E. coli* strain ER2566. Nucleotide sequence of the cloned fragment according to the F. Saenger method revealed no mutations.

The transformants grown at 37°C on the selective agar plates (100 $\mu\text{g}/\text{mL}$ of ampicillin) were used to inoculate an YT-medium containing 100 $\mu\text{g}/\text{mL}$ of ampicillin. When the culture reached a density of 0.7 units at 600 nm, expression of the recombinant enzyme was induced by the addition of 1 mM (final concentration) isopropyl β -D-galactopyranoside (IPTG) and incubation at 37°C was carried out for another 6 h. Cells were harvested by centrifugation at 4,000 rpm for 30 min at 4°C and the expression level in each clone was analyzed by SDS-polyacrylamide gel electrophoresis. Two clones with the highest level of expression of the desired enzyme (up to 50% of the total cellular protein) were selected.

The cell pellet was resuspended in the buffer solution containing 50 mM Tris-HCl, pH 7.7, 4 mM EDTA, 1 mM phenylmethyl-sulphonyl fluoride (PMSF) (*ca.* 7 mL of the buffer per 1 g of cell pellet). The cells were disrupted by sonication for 5-7 min at 20 kHz in ice water. The resulting cell lysate was cleared by centrifugation for 40 min at 15,000 rpm to afford 12 mL of PPM preparation (protein content 230 mg; combined PPM activity 780 units). The specific enzyme activity (*vide infra*) in the cleared lysate was found to be 3.4 $\mu\text{mol}/\text{min}/\text{mg}$ protein. The major portion of PPM was observed in the cleared lysate; presence of PPM in the insoluble pellet may be due to incomplete cell disruption during sonication.

The PPM of clear soluble fraction was further purified with Q-Sepharose HP chromatography to afford 12.5 mL of PPM solution containing 35 mg protein with total 770 units PPM activity (22 $\mu\text{mol}/\text{min}/\text{mg}$ protein). This PPM preparation was employed in experiments on nucleoside synthesis (*vide infra*). The molecular weight of PPM was estimated to be *ca.* 43,000 (*cf.* [10, 11]) by SDS-polyacrylamide gel gradient electrophoresis relative to standard proteins of known molecular weight.


The PPM activity was measured spectrophotometrically by monitoring the formation of uridine resulting from the PPM catalyzed transformation of D-ribose 5-phosphate into α -D-ribofuranose 1-phosphate followed by uridine phosphorylase (UP [6]) (*cf.* [12]). The increase of molar absorptivity at 275 nm by going from the starting uracil to the formed uridine was found to be $\Delta\epsilon = 1800 \text{ M}^{-1}\text{cm}^{-1}$. One unit of PPM activity is defined as the amount of enzyme that transformed 1 μmol uracil into uridine per minute at 23°C in the reaction mixture containing 0.1 M Tris-HCl, 0.1 M MnCl_2 , 0.05 mM glucose 1,6-diphosphate, 7 mM uracil, 10 mM D-ribofuranose 5-phosphate, and an excess of uridine phosphorylase (UP) (3-5 units [6]) (pH 7.5); the reaction was initiated by addition of PPM.

Finally, one-pot enzymatic transformation of D-ribose or 2-deoxy-D-ribose into the pyrimidine and purine nucleosides was studied (Scheme (2)).

First of all, to assess the efficacy of PPM, we have tested the synthesis of inosine (rI) and 1-(β -D-ribofuranosyl) thymine (rT) using D-ribose 5-phosphate and the respective heterobases, hypoxanthine or thymine, as substrates and PPM and the relevant enzymes, PPM/PNP or PPM/TP, as biocatalysts. Based on the literature data [10], the following reaction conditions have been employed in these experiments after a number of preliminary tests: the molar ratio of D-ribose 5-phosphate and base was 1:1; 3 mM MnCl_2 , 10 mM Tris, pH 7.5, all experiments have been performed at 20°C.

The formation of inosine or rT was analyzed by HPLC after 1 and 24 h. It was found that 35% of hypoxanthine is transformed into inosine after 1 h and its quantity only slightly enhanced after 24 h (39%); under similar time intervals, the yield of rT was 8.5 and 16.9%, respectively.

Upon moving to the one-pot synthesis of nucleosides, we noted rather essential differences between that the optimal reaction conditions for RK, PPM and recombinant nucleoside phosphorylases prepared by us earlier [6]. Bearing this in mind, we have optimized the one-pot reaction conditions aiming at the finding out a compromised



Scheme 2. One-pot transformation of D-ribose and 2-deoxy-D-ribose into pyrimidine and purine nucleosides through intermediate formation of respective α -D-pentofuranose 1-phosphates (α -D-PF-1P).

composition of the components allowing satisfactory function of the enzymes under investigation. We have previously shown that magnesium and manganese ions render rather similar effect on the recombinant RK activity and in the present study the latter was employed for the activation of RK and PPM. The data on the concerted action of the aforementioned enzymes in one-pot transformation of D-ribose and 2-deoxy-D-ribose into pyrimidine and purine nucleosides are shown in Table 1.

It is remarkable that (i) the formation of inosine proceeds faster vs that of 2'-deoxyinosine and reached maximum yield

after 30 min (Table 1; Fig. (1)), and (ii) the synthesis of 2'-deoxyribonucleosides of purines by the transglycosylation reaction proceeds with higher efficiency compared to that of ribonucleosides [13].

The formation of thymidine proceeds with higher efficacy and gives rise to the higher final yield of thymidine vs that of rT (Table 1; Fig. (2)). The observed differences in the formation of thymidine and rT may be explained by the lower substrate activity of α -D-ribofuranose 1-phosphate vs its 2-deoxy-counterpart for thymidine phosphorylase (TP). In harmony with this suggestion is the formation of

Table 1. Progress of Nucleoside Syntheses in the Cascade One-Pot Enzymatic Reactions at 20°C [Content of the Corresponding Nucleoside (%) in the Reaction Mixture vs Time of Reaction^{a,b}]

Time of Reaction, h	Inosine (rI)	2'-Deoxy-inosine (dI)	Thymidine (dT)/2'-Deoxyuridine (dU) ^c	1-(β -D-Ribofuranosyl)-thymine (rT)/Uridine(rU) ^c
0.5	45.9	18.8	14.5/0.9	4.7/27.6
1	46.1	27.3	17.6/1.1	8.5/26.6
24	38.4	38.3	-	-
44	-	-	34.7/33.2	19.9/17.5
96	29.4	34.4	-	-

^aAll the reactions have been run under standard conditions. ^bStandard Reaction Conditions: total volume of the reaction mixture 2 mL; 2 mM ATP, 50 mM KCl, 3 mM MnCl₂, 20 mM Tris-HCl (pH 7.5), 2 mM pentose, 2 mM heterobase; run at 20°C; enzymes (respective units): ribokinase 7.65; phosphopentomutase 3.9; thymidine phosphorylase 4.5; uridine phosphorylase 5.4; purine nucleoside phosphorylase 4.68. ^cHPLC Analyses: Breeze chromatograph (Waters, USA); column: Nova-Pak C18, 4 μ m, 4.6 \times 150 mm; isocratic elution with 1.4 % acetonitrile and 0.1 % TFA at a flow rate of 1 mL/min, run time 15 min; UV-detector, eluates were monitored at 254 nm; retention times, R_t (min): hypoxanthine – 2.7; inosine – 5.4; 2'-deoxyinosine – 6.5; thymine – 3.7; thymidine – 8.3; 1-(β -D-ribofuranosyl)thymine – 5.4; uracil – 2.3; uridine – 3.1; 2'-deoxyuridine – 4.2. All the reference nucleosides have been prepared earlier or from commercial sources. ^cThymidine (TP) and uridine (UP) phosphorylases have been employed for the synthesis of thymine and uracil nucleosides, respectively.




Fig. (1). Progress of the synthesis of hypoxanthine nucleosides in the cascade one-pot enzymatic reactions from the relevant pentoses.

corresponding uracil nucleosides catalyzed by uridine phosphorylase (UP). Indeed, as might be expected, α -D-ribofuranose 1-phosphate manifests much higher substrate activity towards UP vs 2-deoxy- α -D-ribofuranose 1-phosphate. However, the final concentration of uridine is ca. half of that of 2'-deoxyuridine. Interplay of a number of factors contributing to the reaction pathway may be responsible for this result.




Fig. (2). Progress of the synthesis of pyrimidine nucleosides in the cascade one-pot enzymatic reactions from the relevant pentoses.

In conclusion, a possibility of the one-pot synthesis of purine and pyrimidine nucleosides employing pure recombinant ribokinase, phosphopentomutase and nucleoside phosphorylases in a cascade transformation of D-pentoses into nucleosides was demonstrated. Taking into account the functional peculiarities of the enzymes under consideration, a careful optimization of the reaction conditions is necessary to achieve high yield of the desired nucleosides. Note that all the reactions of the present study have been run at an 1:1 molar ratio of pentose and heterobase and at 20°C. The studies directed towards assessment of the scope and limitations of this strategy are now in progress. Preliminary results of this study point to reliability to develop practical methods for the preparation of a number of biologically important nucleosides.

ACKNOWLEDGEMENTS

Financial support by the International Science and Technology Centre (project #B-1640) and Foundations for basic research of Russia and Belarus is gratefully

acknowledged. IAM is thankful to the Alexander von Humboldt-Stiftung (Bonn – Bad-Godesberg, Germany) for partial financial support of this study.

REFERENCES

- [1] a) Mikhailopulo, I.A. Biotechnology of nucleic acid constituents - State of the art and perspectives. *Curr. Org. Chem.*, **2007**, *11*, 307-335; b) Lewkowics, E.S.; Iribarren, A.M. Nucleoside phosphorylases. *Curr. Org. Chem.*, **2006**, *10*, 1197-1215.
- [2] a) Roivainen, J.; Elizarova, T.; Lapinjoki, S.; Mikhailopulo, I.A.; Esipov, R.S.; Miroshnikov, A.I. An enzymatic transglycosylation of purine bases. *Nucleosides Nucleotides Nucleic Acids*, **2007**, *26*, 905-909; b) Barai, V.N.; Zinchenko, A.I.; Eroshevskaya, L.A.; Zhernosek, E.V.; De Clercq, E.; Mikhailopulo, I.A. Chemo-enzymatic synthesis of 3-deoxy- β -D-ribofuranosyl purines. *Helv. Chim. Acta*, **2002**, *85*, 1893-1900; c) Barai, V.N.; Zinchenko, A.I.; Eroshevskaya, L.A.; Kalinichenko, E.N.; Kulak, T.I.; Mikhailopulo, I.A. A universal biocatalyst for the preparation of base- and sugar-modified nucleosides via an enzymatic transglycosylation. *Helv. Chim. Acta*, **2002**, *85*, 1901-1908; d) Kaminski, P.A.; Dacher, P.; Dugue, L.; Pochet, S. *In vivo* reshaping the catalytic site of nucleoside 2'-deoxyribosyltransferase for dideoxy- and didehydronucleosides via a single amino acid substitution. *J. Biol. Chem.*, **2008**, *283*, 20053-20059; e) Anand, R.; Kaminski, P.A.; Ealick, S.E. Structures of purine 2'-deoxyribosyltransferase, substrate complexes, and the ribosylated enzyme intermediate at 2.0 Å resolution. *Biochemistry*, **2004**, *43*, 2384-2393.
- [3] a) Ouwerkerk, N.; van Boom, J.H.; Lugtenburg, J.; Raap, J. Chemo-enzymatic synthesis of thymidine 13C-labelled in the 2'-deoxyribose moiety. *Eur. J. Org. Chem.*, **2000**, 861-866; b) Ouwerkerk, N.; Steenweg, M.; De Ruijter, M.; Brouwer, J.; van Boom, J.H.; Lugtenburg, J.; Raap, J. One-pot two-step enzymatic coupling of pyrimidine bases to 2-deoxy-D-ribose-5-phosphate. A new strategy in the synthesis of stable isotope labeled deoxynucleosides. *J. Org. Chem.*, **2002**, *67*, 1480-1489; c) Ogawa, J.; Saito, K.; Sakai, T.; Horinouchi, N.; Kawano, T.; Matsumoto, S.; Sasaki, M.; Mikami, Y.; Shimizu, S. Microbial production of 2-deoxyribose 5-phosphate from acetaldehyde and triosephosphate for the synthesis of 2'-deoxyribonucleosides. *Biosci. Biotechnol. Biochem.*, **2003**, *67*, 933-936; d) Ishige, T.; Honda, K.; Shimizu, S. Whole organism biocatalysis. *Curr. Opin. Chem. Biol.*, **2005**, *9*, 174-180; e) Horinouchi, N.; Ogawa, J.; Kawano, T.; Sakai, T.; Saito, K.; Matsumoto, S.; Sasaki, M.; Mikami, Y.; Shimizu, S. Biochemical retrosynthesis of 2'-deoxyribonucleosides from glucose, acetaldehyde, and a nucleobase. *Appl. Microbiol. Biotechnol.*, **2006**, *71*, 615-621; f) Horinouchi, N.; Ogawa, J.; Kawano, T.; Sakai, T.; Saito, K.; Matsumoto, S.; Sasaki, M.; Mikami, Y.; Shimizu, S. Efficient production of 2-deoxyribose 5-phosphate from glucose and acetaldehyde by coupling of the alcoholic fermentation system of baker's yeast and deoxyriboaldolase-expressing *Escherichia coli*. *Biosci. Biotechnol. Biochem.*, **2006**, *70*, 1371-1378; g) Horinouchi, N.; Ogawa, J.; Kawano, T.; Sakai, T.; Saito, K.; Matsumoto, S.; Sasaki, M.; Mikami, Y.; Shimizu, S. One-pot microbial synthesis of 2'-deoxyribonucleoside from glucose, acetaldehyde, and a nucleobase. *Biotechnol. Lett.*, **2006**, *28*, 877-881.
- [4] a) Komatsu, H.; Awano, H. First stereoselective synthesis of 2-deoxy- α -D-ribose-1-phosphate: Novel application of crystallization-induced asymmetric transformation. *J. Org. Chem.*, **2002**, *67*, 5419-5421; b) Komatsu, H.; Ikeda, I. Synthesis of 2-deoxy- β -D-ribose 1-phosphate, NMR comparison and its enzymatic activity for structural elucidation of synthetic α -isomer. *Nucleosides Nucleotides Nucleic Acids*, **2003**, *22*, 1685-1686; c) Komatsu, H.; Araki, T. Chemo-enzymatic synthesis of 2',3'-dideoxy-3'-fluoro- β -D-guanosine via 2,3-dideoxy-3-fluoro- α -D-ribose 1-phosphate. *Tetrahedron Lett.*, **2003**, *44*, 2899-2901; d) Komatsu, H.; Awano, H.; Ishibashi, H.; Oikawa, I.; Araki, T. Chemo-enzymatic syntheses of natural and unnatural 2'-deoxynucleosides. *Nucleic Acids Res. Suppl. No. 3*, **2003**, 101-102; e) Taverna-Porro, M.; Bouvier, L.A.; Pereira, C.A.; Montserrat, J.M.; Iribarren, A.M. Chemoenzymatic preparation of nucleosides from furanoses. *Tetrahedron Lett.*, **2008**, *49*, 2642-2645.
- [5] Chuvikovskiy, D.V.; Esipov, R.S.; Skoblov, Y.S.; Chupova, L.A.; Muravyova, T.I.; Miroshnikov, A.I.; Lapinjoki, S.; Mikhailopulo,

- I.A. Ribokinase from *E. coli*: Expression, purification, and substrate specificity. *Bioorg. Med. Chem.*, **2006**, *14*, 6327-6332.
- [6] Esipov, R.S.; Gurevich, A.I.; Chuvikovsky, D.V.; Chupova, L.A.; Muravyova, T.I.; Miroshnikov, A.I. Overexpression of *Escherichia coli* genes encoding nucleoside phosphorylases in the pET/B121(DE3) system yields active recombinant enzymes. *Protein Express. Purif.*, **2002**, *24*, 56-60.
- [7] a) Gross, A.; Abril, O.; Lewis, J.M.; Geresh, S.; Whitesides, G.M. Practical synthesis of 5-phospho-D-ribosyl α -1-pyrophosphate (PRPP): Enzymatic routes from ribose 5-phosphate or ribose. *J. Am. Chem. Soc.*, **1983**, *105*, 7428-7434, and references cited therein; b) Tolbert, T.J.; Williamson, J.R. Preparation of specifically deuterated RNA for NMR studies using a combination of chemical and enzymatic synthesis. *J. Am. Chem. Soc.*, **1996**, *118*, 7929-7940.
- [8] Graham, S.M.; Pope, S.C. Selective phosphitylation of the primary hydroxyl group in unprotected carbohydrates and nucleosides. *Org. Lett.*, **1999**, *1*, 733-736, and references cited therein.
- [9] Ahmadibeni, Y.; Parang, K. Solid-phase reagents for selective monophosphorylation of carbohydrates and nucleosides. *J. Org. Chem.*, **2005**, *70*, 1100-1103.
- [10] Hammer-Jespersen, K.; Munch-Petersen, A. Phosphodeoxyribomutase from *Escherichia coli*. Purification and some properties. *Eur. J. Biochem.*, **1970**, *17*, 397-407; Leer, J.C.; Hammer-Jespersen, K. Multiple forms of phosphodeoxyribomutase from *Escherichia coli*. Physical and chemical characterization. *Biochemistry*, **1975**, *14*, 599-607.
- [11] Rashid, N.; Imanaka, H.; Fukui, T.; Atomi, H.; Imanaka, T. Presence of a novel phosphopentomutase and a 2-deoxyribose 5-phosphate aldolase reveals a metabolic link between pentoses and central carbon metabolism in the hyperthermophilic archaeon *Thermococcus kodakaraensis*. *J. Bacteriol.*, **2004**, *186*, 4185-4191.
- [12] a) Tozzi, M.G.; Catalani, R.; Ippata, P.L.; Mura, U. A coupled optical enzyme assay for phosphopentomutase. *Anal. Biochem.*, **1982**, *123*, 265-269; b) Barbas, III, C.F.; Wong, C.-H. Overexpression and substrate specificity studies of phosphodeoxyribomutase and thymidine phosphorylase. *Bioorg. Chem.*, **1991**, *19*, 261-269.
- [13] Mikhailopulo, I.A.; Zinchenko, A.I.; Bokut, S.B.; Dudchik, N.V.; Barai, V.N.; Kalinichenko, E.N.; Rosemeyer, H.; Seela, F. 1-deaza and 3-deazapurines in the reaction of microbiological transglycosylation. *Biotechnol. Lett.*, **1992**, *14*, 885-890.

Received: January 23, 2010

Revised: July 22, 2010

Accepted: July 23, 2010

© Miroshnikov et al.; Licensee Bentham Open.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.