

P-19**Comparisons of Antibacterial Activity of Crude Extracts Between Two Varieties of *Ficus Deltoidea* Blume**

Hasimah Alimon^{1,*}, Siti Noriswana Safiai¹, Nor Nafizah Mohd Noor¹, Norhayati Daud¹ and Saripah Salbiah Syed Abdul Azziz²

¹Department of Biology; ²Department of Chemistry, Faculty of Science and Mathematics, Universiti Pendidikan Sultan Idris, 35900 Tanjong Malim, Perak, Malaysia; E-mail for correspondence: hasimah@fsmu.upsu.edu.my

A study of antibacterial activity was carried out to compare the effectiveness of two varieties of *F. deltoidea*; var. *bilobata* and var. *angustifolia* as antibacterial agents against three bacteria; *P.aureginosa*, *S.aureus*, and *B.subtilis* using methanol and aqueous solvents. The results show both methanol and aqueous leaf extracts of both varieties were able to inhibit the growth of all three bacteria with minimal inhibition concentration of 25 mgmL⁻¹. The stem aqueous extracts of both varieties also showed inhibition zones against two bacteria, *P.aureginosa* and *S.aureus* at minimal inhibition concentration of 100 mgmL⁻¹ but no inhibition zone observed from methanol stem extracts of both varieties.

Keywords: *F. deltoidea*, var. *bilobata*, var. *angustifolia*, antibacterial activities, inhibition zones.
