



The Open Plant Science Journal

Content list available at: www.benthamopen.com/TOPSJ/

DOI: 10.2174/1874294701710010055



RESEARCH ARTICLE

Expression and Construction of Yeast Expression Vector Containing *CsTCTP1* Gene from Cucumber in Yeast Two-Hybrid System

Qiu-Min Chen¹, Na Cui^{1,2,*}, Yang Yu¹, Xiang-Nan Meng¹ and Hai-Yan Fan^{1,2,*}¹College of Bioscience and Biotechnology, Shenyang Agricultural University, Shenyang 110866, PR China²Key Laboratory of Protected Horticulture of Ministry of Education, Shenyang Agricultural University, Shenyang 110866, PR China

Received: December 06, 2016

Revised: January 24, 2017

Accepted: March 21, 2017

Abstract:

Background:

The translationally controlled tumor protein (TCTP) was originally found in tumor tissue, and later found in other tissues. Initially, TCTP was considered a kind of growth-associated protein. Recent studies have shown that TCTP has many biological functions.

Objective:

To verification of *CsTCTP1* gene function by yeast two-hybrid system, the pGBKT7- *CsTCTP1* yeast expression vector was constructed and cytotoxicity and self-activating activity were detected, which could lay the foundation for further studies on gene function and make a preparation for verification of *CsTCTP1* gene function by yeast two-hybrid system.

Method:

Specific PCR, conventional sequencing, heat shock conversion method and TE/LiAC transformation method.

Results:

We constructed a yeast expression vector containing the *CsTCTP1* gene. The *CsTCTP1* coding sequence was inserted into a pGBKT7 vector as a bait protein and then transformed into the Y2HGold yeast strain.

Conclusion:

We found that *CsTCTP1* protein had no cytotoxic effect and could not be self-activated. The constructed bait expression vector can be used in the subsequent yeast two - hybrid detection system.

Keywords: TCTP, PGBKT7 bait vector, Yeast two-hybrid, Y2HGold, Cytotoxic, Self-activated.

INTRODUCTION

The translationally controlled tumor protein (TCTP) is a highly conserved protein that is abundant in plant and animal cells and has different functions. The TCTP was initially identified in mouse tumor cells [1, 2]. Studies indicated that TCTP has a variety of biological functions: calcium binding protein (CBP) [3], histamine release factor [4], tubulin binding protein [5], anti-apoptotic protein [6, 7], anti-oxidation [8], involved in stress response and signal transduction [9, 10] and so on.

* Address correspondence to this authors at the College of Bioscience and Biotechnology, Shenyang Agricultural University, Shenyang 110866, PR China Tel: +86 2488487163, Fax: +86 2488487163; E-mails: syaua@163.com; hyfan74@163.com

In plants, research about TCTP protein function is still scarce. However, the similarity of TCTP structure of plants and animals suggested that they may have similar functions. Plant TCTP proteins play key roles in both abiotic and biotic stress. It was reported that TCTP transcript or protein levels responded to different stress conditions like drought, aphids, cold, high temperature, salt, light, water deficit and so on [11 - 16].

Powdery mildew, caused by *Sphaerotheca fuliginea* (Sf), is a widely distributed and destructive disease of greenhouse and field-grown cucumber plants and causes great yield loss. In a previous study, TCTP showed increased abundance in powdery mildew resistant line B21-a-2-1-2 (highly resistant) by 2-DE [17].

Protein - protein interactions is an important way to reveal the new role of the proteins. Yeast two-hybrid is a common method for protein interaction screening. Y2HGold yeast stain is a common stain that was used in Yeast two-hybrid system, Y2HGold was constructed of four reporter genes--ADE2, AUR1-C, HIS3 and MEL1 in GAL4 UASs (GAL4 upstream activating sequences) and downstream of the promoter through genetic engineering. GAL4 can specially bind with promoter and activate the downstream reporter gene expression. In this study, the coding sequence of *CsTCTP1* was amplified. The PCR product was cloned into the pGBKT7 vector to generate pGBKT7-*CsTCTP1*, and then the bait vector was transformed into Y2HGold yeast stain. Cytotoxicity and self-activation were detected. On this basis, we can further screen proteins interacting with cucumber TCTP protein by yeast two-hybrid system, building networks of protein interactions, for understanding its role in the cell.

MATERIALS AND METHODS

Plant Materials Handling and Growth Conditions

Cucumber seedlings were Jinyan 4th. Cucumber seeds were sterilized with 50% NaClO 15min and 75% ethyl alcohol for 45s. The sterilized seeds were germinated on a solid MS medium in a growth chamber at 24°C with light intensity of 40 lux. Seedlings with two true leaves were carefully transferred into Soil matrix, and then grown in a greenhouse.

Specific Primer Design and Amplification of *CsTCTP1* Gene Sequence

Primer pairs were designed using Primer 5.0 software and synthesized by Shanghai Biological Engineering Co., Ltd. The primers used in this study are listed in Table 1.

Table 1. *CsTCTP1* gene primer sequences, (5' to 3'). Represented restriction sites are underlined.

Gene name	Primer sequence (5' to 3')
<i>CsTCTP1-F</i>	CGA <u>AATTC</u> ATGTTGGTTTATCAGGACC
<i>CsTCTP1-R</i>	CGGATCCTCAGCACTTGACTTC

Total RNA was isolated from the cucumber leaves using RNA plant plus kit (Tiangen). Complementary DNA (cDNA) was synthesized using oliego (dT) primers (Takara) and PrimeScriptTM RTase reverse transcriptase (Takara) for 1 h at 42°C. Resulting cDNA were used for amplified *CsTCTP1* coding sequence. The PCR product was purified using DNA Gel Extraction kit.

The Bait Vector Construction and Transformation

To obtain the pGBKT7-*CsTCTP1* bait vector, the pMD18-T-*CsTCTP1* vector was digested with *EcoRI* and *BamHI*, the pGBKT7 plasmid was also digested with *EcoRI* and *BamHI*. Then, the connection of *CsTCTP1* and pGBKT7 was executed using T₄ DNA ligase.

Transformation of Y2HGold and Self-Activation Detection

The TE/LiAC method [18] was used for preparing Y2HGold competent yeast cells, and the recombinant plasmid was transformed into Y2HGold Yeast stain through Lithium acetate transformation method. The conversion solution was incubated in SD/-Trp, SD/-Trp-Leu and SD/-Trp-His-Ade medium respectively. The positive clones were detected by colony PCR.

RESULTS

Amplification of *CsTCTP1*

CsTCTP1 gene PCR amplification products and cycle products were detected by 1% agarose gel electrophoresis which showed that the specific fragment was about 500 bp, consistent with that expected (Fig. 1).




Fig. (1). Electrophoresis profile of PCR products of *CsTCTP1* gene M: DNA marker DL2000; 1: Water; 2, 3: *CsTCTP1* PCR amplified product.




Fig. (2). Colony PCR of recombinant plasmid *pGBKT7-CsTCTP1* M: DNA marker DL2000; 1: Water; 2-12: Colony PCR products of *CsTCTP1* gene.

Construction of Bait Expression Vector

To construct the recombinant bait vector, we recycle the small fragments of *pMD18-T-CsTCTP1* and large fragments of *pGBKT7* plasmid after double digest with *EcoR* I and *BamH* I. The colony PCR products of recombinant

plasmid pGBKT7-*CsTCTP1* via agarose gel electrophoresis analysis showed 507 bp of the target gene band (Fig. 2). Sequencing results showed that the gene fragments was consistent with the target gene sequence in GenBank (Fig. 3).

CsTCTP1.seqATGTTGGTTTATCAGGACCTCGTCTCTGGTGACGAGCTTCT	41
Sequeineing	TCAGAGGAGGACCTGCATATGGCCATGGAGCCGAATTCATGTTGGTTTATCAGGACCTCGTCTCTGGTGACGAGCTTCT	560
Consensus	atgttggtttatcaggacctcgtctctggtgacgagcttct	
CsTCTP1.seq	CTCGGATTCGTTTCATACAAGGAAATTGAGAATGGAATGATTTGGGAAGTTGAAGGAAAGTGGGTCGTTAAAGGAGCAG	121
Sequeineing	CTCGGATTCGTTTCATACAAGGAAATTGAGAATGGAATGATTTGGGAAGTTGAAGGAAAGTGGGTCGTTAAAGGAGCAG	640
Consensus	ctcggattcgtttccatacaaggaattgagaatggaatgatttgggaagttgaaggaaagtgggtcgttaaaggagcag	
CsTCTP1.seq	TTGATGTGGATATCGGTGCTAATCCTTCAGCTGAAGGTGGCGGTGATGATGAAGGTGTTGATGATCAGGCTGTGAAGGTG	201
Sequeineing	TTGATGTGGATATCGGTGCTAATCCTTCAGCTGAAGGTGGCGGTGATGATGAAGGTGTTGATGATCAGGCTGTGAAGGTG	720
Consensus	ttgatgtggatatacgggtgctaataccttcagctgaaggtggcggtgatgatgaaggtgttgatgatcaggctgtgaaggtg	
CsTCTP1.seq	GTCGACATTGTTGACACCTTCCGTCCTTCAGGAGCAACCTTCCATGGACAAGAAGGTTTTCCTCACATGCATTAAAGTA	281
Sequeineing	GTCGACATTGTTGACACCTTCCGTCCTTCAGGAGCAACCTTCCATGGACAAGAAGGTTTTCCTCACATGCATTAAAGTA	800
Consensus	gtcgcacattgttgacaccttccgtcttcaggagcaaccttccatggacaagaaggttttcctcacatgcattaagaagta	
CsTCTP1.seq	TATCAAAAAATTGACACCTTTGCTGAAAGGAGAGCAACAAGAGGCATTTAAGAGTAAAAATTGAGGGAGCAGTTAAGTTC	361
Sequeineing	TATCAAAAAATTGACACCTTTGCTGAAAGGAGAGCAACAAGAGGCATTTAAGAGTAAAAATTGAGGGAGCAGTTAAGTTC	880
Consensus	tatcaaaaaattgacacctttgctgaaaggagagcaacaagaggcattttaagagtaaaaattgagggagcagttaagttcc	
CsTCTP1.seq	TACTTCCAAGGTC AAGGACTTGCAATTCITCGTGGGGAGAGTATGGCCGATGACAGCGCTATGGTATTTGCATACTAC	441
Sequeineing	TACTTCCAAGGTC AAGGACTTGCAATTCITCGTGGGGAGAGTATGGCCGATGACAGCGCTATGGTATTTGCATACTAC	960
Consensus	tacttccaaggtaaggacttgcaattcttctggtggggagagtatggccgatgacagcgctatggtatgttgcatactac	
CsTCTP1.seq	AAGGAAGGAGCTACTGACCCAACATTTTATACATTGCCCTGGTTTGAAGGAAGTCAAGTGCTGA.....	507
Sequeineing	AAGGAAGGAGCTACTGACCCAACATTTTATACATTGCCCTGGTTTGAAGGAAGTCAAGTGCTGAGGATCCGTCGACCT	1040
Consensus	aaggaaggagctactgacccaacatTTTTATACATTGCCCTGGTTTGAAGGAAGTCAAGTGCTGA	

Fig. (3). Sequencing results recombinant vector pGBKT7-*CsTCTP1*.

Cytotoxic and Self-activation Detection

To detect whether *CsTCTP1* protein has cytotoxic or self-activation, we transformed pGBKT7-*CsTCTP1* bait expression vector into Y2HGold yeast stain, and incubated it in SD/-Trp, SD/-Trp-Leu and SD/-Trp-His-Ade medium at 30°C. The results showed that pGBKT7-*CsTCTP1* and pGBKT7 stain could grow on SD/-Trp medium. The yeast growth curve of pGBKT7-*CsTCTP1* transformants was the same as pGBKT7 transformants. Both of transformants could not grow on SD/-Trp-Leu and SD/-Trp-His-Ade medium, which indicated that *CsTCTP1* protein had no cytotoxic and no self-activation (Figs. 4, 5) which could lay the foundation for screening *CsTCTP1* interacting target proteins using the yeast two-hybrid.




Fig. (4). Yeast growth curve of *pGBKT7-CsTCTP1* and pGBKT7 transformants.

DISCUSSION

As previously mentioned, although TCTP has many biological functions, its interaction mechanism is unclear, particularly in plants. TCTP gene function analysis of *Arabidopsis* from the pros and cons, both proved TCTP was related to plant growth, and after the *Drosophila* TCTP gene was transferred to transgenic *Arabidopsis* where TCTP had been knocked out, as it can restore growth [19]. Studies have shown that TCTP could interact with TSC-22 and p53 and affect the stability of these two proteins. Overexpression of TCTP could promote the degradation of TSC-22, thereby inhibiting apoptosis mediated by TSC [20]. Human hTCTP gene could interact with HRheb in vitro and promote hRheb to release GDP. hTCTP can also improve mTORC1 activity in vivo [21]. TCTP can also increase NADPH-dependent reactive oxygen species (ROS), and stimulate the cytoskeleton and cell movement [22].

Brioudes *et al.* conducted a positive and negative identification of the TCTP gene of *Arabidopsis thaliana*. It was proved that TCTP was related to growth, and the TCTP gene of *Drosophila* was transformed into the transgenic *Arabidopsis thaliana* knocked out of TCTP gene, its growth rate can be restored, indicating that TCTP gene has the same functions both in plants and animals [19]. A recent study showed that tobacco TCTP protein could interact with ethylene receptor tobacco histidine kinase1 enhancing plant growth through promotion of cell proliferation. Our study found that cucumber TCTP may have important effect on powdery mildew resistance. We are able to detect whether cucumber TCTP protein could interact with other protein to exercise its functions. It has been found that the constructed bait expression vector has no toxicity and self-activating activity and can be used in the further experiments, which would lay a foundation for further studies of TCTP gene function and protein interaction mechanism in plants and animals.

Protein - protein interactions is an important way to study protein function, yeast two-hybrid protein interaction studies is a common method. In this work, the three-reporter gene was used to verify the cytotoxic effect and self-activation of fusion protein in yeast, which induced the false positive in further yeast two hybrid assays. Our study intends to make CsTCTP1 protein as a bait to screen the interaction proteins, and understand how the CsTCTP1 protein exercises their functions.




Fig. (5). Cytotoxic and self-activation detection. A, B: pGBKT7-*CsTCTP1* and pGBKT7 transformants incubated at SD/-Trp medium. C, D: pGBKT7-*CsTCTP1* and pGBKT7 transformants incubated at SD/-Trp-Leu medium. E, F: pGBKT7-*CsTCTP1* and pGBKT7 transformants incubated at SD/-Trp-His-Ade medium.

CONCLUSION

We have successfully amplified and purified of *CsTCTP1* gene in cucumber coding sequence, and constructed the yeast two-hybrid yeast expression vector. The bait vector was successfully transformed to Y2HGGold, which has no cytotoxic effect and no self-activation. Our study is a preparation for screening *CsTCTP1* interaction proteins by yeast two-hybrid.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

HUMAN AND ANIMAL RIGHTS

No Animals/Humans were used for studies that are base of this research.

CONSENT FOR PUBLICATION

Not applicable.

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

This study was supported by the Public Research Fund of Science Cause of Liaoning Province (2014002013); Science and Technology Research Projects of Education Department of Liaoning Province.

REFERENCES

- [1] Bommer UA, Thiele BJ. The translationally controlled tumour protein (TCTP). *Int J Biochem Cell Biol* 2004; 36(3): 379-85. [J]. [[http://dx.doi.org/10.1016/S1357-2725\(03\)00213-9](http://dx.doi.org/10.1016/S1357-2725(03)00213-9)] [PMID: 14687915]
- [2] Yenofsky R, Cereghini S, Krowczynska A, Brawerman G. Regulation of mRNA utilization in mouse erythroleukemia cells induced to differentiate by exposure to dimethyl sulfoxide. *Mol Cell Biol* 1983; 3(7): 1197-203. [<http://dx.doi.org/10.1128/MCB.3.7.1197>] [PMID: 6577280]
- [3] Xu A, Bellamy AR, Taylor JA. Expression of translationally controlled tumour protein is regulated by calcium at both the transcriptional and post-transcriptional level. *Biochem J* 1999; 342(Pt 3): 683-9. [<http://dx.doi.org/10.1042/bj3420683>] [PMID: 10477280]
- [4] MacDonald SM, Bhisutthibhan J, Shapiro TA, *et al.* Immune mimicry in malaria: Plasmodium falciparum secretes a functional histamine-releasing factor homolog in vitro and in vivo. *Proc Natl Acad Sci USA* 2001; 98(19): 10829-32. [<http://dx.doi.org/10.1073/pnas.201191498>] [PMID: 11535839]
- [5] Gachet Y, Tournier S, Lee M, Lazaris-Karatzas A, Poulton T, Bommer UA. The growth-related, translationally controlled protein P23 has properties of a tubulin binding protein and associates transiently with microtubules during the cell cycle. *J Cell Sci* 1999; 112(Pt 8): 1257-71. [PMID: 10085260]
- [6] Liu H, Peng HW, Cheng YS, Yuan HS, Yang-Yen HF. Stabilization and enhancement of the antiapoptotic activity of mcl-1 by TCTP. *Mol Cell Biol* 2005; 25(8): 3117-26. [<http://dx.doi.org/10.1128/MCB.25.8.3117-3126.2005>] [PMID: 15798198]
- [7] Susini L, Besse S, Duflaut D, *et al.* TCTP protects from apoptotic cell death by antagonizing bax function. *Cell Death Differ* 2008; 15(8): 1211-20. [<http://dx.doi.org/10.1038/cdd.2008.18>] [PMID: 18274553]
- [8] Gnanasekar M, Ramaswamy K. Translationally controlled tumor protein of *Brugia malayi* functions as an antioxidant protein. *Parasitol Res* 2007; 101(6): 1533-40. [<http://dx.doi.org/10.1007/s00436-007-0671-z>] [PMID: 17687568]
- [9] Shen QH, Saijo Y, Mauch S, *et al.* Nuclear activity of MLA immune receptors links isolate-specific and basal disease-resistance responses. *Science* 2007; 315(5815): 1098-103. [<http://dx.doi.org/10.1126/science.1136372>] [PMID: 17185563]
- [10] Gnanasekar M, Dakshinamoorthy G, Ramaswamy K. Translationally controlled tumor protein is a novel heat shock protein with chaperone-like activity. *Biochem Biophys Res Commun* 2009; 386(2): 333-7. [<http://dx.doi.org/10.1016/j.bbrc.2009.06.028>] [PMID: 19523440]
- [11] Kim YM, Han YJ, Hwang OJ, *et al.* Overexpression of Arabidopsis translationally controlled tumor protein gene *AtTCTP* enhances drought tolerance with rapid ABA-induced stomatal closure. *Mol Cells* 2012; 33(6): 617-26. [<http://dx.doi.org/10.1007/s10059-012-0080-8>] [PMID: 22610367]

- [12] Zhang JM, Ma LF, Li W, Zhang J, Li DD, Li XB. Cotton TCTP1 gene encoding a translationally controlled tumor protein participates in plant response and tolerance to aphids. *Plant Cell Tissue Organ Cult* 2014; 117(2): 145-56. [<http://dx.doi.org/10.1007/s11240-014-0426-3>]
- [13] Cao B, Lu Y, Chen G, Lei J. Functional characterization of the translationally controlled tumor protein (TCTP) gene associated with growth and defense response in cabbage. *Plant Cell Tissue Organ Cult* 2010; 103(2): 217-26. [<http://dx.doi.org/10.1007/s11240-010-9769-6>]
- [14] Santa Brigida AB, dos Reis SP, Costa CdeN, Cardoso CM, Lima AM, de Souza CR. Molecular cloning and characterization of a cassava translationally controlled tumor protein gene potentially related to salt stress response. *Mol Biol Rep* 2014; 41(3): 1787-97. [<http://dx.doi.org/10.1007/s11033-014-3028-6>] [PMID: 24413992]
- [15] Sage-Ono K, Ono M, Harada H, Kamada H. Dark-induced accumulation of mRNA for a homolog of translationally controlled tumor protein (TCTP) in Pharbitis. *Plant Cell Physiol* 1998; 39(3): 357-60. [<http://dx.doi.org/10.1093/oxfordjournals.pcp.a029377>] [PMID: 9588028]
- [16] Vincent D, Ergül A, Bohlman MC, *et al*. Proteomic analysis reveals differences between *Vitis vinifera* L. cv. Chardonnay and cv. Cabernet Sauvignon and their responses to water deficit and salinity. *J Exp Bot* 2007; 58(7): 1873-92. [<http://dx.doi.org/10.1093/jxb/erm012>] [PMID: 17443017]
- [17] Fan H, Ren L, Meng X, Song T, Meng K, Yu Y. Proteome-level investigation of *Cucumis sativus*-derived resistance to *Sphaerotheca fuliginea*. *Acta Physiol Plant* 2014; 36(7): 1781-91. [<http://dx.doi.org/10.1007/s11738-014-1552-6>]
- [18] Bartel PL, Fiwlds S. The Yeast Two-Hybrid System. Oxford, New York: Oxford University Press 1997; pp. 425-8.
- [19] Brioudes F, Thierry AM, Chambrier P, Mollereau B, Bendahmane M. Translationally controlled tumor protein is a conserved mitotic growth integrator in animals and plants. *Proc Natl Acad Sci USA* 2010; 107(37): 16384-9. [<http://dx.doi.org/10.1073/pnas.1007926107>] [PMID: 20736351]
- [20] Lee JH, Rho SB, Park SY, Chun T. Interaction between fortilin and transforming growth factor-beta stimulated clone-22 (TSC-22) prevents apoptosis *via* the destabilization of TSC-22. *FEBS Lett* 2008; 582(8): 1210-8. [<http://dx.doi.org/10.1016/j.febslet.2008.01.066>] [PMID: 18325344]
- [21] Dong X, Yang B, Li Y, Zhong C, Ding J. Molecular basis of the acceleration of the GDP-GTP exchange of human ras homolog enriched in brain by human translationally controlled tumor protein. *J Biol Chem* 2009; 284(35): 23754-64. [<http://dx.doi.org/10.1074/jbc.M109.012823>] [PMID: 19570981]
- [22] Jung J, Kim HY, Kim M, Sohn K, Kim M, Lee K. Translationally controlled tumor protein induces human breast epithelial cell transformation through the activation of Src. *Oncogene* 2011; 30(19): 2264-74. [<http://dx.doi.org/10.1038/onc.2010.604>] [PMID: 21278788]