




The Open Dermatology Journal

Supplementary Material

Content list available at: www.benthamopen.com/TODJ/

DOI: 10.2174/187437220181201054



0.1MG/ML Tamoxifen Gel Improves Plaque Psoriasis. An Open Study

Ayman Abdelmaksoud¹, Angela Filoni², Domenico Bonamonte², Giuseppe Giudice³, Azza Abdelaziz⁴, Mohammed Zaki⁵, Maurizio Congedo⁶ and Michelangelo Vestita^{3,*}

¹Mansoura Dermatology, Venerology and Leprology Hospital, Mansoura, Egypt.

²Department of Biomedical Science and Human Oncology University of Bari, Section of Dermatology, 11, Piazza Giulio Cesare, Bari, 70124, Italy

³Department of Emergency and Organ Transplantation, Section of Plastic and Reconstructive Surgery, University of Bari, 11, Piazza Giulio Cesare, Bari, 70124, Italy

⁴Department of pathology, Faculty of Medicine, Mansoura University, Mansoura, Egypt.

⁵Department of pharmacognosy, Faculty of Pharmacy, Alazhar University, Cairo, Egypt.

⁶Section of Dermatology, Vito Fazzi Hospital, Piazza Filippo Muratore, Lecce, 73100, Italy

PREPARATION OF TOPICAL LIPOSOMAL TAMOXIFEN CITRATE GEL (0.1MG/ML)

1 mg of tamoxifen citrate was levigated with 10 ml of the liposomal base (obtained from Flukae Chemika) for 10 minutes, then, left for about 2 hours.

Carboxymethyl cellulose sodium Gel (Na CMC gel) was prepared by suspending 3 mg of Na CMC gel in 100 ml distilled water, and allowing to swell in a refrigerator overnight.

The gel base was added to the liposomal tamoxifen citrate mixture in geometric proportions, with continuous levigation after each addition, to obtain 100 mg of the final formula.