



The Open Orthopaedics Journal

Content list available at: www.benthamopen.com/TOORTHJ/

DOI: 10.2174/1874325001812010396, 2018, 12, 03-00



RESEARCH ARTICLE

Correlations between Capsular Changes and ROM Restriction in Frozen Shoulder Evaluated by Plain MRI and MR Arthrography

Kenji Kanazawa¹, Yoshihiro Hagiwara^{2,*}, Takuya Sekiguchi³, Kazuaki Suzuki⁴, Masashi Koide⁵, Akira Ando⁵ and Yutaka Yabe²

¹Department of Orthopaedic Surgery, South Miyagi Medical Center, Oogawara, Miyagi, Japan

²Department of Orthopaedic Surgery, Tohoku University School of Medicine, Sendai, Miyagi, Japan

³Department of Orthopaedic Surgery, Iwate Prefectural Central Hospital, Morioka, Iwate, Japan

⁴Department of Orthopaedic Surgery, JR Sendai Hospital, Sendai, Miyagi, Japan

⁵Department of Orthopaedic Surgery, Matsuda Hospital, Sendai, Miyagi, Japan

Received: June 8, 2018

Revised: September 13, 2018

Accepted: September 13, 2018

Supplementary Table 1. Correlation coefficient difference between ROM and thickness of CHL, axillary area, capsular area and thickness of capsule in axillary recess.

			FF	LE	ER	AER	AIR	HBB	HF
Thickness of CHL	MRI	R	-0.31	-0.08	-0.24	0.09	0.28	-0.1	0.1
		P	0.139	0.714	0.269	0.681	0.18	0.639	0.65
Axillary area	MRI	R	0.19	0.36	0.25	0.24	-0.21	0.18	0.06
		P	0.385	0.087	0.235	0.255	0.337	0.4	0.768
Thickness of capsule in axillary recess									
Humeral side	MRI	R	0.13	<0.01	-0.2	-0.11	0.09	-0.1	0.12
		P	0.559	0.995	0.339	0.607	0.673	0.625	0.57
	MRA	R	0.01	<-0.01	-0.08	0.01	-0.02	<0.01	0.03
		P	0.954	0.999	0.702	0.973	0.942	0.986	0.872

* Address correspondence to this author at the Department of Orthopaedic Surgery, Tohoku University School of Medicine, 1-1 Seiryomachi, Aoba-ku, Sendai, 980-8574, Japan; Tel: +81-22-717-7245; Fax: +81-22-717-7248; E-mail: hagi@med.tohoku.ac.jp